

Applications of Mathematics

Sadek Gala

Uniqueness of weak solutions of the Navier-Stokes equations

Applications of Mathematics, Vol. 53 (2008), No. 6, 561–582

Persistent URL: <http://dml.cz/dmlcz/140341>

Terms of use:

© Institute of Mathematics AS CR, 2008

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

UNIQUENESS OF WEAK SOLUTIONS OF THE
NAVIER-STOKES EQUATIONS

SADEK GALA, Mostaganem

(Received February 13, 2007, in revised version July 7, 2007)

Abstract. Consider the Navier-Stokes equation with the initial data $a \in L^2_\sigma(\mathbb{R}^d)$. Let u and v be two weak solutions with the same initial value a . If u satisfies the usual energy inequality and if $\nabla v \in L^2((0, T); \dot{X}_1(\mathbb{R}^d)^d)$ where $\dot{X}_1(\mathbb{R}^d)$ is the multiplier space, then we have $u = v$.

Keywords: Navier-Stokes equations, solution uniqueness, weak Leray-Hopf solution, multiplier space

MSC 2010: 35Q30, 76D05, 76D03

1. INTRODUCTION

Consider the Navier-Stokes equation in $(0, T) \times \mathbb{R}^d$ with $0 < T < \infty$ and $d \geq 3$:

$$(1.1) \quad \begin{aligned} \partial_t u + (u \cdot \nabla)u - \Delta u + \nabla p &= 0, & (x, t) \in \mathbb{R}^d \times (0, \infty), \\ \nabla \cdot u &= 0, & (x, t) \in \mathbb{R}^d \times (0, \infty), \\ u(x, 0) &= a(x), & x \in \mathbb{R}^d, \end{aligned}$$

where $u = u(x, t)$ is the velocity field, $p = p(x, t)$ is the scalar pressure and $a(x)$ with $\operatorname{div} a = 0$ in the sense of distributions is the initial velocity field. For simplicity, we assume that the external force has a scalar potential and is included in the pressure gradient.

In their famous paper, Leray [11] and Hopf [5] constructed a weak solution u of (1.1) for arbitrary $a \in L^2_\sigma$. The solution is called the Leray-Hopf weak solution. In the general case the problem of uniqueness of Leray-Hopf's weak solutions is still an open question. Masuda [13] extended Serrin's class for uniqueness of weak

solutions and made it clear that the class $L^\infty((0, T); L^d(\mathbb{R}^d))$ plays an important role for the uniqueness of weak solutions. Kozono-Sohr [7] showed that uniqueness holds in $L^\infty((0, T); L^d)$.

Foias [3] and Serrin [14] introduced the class $L^\alpha((0, \infty); L^q)$ and showed that under the additional assumption

$$u \in L^\alpha((0, \infty); L^q) \quad \text{for } \frac{2}{\alpha} + \frac{d}{q} = 1 \quad \text{with } q > d,$$

u is the only weak solution.

The purpose of this paper is to improve the criterion on uniqueness of weak solutions to the class $L^2((0, T); \dot{X}_1(\mathbb{R}^d)^d)$ (Definition 2). We know that for every $a \in L^2_\sigma(\mathbb{R}^d)$ there is at least one weak solution u of (1.1) satisfying the energy inequality. Here we mean by the weak solution a function $u \in L^\infty((0, T); L^2_\sigma) \cap L^2((0, T); \dot{H}^1_\sigma)$ satisfying (1.1) in the sense of distributions (Definition 4). For more facts concerning uniqueness of weak solutions we refer to the celebrated paper of Kozono and Sohr [7].

1.1. *BMO and Hardy space* $\mathcal{H}^1(\mathbb{R}^d)$

We recall that a locally summable function g on \mathbb{R}^d is said to have bounded mean oscillation if

$$\|g\|_{BMO} = \sup_{x,R} \frac{1}{|B(x,R)|} \int_{B(x,R)} |g(y) - g_{B(x,R)}| \, dy < \infty,$$

where

$$g_{B(x,R)} = \frac{1}{|B(x,R)|} \int_{B(x,R)} g(y) \, dy.$$

The class of functions of bounded mean oscillation is denoted by *BMO* and often is referred to as the John-Nirenberg space.

Note that

$$\|g\|_{BMO} = 0 \quad \text{if and only if } g = \text{const.}$$

It is thus natural to consider the quotient space BMO/\mathbb{R} with the norm induced by $\|\cdot\|_{BMO}$. Then BMO/\mathbb{R} is a Banach space, which will also be denoted *BMO* for simplicity. We easily see that $L^\infty \subset BMO$ with continuous injection. For $f(x) = \log|x|$ we have $f \in BMO$ but $f \notin L^\infty$, so *BMO* is strictly larger than L^∞ .

Next, we recall the definition and some of the main properties of Hardy spaces $\mathcal{H}^p(\mathbb{R}^d)$ introduced by E. Stein and G. Weiss [16] (for more facts on these spaces see C. Fefferman and E. Stein [4]).

Definition 1 ([4]). Let $0 < p < \infty$ and let $\varphi \in \mathcal{S}(\mathbb{R}^d)$ satisfy $\int_{\mathbb{R}^d} \varphi \, dx = 1$. A tempered distribution f belongs to the Hardy space $\mathcal{H}^p(\mathbb{R}^d)$ if

$$(1.2) \quad f^*(x) = \sup_{t>0} |(\varphi_t * f)(x)| \in L^p(\mathbb{R}^d),$$

where $\varphi_t(x) = t^{-d}\varphi(t^{-1}x)$.

It is known that if $f \in \mathcal{H}^p(\mathbb{R}^d)$, then (1.2) holds for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$ satisfying $\int_{\mathbb{R}^d} \varphi \, dx = 1$. The (quasi)-norm of $\mathcal{H}^p(\mathbb{R}^d)$ is defined, up to equivalence, by

$$\|f\|_{\mathcal{H}^p(\mathbb{R}^d)} = \|f^*(x)\|_{L^p(\mathbb{R}^d)} = \left(\int_{\mathbb{R}^d} |f^*(x)|^p \, dx \right)^{1/p}.$$

We know by [4], [15] that if $1 \leq p < \infty$, then \mathcal{H}^p is a Banach space:

$$\begin{aligned} \mathcal{H}^p(\mathbb{R}^d) &= L^p(\mathbb{R}^d) \quad \text{for } 1 < p < \infty, \\ \mathcal{H}^1(\mathbb{R}^d) &\subset L^1(\mathbb{R}^d) \quad \text{with continuous injection,} \end{aligned}$$

and that $\mathcal{H}^p(\mathbb{R}^d)$, $0 < p < 1$, are quasi-Banach spaces in the quasi-norm $\|\cdot\|_{\mathcal{H}^p(\mathbb{R}^d)}$.

The crucial fact for our purpose is the boundedness of the Riesz transforms R_j on all of the spaces \mathcal{H}^p . Furthermore, an L^1 -function f on \mathbb{R}^d belongs to $\mathcal{H}^1(\mathbb{R}^d)$ if and only if its Riesz transforms $R_j f$ all belong to $L^1(\mathbb{R}^d)$ and

$$\|f\|_{\mathcal{H}^1(\mathbb{R}^d)} \cong \|f\|_{L^1(\mathbb{R}^d)} + \sum_{j=1}^d \|R_j f\|_{L^1(\mathbb{R}^d)} \quad (\text{equivalent norms}).$$

Notice that all functions $f \in \mathcal{H}^1(\mathbb{R}^d)$ satisfy

$$(1.3) \quad \int_{\mathbb{R}^d} f(x) \, dx = 0.$$

Indeed, the assumption $f \in \mathcal{H}^1(\mathbb{R}^d)$ implies that the Fourier transforms

$$\hat{f}(\xi) = \int f(x) e^{-ix\xi} \, dx \quad \text{and} \quad \widehat{R_j f}(\xi) = \frac{i\xi_j}{|\xi|} \hat{f}(\xi) \quad (j = 1, \dots, d),$$

are all continuous on \mathbb{R}^d , so $\hat{f}(0) = 0$ and (1.3) is proved.

A fundamental theorem in the theory of Hardy spaces $\mathcal{H}^1(\mathbb{R}^d)$ developed by C. Fefferman and E. Stein [4] asserts:

Theorem 1 (Fefferman). *The dual space of $\mathcal{H}^1(\mathbb{R}^d)$ is BMO. More precisely, L is a continuous linear functional on $\mathcal{H}^1(\mathbb{R}^d)$ if and only if it can be represented as*

$$L(f) = \int_{\mathbb{R}^d} fg$$

for some function g in BMO. Moreover, for any $g \in BMO$ and any $f \in \mathcal{H}^1(\mathbb{R}^d)$ we have

$$(1.4) \quad \left| \int_{\mathbb{R}^d} fg \, dx \right| \leq c(d) \|f\|_{\mathcal{H}^1} \|g\|_{BMO}.$$

Let $\gamma > 1$. We define the maximal function of f depending on γ ,

$$M_\gamma f(x) = \sup_{t>0} \left(\frac{1}{|B_t(x)|} \int_{B_t(x)} |f(y)|^\gamma \, dy \right)^{1/\gamma}.$$

We begin by establishing the following result which is a variant of the Hardy-Littlewood maximal theorem. We need

Lemma 1. *If $\gamma < p \leq \infty$, then*

$$M_\gamma : L^p(\mathbb{R}^d) \rightarrow L^p(\mathbb{R}^d) \quad \text{is bounded.}$$

Proof. See [15]. □

In [1], Coifman, Lions, Meyer and Semmes showed that the Hardy spaces can be used to analyze the regularity of various nonlinear quantities by the compensated compactness theory due to L. Murat [12] and F. Tartar [17]. Since then, these spaces play an important role in studying the regularity of solutions to partial differential equations. In particular, it was shown that for exponents p, q with $1 < p < \infty$, $1/p + 1/q = 1$, and vector fields $u \in L^p(\mathbb{R}^d)^d$, $v \in L^q(\mathbb{R}^d)^d$ with $\operatorname{div} u = 0$, $\operatorname{curl} v = 0$ in the sense of distributions, the scalar product $u \cdot v$ belongs to the Hardy space $\mathcal{H}^1(\mathbb{R}^d)$. Moreover, there exists a positive constant C such that

$$\|u \cdot v\|_{\mathcal{H}^1(\mathbb{R}^d)} \leq C \|u\|_{L^p} \|v\|_{L^q}.$$

The main purpose of this subsection is to prove two facts about the div-curl lemma without imposing any a priori assumptions on exact cancellation, namely, the divergence and curl need not be zero. Our results will lead to $\operatorname{div}(uv)$ being in the Hardy space $\mathcal{H}^1(\mathbb{R}^d)$.

The proof will be divided into two parts. In part 1, we consider the case u and v being supported on the ball $|x| \leq R_0$, where $R_0 > 1$ is a positive constant to be determined later, while in Part 2, the general case follows by partition of unity. In order to simplify the presentation, we take $p = q = 2$.

The Sobolev space $H_p^1(\mathbb{R}^d)$, $1 \leq p < \infty$, consists of functions $f \in L^p(\mathbb{R}^d)$ such that $|\nabla f| \in L^p(\mathbb{R}^d)$. It is a Banach space with respect to the norm

$$\|f\|_{H_p^1} = \|f\|_{L^p} + \|\nabla f\|_{L^p}.$$

Specifically, we will prove

Theorem 2. *Let $u \in H_p^1(\mathbb{R}^d)^d$ and $v \in H_q^1(\mathbb{R}^d)$, $p > 1$, $1/p + 1/q = 1$. Then there exists a positive constant $C(d)$ such that*

$$(1.5) \quad \|\operatorname{div}(uv)\|_{\mathcal{H}^1(\mathbb{R}^d)} \leq C(\|u\|_{L^p} \|\nabla v\|_{L^q} + \|\operatorname{div} u\|_{L^p} \|v\|_{L^q}).$$

Remark 1. Such inequalities and their generalizations are useful in hydrodynamics. The reader is referred, in particular, to [1], [2].

Theorem 2 is a generalized version of the “div-curl” lemma ([1], Theorem II.1). Observe that when $\operatorname{div} u = 0$, Theorem 2 reduces to the classical div-curl lemma [1].

The following result due to [1] shows the importance of the Hardy space theory in estimating the non-linear term $u \cdot \nabla v$ attached to the Navier-Stokes equations. This produces a useful tool for PDE.

Lemma 2. *Let $1 < p < \infty$, $1 < q < d$ and $1/r = 1/p + 1/q < 1/d + 1$. If $u \in L^p(\mathbb{R}^d)^d$ with $\nabla \cdot u = 0$ and $\nabla v \in L^q(\mathbb{R}^d)$, then*

$$u \cdot \nabla v \in \mathcal{H}^r(\mathbb{R}^d)$$

and

$$\|u \cdot \nabla v\|_{\mathcal{H}^r(\mathbb{R}^d)} \leq C\|u\|_{L^p} \|\nabla v\|_{L^q}.$$

Proof. The result is due to [1]; but we give here a detailed proof for the reader’s convenience. Observe that

$$f = u \cdot \nabla v = \nabla \cdot (u \otimes (v - c))$$

for an arbitrary constant vector c . So we get

$$(\varphi_t * f)(x) = t^{-d-1} \int_{B_t(x)} (\nabla \varphi)(t^{-1}(x - y)) u(y) (v(y) - m_B(v)) dy,$$

where

$$m_B(v) = \frac{1}{|B_t(x)|} \int_{B_t(x)} v(y) \, dy.$$

Taking

$$1 < \gamma < \infty, \quad 1 < \beta < d \quad \text{with} \quad \frac{1}{\gamma} + \frac{1}{\beta} = 1 + \frac{1}{d}$$

and writing

$$\frac{1}{\beta^*} = \frac{1}{\beta} - \frac{1}{d},$$

we see by the Poincaré-Sobolev inequality that

$$\begin{aligned} |(\varphi_t * f)(x)| &\leq \frac{C}{t^{d+1}} \left(\int_{B_t(x)} |u(y)|^\gamma \, dy \right)^{1/\gamma} \left(\int_{B_t(x)} |v(y) - m_B(v)|^{\beta^*} \, dy \right)^{1/\beta^*} \\ &\leq \frac{C}{t^{d+1}} \left(\int_{B_t(x)} |u(y)|^\gamma \, dy \right)^{1/\gamma} \left(\int_{B_t(x)} |\nabla v(y)|^\beta \, dy \right)^{1/\beta} \\ &= C \left(\frac{1}{|B_t(x)|} \int_{B_t(x)} |u(y)|^\gamma \, dy \right)^{1/\gamma} \left(\frac{1}{|B_t(x)|} \int_{B_t(x)} |\nabla v(y)|^\beta \, dy \right)^{1/\beta} \\ &\leq C(M_\gamma u)(x) \cdot (M_\beta(\nabla v))(x). \end{aligned}$$

We thus obtain

$$\sup_{t>0} |(\varphi_t * f)(x)| \leq C(M_\gamma u)(x) \cdot (M_\beta(\nabla v))(x).$$

Since we can take γ and β such that

$$1 < \gamma < p, \quad 1 < \beta < q < d,$$

it follows from Lemma 1 that

$$\|M_\gamma u\|_{L^p} \leq C\|u\|_{L^p}, \quad \|M_\beta(\nabla v)\|_{L^q} \leq C\|\nabla v\|_{L^q}.$$

Lemma 2 now follows from Hölder's inequality

$$\|f \cdot g\|_{L^r} \leq \|f\|_{L^p} \|g\|_{L^q} \quad \left(0 < p < \infty, \quad 0 < q < \infty, \quad \frac{1}{r} = \frac{1}{p} + \frac{1}{q} \right).$$

This completes the proof of the lemma. □

We are now in position to prove Theorem 2.

P r o o f of Theorem 2. We distinguish three cases.

Case A. Let us assume first that

$$\nabla \cdot u = 0.$$

In this case we get

$$\operatorname{div}(vu) = (\nabla v) \cdot u + v \operatorname{div} u = u \cdot \nabla v.$$

Then we have $u \in L^p(\mathbb{R}^d)^d$, $\nabla v \in L^q(\mathbb{R}^d)$ with $\operatorname{div} u = 0$, $\operatorname{curl}(\nabla v) = 0$ in the sense of distributions. It follows from Lemma 2 that

$$u \cdot \nabla v \in \mathcal{H}^1(\mathbb{R}^d)$$

and there exists an absolute constant C such that

$$\|\operatorname{div}(vu)\|_{\mathcal{H}^1(\mathbb{R}^d)} \leq C \|u\|_{L^p} \|\nabla v\|_{L^q}.$$

Case B. We may of course introduce an additional assumption that u and v are supported on the ball $|x| \leq R_0$. In order to simplify the presentation, we take $p = q = 2$. We shall write Ω for the ball in \mathbb{R}^d of radius R_0 centered at the origin. By $H_0^1(\Omega)$ we denote the closed subspace of $H^1(\Omega)$ which is the closure of $C_0^\infty(\Omega)$ in the H^1 norm. Let

$$g = \operatorname{div} u \in L^2(\mathbb{R}^d).$$

By the classical result (see e.g. [18]) we know that

$$g = \partial_1 g_1 + \dots + \partial_d g_d,$$

where g_1, \dots, g_d belong to $H_0^1(\Omega)$. Set

$$G = (g_1, \dots, g_d) \quad \text{and} \quad r = u - G.$$

Then it follows that

$$\operatorname{div} r = 0 \quad \text{and} \quad r \in L^2(\Omega).$$

Using Lemma 2 we infer

$$\operatorname{div}(rv) \in \mathcal{H}^1(\mathbb{R}^d).$$

Further we set

$$f = \operatorname{div}(Gv).$$

Using Lemma 3 below, we conclude that $f \in \mathcal{H}^1(\mathbb{R}^d)$.

Case C. The general case. We call φ a smooth bump function with compact support provided

$$1 = \sum_{k \in \mathbb{Z}^d} \varphi^2(x - k).$$

If f and g are two functions, we thus have

$$f(x)g(x) = \sum_{k \in \mathbb{Z}^d} f(x)\varphi^2(x - k)g(x) = \sum_{k \in \mathbb{Z}^d} f_k(x)g_k(x),$$

where

$$f_k(x) = \varphi(x - k)f(x) \quad \text{and} \quad g_k(x) = \varphi(x - k)g(x).$$

Now set

$$u_k(x) = \varphi(x - k)u(x) \quad \text{and} \quad v_k(x) = \varphi(x - k)v(x)$$

for $k \in \mathbb{Z}^d$. We then have

$$\operatorname{div}(uv) = \sum_{k \in \mathbb{Z}^d} (u_k v_k) = \sum_{k \in \mathbb{Z}^d} w_k, \quad w_k = \operatorname{div}(u_k v_k).$$

We are going to check that

$$\sum_{k \in \mathbb{Z}^d} \|w_k\|_{\mathcal{H}^1(\mathbb{R}^d)} < \infty.$$

To do this, we apply the local version (Case A). It follows that

$$\begin{aligned} \|w_k\|_{\mathcal{H}^1(\mathbb{R}^n)} &\leq C(\|u_k\|_{L^2} + \|\operatorname{div} u_k\|_{L^2})(\|v_k\|_{L^2} + \|\operatorname{div} v_k\|_{L^2}) \\ &= \varepsilon_k \in l^1(\mathbb{Z}^d), \end{aligned}$$

where

$$\varepsilon_k = C(\|u_k\|_{L^2} + \|\operatorname{div} u_k\|_{L^2})(\|v_k\|_{L^2} + \|\operatorname{div} v_k\|_{L^2}).$$

Up to now we have proved

$$(1.6) \quad \|\operatorname{div}(uv)\|_{\mathcal{H}^1(\mathbb{R}^d)} \leq C(\|u\|_{L^2} + \|\operatorname{div} u\|_{L^2})(\|v\|_{L^2} + \|\operatorname{div} v\|_{L^2}).$$

This automatically yields the estimate

$$(1.7) \quad \|\operatorname{div}(uv)\|_{\mathcal{H}^1(\mathbb{R}^d)} \leq C(\|u\|_{L^2} \|\nabla v\|_{L^2} + \|v\|_{L^2} \|\operatorname{div} u\|_{L^2}).$$

To see this, we may replace u in the above inequality by

$$u = \delta^{1/2-d/2} u\left(\frac{x}{\delta}\right) \quad \text{whenever } 0 < \delta < \infty,$$

and similarly v by

$$v_\delta = \delta^{1/2-d/2} v\left(\frac{x}{\delta}\right) \quad \text{whenever } 0 < \delta < \infty.$$

Thus the left-hand side of (1.6) fortunately does not change, while on the right-hand side we get rid of the undesirable terms by letting δ to be equal either to 0 or to $+\infty$. This completes the proof. \square

Now we turn to the proof of Lemma 3. One can show that every function $f \in L^p(\mathbb{R}^d)$, $p \in (1, +\infty]$, with compact support and $\int f \, dx = 0$ belongs to $\mathcal{H}^1(\mathbb{R}^d)$. In particular, we have

Lemma 3. *If $d^* = d/(d-1)$, $f \in L^{d^*}$, $\text{supp } f \subset \overline{\Omega}$ and*

$$\int f \, dx = 0,$$

then $f \in \mathcal{H}^1(\mathbb{R}^d)$.

Proof. We have

$$f = \text{div}(G)v + G \cdot \nabla v$$

and we have to prove that both the terms belong to L^{d^*} . We consider the first term on the right-hand side. Since $\nabla v \in L^2$, we have

$$\text{div}(G) \in L^2 \quad \text{and} \quad v \in L^q \quad \text{where} \quad \frac{1}{2} - \frac{1}{q} = \frac{1}{d}.$$

Thus,

$$v \, \text{div}(G) \in L^{d^*}.$$

A similar argument works on the second term and this completes the proof of the lemma. \square

1.2. Multipliers and Morrey-Campanato spaces

In this section we give a description of the multiplier space \dot{X}_r introduced recently by P. G. Lemarié-Rieusset in his work [9] (see also [10]). The space \dot{X}_r of pointwise multipliers which map L^2 into \dot{H}^{-r} is defined in the following way.

Definition 2. For $0 \leq r < d/2$ we define the homogeneous space \dot{X}_r by

$$\dot{X}_r = \{f \in L^2_{\text{loc}} : \forall g \in \dot{H}^r \, fg \in L^2\},$$

where we denote by $\dot{H}^r(\mathbb{R}^d)$ the completion of the space $\mathcal{D}(\mathbb{R}^d)$ with respect to the norm $\|u\|_{\dot{H}^r} = \|(-\Delta)^{r/2}u\|_{L^2}$.

The norm of \dot{X}_r is given by the operator norm of pointwise multiplication

$$\|f\|_{\dot{X}_r} = \sup_{\|g\|_{\dot{H}^r} \leq 1} \|fg\|_{L^2}.$$

Similarly, we define the nonhomogeneous space X_r for $0 \leq r < d/2$ equipped with the norm

$$\|f\|_{X_r} = \sup_{\|g\|_{H^r} \leq 1} \|fg\|_{L^2}.$$

We have the homogeneity properties : $\forall x_0 \in \mathbb{R}^d$

$$\begin{aligned} \|f(x + x_0)\|_{X_r} &= \|f\|_{X_r}, \\ \|f(x + x_0)\|_{\dot{X}_r} &= \|f\|_{\dot{X}_r}, \\ \|f(\lambda x)\|_{X_r} &\leq \frac{1}{\lambda^r} \|f\|_{X_r}, \quad 0 < \lambda \leq 1, \\ \|f(\lambda x)\|_{\dot{X}_r} &\leq \frac{1}{\lambda^r} \|f\|_{\dot{X}_r}, \quad \lambda > 0. \end{aligned}$$

The imbeddings

$$\begin{aligned} L^{d/t} &\subset X_r, \quad 0 \leq r < \frac{d}{2}, \quad 0 \leq t \leq r, \\ L^{d/r} &\subset \dot{X}_r, \quad 0 \leq r < \frac{d}{2} \end{aligned}$$

hold.

Example 1. If $u(x) \in \mathcal{D}(\mathbb{R}^d)$, $\varphi(x) = \left(\sum_{k=1}^d |x_k|^{\gamma_k} \right)^{-1}$, $\gamma_k > 0$, $d > 2$ and $\sum_{k=1}^d \gamma_k^{-1} = d/2$, then

$$\int_{\mathbb{R}^d} \varphi(x) |u(x)|^2 dx \leq C \int_{\mathbb{R}^d} |\nabla u(x)|^2 dx$$

and $\varphi \in X_{1t}(\mathbb{R}^d)$.

Indeed, the inequality

$$\begin{aligned} &\int_{\lambda < |x| < 2\lambda} \varphi(x) |u(x)|^2 dx \\ &\leq \left[\int_{\lambda < |x| < 2\lambda} |u(x)|^{2d/(d-2)} dx \right]^{(d-2)/2} \left[\int_{\lambda < |x| < 2\lambda} \varphi(x)^{d/2} dx \right]^{2/d} \end{aligned}$$

and the Sobolev theorem imply that for $\lambda > 0$

$$\int_{\lambda < |x| < 2\lambda} \varphi(x) |u(x)|^2 dx \leq C \left[\int_{\lambda < |x| < 2\lambda} |\nabla u(x)|^2 dx + \int_{\lambda < |x| < 2\lambda} \frac{|u(x)|^2}{|x|^2} dx \right] \\ \times \left[\int_{\lambda < |x| < 2\lambda} \varphi(x)^{d/2} dx \right]^{2/d},$$

where C does not depend on λ . Let us estimate the integral

$$S(\lambda) = \int_{\lambda < |x| < 2\lambda} \varphi(x)^{d/2} dx.$$

The domain $\lambda < |x| < 2\lambda$ can be represented as a finite sum of domains $\Omega_{j\lambda}$ such that $|x_j| > \frac{1}{2}\lambda$ if $x \in \Omega_{j\lambda}$ for $j = 1, \dots, d$. Let for instance $|x_1| > \frac{1}{2}\lambda$. Then

$$\int_{\Omega_{j\lambda}} \varphi(x)^{d/2} dx \leq \frac{3}{2}\lambda \int_{\lambda < |x| < 2\lambda} \frac{dx_1 \dots dx_d}{((\frac{1}{2}\lambda)^{\gamma_1} + |x_2|^{\gamma_2} + \dots + |x_d|^{\gamma_d})^{d/2}}.$$

The substitution $x_j = t_j(\frac{1}{2}\lambda)^{\gamma_1/\gamma_j}$ gives the relations

$$S(\lambda) \leq C \int_{\mathbb{R}^{d-1}} \frac{dt_1 \dots dt_d}{(1 + |t_2|^{\gamma_2} + \dots + |t_d|^{\gamma_d})^{d/2}} \leq C,$$

since the integral is converging. To see this, set $t_s = \tau_s^{1/\gamma_s}$. Then

$$\int_{\mathbb{R}^{d-1}} \frac{dt_1 \dots dt_d}{(1 + |t_2|^{\gamma_2} + \dots + |t_d|^{\gamma_d})^{d/2}} \\ \leq C \int_{\mathbb{R}^{d-1}} \frac{|\tau|^{1/\gamma_2 + \dots + 1/\gamma_d - (d-1)}}{(1 + |\tau|)^{d/2}} d\tau_1 \dots d\tau_d \\ \leq C \int_0^\infty \frac{d|\tau|}{(1 + |\tau|)^{1/\gamma+1}} < \infty.$$

Therefore,

$$\int_{\lambda < |x| < 2\lambda} \varphi(x) |u(x)|^2 dx \leq C_5 \left[\int_{\lambda < |x| < 2\lambda} |u(x)|^2 dx + \int_{\lambda < |x| < 2\lambda} \frac{|u(x)|^2}{|x|^2} dx \right].$$

Setting $\lambda = 2^m$, $m \in \mathbb{Z}$ and assuming these inequalities for all m , we obtain that

$$\int_{\mathbb{R}^d} \varphi(x) |u(x)|^2 dx \leq C \left(\int_{\mathbb{R}^d} |\nabla u(x)|^2 dx + \int_{\mathbb{R}^d} \frac{|u(x)|^2}{|x|^2} dx \right).$$

By Hardy's inequality in \mathbb{R}^d , $d \geq 3$, we have

$$\int_{\mathbb{R}^d} \frac{|u(x)|^2}{|x|^2} dx \leq \frac{4}{(d-2)^2} \int_{\mathbb{R}^d} |\nabla u(x)|^2 dx, \quad u(x) \in \mathcal{D}(\mathbb{R}^d),$$

and hence

$$\int_{\mathbb{R}^d} \varphi(x)|u(x)|^2 dx \leq C \int_{\mathbb{R}^d} |\nabla u(x)|^2 dx.$$

Now we recall the definition of the Morrey-Campanato spaces ([6], [19]):

Definition 3. For $1 < p \leq q \leq +\infty$, the Morrey-Campanato space $\mathcal{M}_{p,q}$ is defined by

$$(1.8) \quad \mathcal{M}_{p,q} = \{f \in L^p_{\text{loc}}(\mathbb{R}^d) : \|f\|_{\mathcal{M}_{p,q}} = \sup_{x \in \mathbb{R}^d} \sup_{0 < R \leq 1} R^{d/q-d/p} \|f(y)1_{B(x,R)}(y)\|_{L^p(dy)} < \infty\}.$$

Let us define the homogeneous Morrey-Campanato spaces $\dot{\mathcal{M}}_{p,q}$ for $1 < p \leq q \leq +\infty$ by

$$(1.9) \quad \|f\|_{\dot{\mathcal{M}}_{p,q}} = \sup_{x \in \mathbb{R}^d} \sup_{R > 0} R^{d/q-d/p} \left(\int_{B(x,R)} |f(y)|^p dy \right)^{1/p}.$$

It is easy to check the following properties:

$$\|f(\lambda x)\|_{\mathcal{M}_{p,q}} = \frac{1}{\lambda^{d/q}} \|f\|_{\mathcal{M}_{p,q}}, \quad 0 < \lambda \leq 1,$$

$$\|f(\lambda x)\|_{\dot{\mathcal{M}}_{p,q}} = \frac{1}{\lambda^{d/q}} \|f\|_{\dot{\mathcal{M}}_{p,q}}, \quad \lambda > 0.$$

We shall use the following classical results [6].

a) For $1 \leq p \leq p'$, $p \leq q \leq +\infty$ and for all functions f such that $f \in \dot{\mathcal{M}}_{p,q} \cap L^\infty$ we have

$$\|f\|_{\dot{\mathcal{M}}_{p',q \cdot (p'/p)}} \leq \|f\|_{L^\infty}^{1-p/p'} \|f\|_{\dot{\mathcal{M}}_{p,q}}^{p/p'}.$$

b) For p, q, p', q' such that $1/p + 1/p' \leq 1$, $1/q + 1/q' \leq 1$, $f \in \dot{\mathcal{M}}_{p,q}$, $g \in \dot{\mathcal{M}}_{p',q'}$ we have

$$fg \in \dot{\mathcal{M}}_{p'',q''} \quad \text{with} \quad 1/p + 1/p' = 1/p'', \quad 1/q + 1/q' = 1/q''.$$

c) For $1 \leq p \leq d$ we have

$$\forall \lambda > 0, \quad \|\lambda f(\lambda x)\|_{\dot{\mathcal{M}}_{p,d}} = \|f\|_{\dot{\mathcal{M}}_{p,d}}.$$

d) If $p' < p$ then

$$\begin{aligned}\dot{\mathcal{M}}_{p,q} &\subset \mathcal{M}_{p,q}, \\ \dot{\mathcal{M}}_{p,q} &\subset \mathcal{M}_{p',q}.\end{aligned}$$

e) If $q_2 < q_1$ then

$$\begin{aligned}\mathcal{M}_{p,q_1} &\subset \mathcal{M}_{p,q_2}, \\ L^q = \dot{\mathcal{M}}_{q,q} &\subset \dot{\mathcal{M}}_{p,q}, \quad p \leq q.\end{aligned}$$

We have the following comparison between multipliers and Morrey-Campanato spaces:

Proposition 1. For $0 \leq r < d/2$, we have

$$\begin{aligned}X_r &\subseteq \mathcal{M}_{2,d/r}, \\ \dot{X}_r &\subseteq \dot{\mathcal{M}}_{2,d/r}.\end{aligned}$$

Proof. Let $f \in X_r$, $0 < R \leq 1$, $x_0 \in \mathbb{R}^d$ and $\phi \in \mathcal{D}$, $\phi \equiv 1$ on $B(x_0/R, 1)$. We have

$$\begin{aligned}R^{r-d/2} \left(\int_{|x-x_0| \leq R} |f(x)|^2 dx \right)^{1/2} &= R^r \left(\int_{|y-x_0/R| \leq 1} |f(Ry)|^2 dy \right)^{1/2} \\ &\leq R^r \left(\int_{y \in \mathbb{R}^d} |f(Ry)\phi(y)|^2 dy \right)^{1/2} \\ &\leq R^r \|f(Ry)\|_{X_r} \|\phi\|_{H^r} \\ &\leq \|f(y)\|_{X_r} \|\phi\|_{H^r} \leq C \|f(y)\|_{X_r}.\end{aligned}$$

We observe that the same proof is also valid for homogeneous spaces. □

Additionally, for $2 < p \leq d/r$ and $0 \leq r < d/2$ we have the following inclusion relations:

$$L^{d/r}(\mathbb{R}^d) \subset L^{d/r,\infty}(\mathbb{R}^d) \subset \dot{\mathcal{M}}_{p,d/r}(\mathbb{R}^d) \subset \dot{X}_r(\mathbb{R}^d) \subset \dot{\mathcal{M}}_{2,d/r}(\mathbb{R}^d),$$

where $L^{p,\infty}$ denotes the usual Lorentz (weak L^p) space. For the definition and basic properties of Lorentz spaces $L^{p,q}$ we refer to [16].

2. UNIQUENESS THEOREM

Before turning our attention to uniqueness issues, we start with some prerequisites for our main result. Let

$$C_{0,\sigma}^\infty(\mathbb{R}^d) = \{\varphi \in (C_0^\infty(\mathbb{R}^d))^d : \operatorname{div} \varphi = 0\} \subseteq (C_0^\infty(\mathbb{R}^d))^d.$$

The subspace

$$L_\sigma^2(\mathbb{R}^d) = \overline{C_{0,\sigma}^\infty(\mathbb{R}^d)}^{\|\cdot\|_{L^2}} = \{u \in L^2(\mathbb{R}^d)^d : \operatorname{div} u = 0\}$$

is obtained as the closure of $C_{0,\sigma}^\infty$ with respect to the L^2 -norm $\|\cdot\|_{L^2}$. H_σ^r denotes the closure of $C_{0,\sigma}^\infty$ with respect to the norm

$$\|u\|_{H^r} = \|u\|_{L^2} + \|(1 - \Delta)^{r/2} u\|_{L^2} \quad \text{for } r \geq 0.$$

Our definition of the Leray-Hopf weak solutions (see e.g., [8], [7]) now reads:

Definition 4 (weak solutions). Let $a \in L_\sigma^2$ and $T > 0$. A measurable function u is called a weak solution of (1.1) on $(0, T)$ if u has the following properties:

1. $u \in L^\infty((0, T); L_\sigma^2) \cap L^2((0, T); \dot{H}_\sigma^1)$ for all $T > 0$;
2. $u(t)$ is continuous in time in the weak topology of L_σ^2 with

$$\langle u(t), \phi \rangle \rightarrow \langle a, \phi \rangle \quad \text{as } t \rightarrow 0^+$$

for all $\phi \in L_\sigma^2$;

3. for any $0 \leq s \leq t \leq T$, u satisfies the identity

$$(2.1) \quad \int_s^t \{-\langle u, \partial_\tau \phi \rangle + \langle u \cdot \nabla u, \phi \rangle + \langle \nabla u, \nabla \phi \rangle\} d\tau = -\langle u(t), \phi(t) \rangle + \langle u(s), \phi(s) \rangle$$

for all $\phi \in H^1((s, t); H_\sigma^1)$. Here $\langle \cdot, \cdot \rangle$ denotes the scalar product and $\|\cdot\|_{L^2}$ denotes the norm in $L^2(\mathbb{R}^d)^d$.

Remark 2. For u and ϕ as above, the integral

$$\int_0^T \langle u \cdot \nabla u, \phi \rangle d\tau$$

is well defined since by the Sobolev inequality we have

$$\|u\|_{L^{2d/(d-2)}} \leq C \|\nabla u\|_{L^2}$$

so that

$$\begin{aligned} \left| \int_0^T \langle u \cdot \nabla u, \phi \rangle d\tau \right| &\leq \int_0^T \|u\|_{L^{2d/(d-2)}} \|\nabla u\|_{L^2} \|\phi\|_{L^d} d\tau \\ &\leq C \sup_{0 < t < T} \|\phi\|_{L^d} \int_0^T \|\nabla u\|_{L^2}^2 d\tau. \end{aligned}$$

Existence of weak solutions has been established by Leray in [11] for the initial velocity in $L^2_\sigma(\mathbb{R}^d)$. The result is the following

Theorem 3 (Leray-Hopf). *Let $T > 0$. Then, for any given $a \in L^2_\sigma(\mathbb{R}^d)$, there exists at least one weak solution u to (1.1) on $(0, T)$ such that*

$$(2.2) \quad \|u(t)\|_{L^2}^2 + 2 \int_0^t \|\nabla u(s)\|_{L^2}^2 ds \leq \|a\|_{L^2}^2, \quad 0 \leq t < T$$

and

$$\|u(t) - a\|_{L^2} \rightarrow 0 \quad \text{as } t \rightarrow +0.$$

Let us introduce the class $L^s((0, T); L^\gamma)$ with the norm $\|\cdot\|_{L^s((0, T); L^\gamma)}$ by

$$\|u\|_{L^s((0, T); L^\gamma)} = \left(\int_0^T \|u(t)\|_{L^\gamma}^s dt \right)^{1/s}.$$

The classical result on uniqueness of weak solutions in the class $L^s((0, T); L^\gamma)$ was given by Foias, Serrin and Masuda [3], [14], [13].

Theorem 4 (Foias-Serrin-Masuda). *Let $a \in L^2_\sigma(\mathbb{R}^d)$. Let u and v be two weak solutions of (1.1) on $(0, T)$. Suppose that u satisfies*

$$(2.3) \quad u \in L^s((0, T); L^\gamma) \quad \text{for } \frac{2}{s} + \frac{d}{\gamma} = 1 \quad \text{with } d < \gamma < \infty.$$

Assume that v fulfils the energy inequality (2.2) for $0 \leq t < T$. Then we have $u = v$ on $[0, T)$.

Remark 3. In Theorem 4, v need not belong to the class (2.3). On the other hand, every weak solution u with (2.3) fulfils the energy identity

$$(2.4) \quad \|u(t)\|_{L^2}^2 + 2 \int_0^t \|\nabla u(s)\|_{L^2}^2 ds = \|a\|_{L^2}^2, \quad 0 \leq t \leq T.$$

It seems to be an interesting question whether every weak solution satisfies the energy inequality (2.2).

Remark 4. The class (2.3) is important from the view point of scaling invariance for the Navier-Stokes equations. It can be easily seen that if (u, p) is a pair of the solution to (1.1) on $\mathbb{R}^d \times (0, T)$, then so is the family $\{u_\lambda, p_\lambda\}_{\lambda>0}$ where

$$u_\lambda(x, t) = \lambda u(\lambda x, \lambda^2 t), \quad p_\lambda(x, t) = \lambda^2 u(\lambda x, \lambda^2 t).$$

Scaling invariance means that

$$\|u_\lambda\|_{L^s((0, \infty); L^\gamma)} = (\lambda^{1-(2/s+d/\gamma)}) \|u\|_{L^s((0, \infty); L^\gamma)} = \|u\|_{L^s((0, \infty); L^\gamma)} \quad \text{for all } \lambda > 0$$

holds if and only if

$$\frac{2}{s} + \frac{d}{\gamma} = 1.$$

We shall next deal with the critical case with $s = \infty$ and $\gamma = d$ in (2.3).

Theorem 5 (Masuda [13], Kozono-Sohr [7]). *Let $a \in L^2_\sigma(\mathbb{R}^d)$. Let u and v be two weak solutions of (1.1) on $(0, T)$. Suppose that*

$$(2.5) \quad u \in L^\infty((0, T); L^d)$$

and that v fulfils the energy inequality (2.2) for all $0 \leq t < T$. Then we have $u = v$ on $[0, T)$.

Remark 5. Masuda [13] proved that if $u \in L^\infty((0, T); L^d)$ is continuous from the right on $[0, T)$ in the norm of L^d , then $u = v$ holds on $[0, T)$. Later on, Kozono-Sohr [7] showed that every weak solution in $L^\infty((0, T); L^d)$ of (1.1) on $(0, T)$ becomes necessarily continuous from the right in the norm of L^d .

The same result holds when for $\gamma = +\infty$ we replace the assumption

$$u \in L^2((0, T); L^\infty)$$

by the weaker assumption

$$u \in L^2((0, T); BMO(\mathbb{R}^d)^d).$$

The replacement of the hypothesis $u \in L^2((0, T); L^\infty)$ by $u \in L^2((0, T); BMO(\mathbb{R}^d)^d)$ was recently discussed in a similar context by Kozono and Taniuchi [8]. Moreover, we have

Theorem 6 (Kozono-Taniuchi). *Let $a \in L^2_\sigma(\mathbb{R}^d)$ and let u, v be two weak solutions of (1.1) on $(0, T)$. Suppose that*

$$(2.6) \quad u \in L^2((0, T); BMO(\mathbb{R}^d)^d)$$

and that v fulfils the energy inequality (2.2) for $0 \leq t < T$. Then we have $u = v$ on $[0, T]$.

Remark 6. By Theorem 4, every weak solution in $L^2((0, T); L^\infty)$ is unique.

Our result on uniqueness of the weak solution now reads:

Theorem 7. *Let $a \in L^2(\mathbb{R}^d)^d$ with $\nabla \cdot a = 0$. Assume that there exists a solution u for the Navier-Stokes equation on $(0, T) \times \mathbb{R}^d$ (for some $T \in (0, +\infty]$) with the initial data a so that*

$$u \in L^\infty((0, T); L^2_\sigma(\mathbb{R}^d)^d) \cap L^2((0, T); \dot{H}^1_\sigma(\mathbb{R}^d)^d)$$

and

$$\nabla u \in L^2((0, T); \dot{X}_1(\mathbb{R}^d)^d).$$

Then u is the unique Leray-Hopf solution associated with a on $[0, T]$.

The same result holds when the assumption $\nabla u \in L^2((0, T); \dot{X}_1(\mathbb{R}^d)^d)$ is replaced by $u \in L^2((0, T); BMO(\mathbb{R}^d)^d)$.

The following corollary, which is an immediate consequence of Theorem 7, gives a simpler sufficient condition in terms of the Lorentz spaces.

Corollary 1. *Let $a \in L^2(\mathbb{R}^d)^d$ with $\nabla \cdot a = 0$. Assume that there exists a solution u for the Navier-Stokes equation on $(0, T) \times \mathbb{R}^d$ (for some $T \in (0, +\infty]$) with initial data a so that*

$$u \in L^\infty((0, T); L^2_\sigma(\mathbb{R}^d)^d) \cap L^2((0, T); \dot{H}^1_\sigma(\mathbb{R}^d)^d)$$

and

$$\nabla u \in L^2((0, T); L^{d, \infty}(\mathbb{R}^d)^d),$$

where $L^{p, \infty}$ denotes the usual Lorentz (weak L^p) space. Then u is the unique Leray-Hopf solution associated with a on $[0, T]$.

The same result again holds when the assumption $\nabla u \in L^2((0, T); L^{d, \infty}(\mathbb{R}^d)^d)$ is replaced by $u \in L^2((0, T); L^d(\mathbb{R}^d)^d)$.

The following lemmas play a fundamental role in estimating the nonlinear term.

Lemma 4. Let $f \in H^1(\mathbb{R}^d)$, $g(x) = (g_i(x))_{i=1}^d$ with $\nabla \cdot g = 0$ and $g \in L^2(\mathbb{R}^d)^d$. Furthermore, we assume that $\nabla h \in \dot{X}_1(\mathbb{R}^d)$. Then there exists a constant $C(d) > 0$ independent of f , g and h such that

$$(2.7) \quad \left| \int_{\mathbb{R}^d} fg \cdot \nabla h \, dx \right| \leq C \|\nabla f\|_{L^2(\mathbb{R}^d)} \|g\|_{L^2(\mathbb{R}^d)^d} \|\nabla h\|_{\dot{X}_1(\mathbb{R}^d)}$$

and

$$(2.8) \quad \left| \int_{\mathbb{R}^d} \nabla f \cdot gh \, dx \right| \leq C \|\nabla f\|_{L^2(\mathbb{R}^d)} \|g\|_{L^2(\mathbb{R}^d)^d} \|\nabla h\|_{\dot{X}_1(\mathbb{R}^d)}.$$

Proof. The proof is easy, due to the definition of $\dot{X}_1(\mathbb{R}^d)$. Supposing that $\nabla h \in \dot{X}_1(\mathbb{R}^d)$ and using the Cauchy-Schwarz inequality, we get

$$\begin{aligned} \left| \int_{\mathbb{R}^d} fg \cdot \nabla h \, dx \right| &\leq \left(\int_{\mathbb{R}^d} |f|^2 |\nabla h|^2 \, dx \right)^{1/2} \|g\|_{L^2(\mathbb{R}^d)^d} \\ &\leq C \|\nabla h\|_{\dot{X}_1(\mathbb{R}^d)} \left(\int_{\mathbb{R}^d} |\nabla f|^2 \, dx \right)^{1/2} \|g\|_{L^2(\mathbb{R}^d)^d}, \end{aligned}$$

where the constant C is independent of f , g and h . Thus the lemma is proved in the case of (2.7). The proof is similar in the case of (2.8). \square

The same result holds when we replace the assumption $\nabla h \in \dot{X}_1(\mathbb{R}^d)$ by the assumption $h \in H^1(\mathbb{R}^d) \cap BMO(\mathbb{R}^d)$. Indeed, we know that

$$h(x) = \log |x| \in BMO$$

and

$$|\nabla h|^2 \leq \frac{1}{|x|^2},$$

hence by Hardy's inequality in \mathbb{R}^d ($d \geq 3$) we have

$$\int_{\mathbb{R}^d} \frac{|f(x)|^2}{|x|^2} \, dx \leq C(d) \int_{\mathbb{R}^d} |\nabla f|^2 \, dx, \quad \forall f \in H^1(\mathbb{R}^d).$$

This remark suggests that the lemma will also hold when we replace the $\dot{X}_1(\mathbb{R}^d)$ -norm of ∇h by the BMO -norm of h . In fact, the following is a combination of the compensated compactness results of Coifman, Lions, Meyer and Semmes [1] and the duality of the space BMO .

Lemma 5. Let $f \in H^1(\mathbb{R}^d)$, $g = (g_i(x))_{i=1}^d$ with $\nabla \cdot g = 0$ and $g \in L^2(\mathbb{R}^d)^d$ and let $h \in H^1(\mathbb{R}^d) \cap BMO(\mathbb{R}^d)$. Then there exists a constant $C(d) > 0$ independent of f , g and h such that

$$(2.9) \quad |\langle g \cdot \nabla f, h \rangle| \leq C \|\nabla f\|_{L^2(\mathbb{R}^d)} \|g\|_{L^2(\mathbb{R}^d)^d} \|h\|_{BMO(\mathbb{R}^d)}.$$

Proof. It is an immediate consequence of Lemma 2 and the duality inequality (1.4):

$$\begin{aligned} |\langle g \cdot \nabla f, h \rangle| &\leq C \|g \cdot \nabla f\|_{\mathcal{H}^1(\mathbb{R}^d)} \|h\|_{BMO(\mathbb{R}^d)} \\ &\leq C \|\nabla f\|_{L^2(\mathbb{R}^d)} \|g\|_{L^2(\mathbb{R}^d)^d} \|h\|_{BMO(\mathbb{R}^d)}. \end{aligned}$$

□

Next we recall the following well-known result:

Lemma 6 (Poincaré inequality). Suppose Q is a cube in \mathbb{R}^d of side length ϱ and f is C^2 on Q with $\nabla f \in L^2(Q)$. There exists c independent of f such that

$$(2.10) \quad \int_Q |f - m_Q f|^2 \, dy \leq c \varrho^2 \int_Q |\nabla f(y)|^2 \, dy,$$

where $m_Q f = 1/|Q| \int_Q f(y) \, dy$ is the integral mean of f on Q .

Combining this result with Proposition 1 gives

Proposition 2. If $f \in H^1(\mathbb{R}^d)$ and $\nabla f \in \dot{X}_1(\mathbb{R}^d)$, then

$$f \in BMO(\mathbb{R}^d).$$

Proof. Since $\dot{X}_1(\mathbb{R}^d) \subset \dot{\mathcal{M}}_{2,d}(\mathbb{R}^d)$, it follows that

$$\nabla f \in \dot{\mathcal{M}}_{2,d}(\mathbb{R}^d).$$

By the classical Poincaré inequality (2.10) we have

$$\begin{aligned} \int_{B(x,R)} |f(y) - m_{B(x,R)} f(y)|^2 \, dy &\leq CR^2 \int_{B(x,R)} |\nabla f(y)|^2 \, dy \\ &\leq CR^d \|\nabla f\|_{\dot{\mathcal{M}}_{2,d}}^2 \end{aligned}$$

for every ball $B(x, R)$ of any radius R and

$$\begin{aligned} \|f\|_{BMO}^2 &= \sup_{x \in \mathbb{R}^d} \sup_{R > 0} \frac{1}{|B(x, R)|} \int_{B(x, R)} |f(y) - m_{B(x, R)} f(y)|^2 \, dy \\ &\leq C \|\nabla f\|_{\dot{\mathcal{M}}_{2,d}}^2 \leq C \|\nabla f\|_{\dot{X}_1(\mathbb{R}^d)}^2. \end{aligned}$$

□

We are now in position to prove the main result.

P r o o f of Theorem 7. Let v be another weak solution of (1.1) associated to a on $(0, T)$ (with the associated pressure p) such that

$$v \in L^\infty((0, T); L^2_\sigma(\mathbb{R}^d)^d) \cap L^2((0, T); \dot{H}^1_\sigma(\mathbb{R}^d)^d)$$

and

$$\nabla v \in L^2((0, T); \dot{X}_1(\mathbb{R}^d)^d).$$

We consider the difference $w = u - v$ and we obtain

$$(2.11) \quad \begin{aligned} \partial_t w - \Delta w + \nabla p_w &= -[w \cdot \nabla v + u \cdot \nabla w], \\ \operatorname{div} w &= 0, \\ w(x, 0) &= 0. \end{aligned}$$

If we take the inner product $\langle \cdot, \cdot \rangle$ of L^2 with w , we get

$$\frac{1}{2} \frac{d}{dt} \|w\|_{L^2}^2 + \|\nabla w\|_{L^2}^2 = -\langle w \cdot \nabla v, w \rangle.$$

Integration by parts followed by integration over time then lead to

$$(2.12) \quad \|w(t)\|_{L^2}^2 + 2 \int_0^t \|\nabla w\|_{L^2}^2 d\tau = -2 \int_0^t \langle w \cdot \nabla v, w \rangle d\tau = 2 \int_0^t \langle w \cdot \nabla w, v \rangle d\tau$$

for all $0 \leq t < T$. Lemma 4 with

$$g = w, \quad \nabla f = \nabla w \quad \text{and} \quad h = v$$

yields directly

$$|\langle w \cdot \nabla w, v \rangle| \leq C \|\nabla w\|_{L^2(\mathbb{R}^d)} \|w\|_{L^2(\mathbb{R}^d)^d} \|\nabla v\|_{\dot{X}_1(\mathbb{R}^d)}.$$

We thus observe that by the Young inequality ($ab \leq a^2/2 + b^2/2$, $a, b \geq 0$) it follows that

$$\left| \int_0^t \langle w \cdot \nabla w, v \rangle d\tau \right| \leq \frac{1}{2} \int_0^t \|\nabla w\|_{L^2(\mathbb{R}^d)}^2 d\tau + \frac{C}{2} \int_0^t \|w\|_{L^2(\mathbb{R}^d)^d}^2 \|\nabla v\|_{\dot{X}_1(\mathbb{R}^d)}^2 d\tau.$$

Hence by (2.12)

$$\|w(t)\|_{L^2}^2 + \int_0^t \|\nabla w\|_{L^2}^2 d\tau \leq C \int_0^t \|w\|_{L^2(\mathbb{R}^d)^d}^2 \|\nabla v\|_{\dot{X}_1(\mathbb{R}^d)}^2 d\tau$$

for all $t > 0$. Since $\nabla v \in L^2((0, T); \dot{X}_1(\mathbb{R}^d)^d)$ and $w(0) = 0$, it follows from the Gronwall inequality that

$$\|w(t)\|_{L^2}^2 \leq \|w(0)\|_{L^2}^2 \exp\left(C \int_0^t \|\nabla v\|_{\dot{X}_1(\mathbb{R}^d)}^2 d\tau\right)$$

and thus

$$\|w(t)\|_{L^2}^2 = 0, \quad 0 \leq t < T,$$

which implies the uniqueness of weak solutions.

The proof when

$$u \in L^2((0, T); BMO(\mathbb{R}^d)^d)$$

is quite similar. We apply Lemma 5 with

$$g = w \quad \nabla f = \nabla w \quad \text{and} \quad h = v$$

which yields directly

$$|\langle w \cdot \nabla w, v \rangle| \leq C \|\nabla w\|_{L^2(\mathbb{R}^d)} \|w\|_{L^2(\mathbb{R}^d)^d} \|v\|_{BMO(\mathbb{R}^d)}.$$

Using again Young's inequality, we get

$$\left| \int_0^t \langle w \cdot \nabla w, v \rangle d\tau \right| \leq \frac{1}{2} \int_0^t \|\nabla w\|_{L^2(\mathbb{R}^d)}^2 d\tau + \frac{C}{2} \int_0^t \|w\|_{L^2(\mathbb{R}^d)^d}^2 \|v\|_{BMO(\mathbb{R}^d)}^2 d\tau.$$

Hence it follows from (2.12) that

$$\|w(t)\|_{L^2}^2 + \int_0^t \|\nabla w\|_{L^2}^2 d\tau \leq C \int_0^t \|w\|_{L^2(\mathbb{R}^d)^d}^2 \|v\|_{BMO(\mathbb{R}^d)}^2 d\tau$$

for all $0 \leq t < T$. Since $v \in L^2((0, T); BMO(\mathbb{R}^d)^d)$ and $w(0) = 0$, the Gronwall inequality yields

$$\|w(t)\|_{L^2}^2 = 0, \quad 0 \leq t < T,$$

from which we get the desired uniqueness. □

References

- [1] *R. Coifman, P.-L. Lions, Y. Meyer, S. Semmes*: Compensated compactness and Hardy spaces. *J. Math. Pures Appl.* 72 (1993), 247–286.
- [2] *P. Constantin*: Remarks on the Navier-Stokes equations. In: *New Perspectives in Turbulence*. Springer, New York, 1991, pp. 229–261.
- [3] *C. Foias*: Une remarque sur l'unicité des solutions des équations de Navier-Stokes en dimension n . *Bull. Soc. Math. Fr.* 89 (1961), 1–8. (In French.)
- [4] *C. Fefferman, E. M. Stein*: H^p spaces of several variables. *Acta Math.* 129 (1972), 137–193.
- [5] *E. Hopf*: Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen. *Math. Nachr.* 4 (1951), 213–231. (In German.)
- [6] *T. Kato*: Strong L^p solutions of the Navier-Stokes equation in Morrey spaces. *Bol. Soc. Bras. Mat.* 22 (1992), 127–155.
- [7] *H. Kozono, H. Sohr*: Remark on uniqueness of weak solutions to the Navier-Stokes equations. *Analysis* 16 (1996), 255–271.
- [8] *H. Kozono, Y. Taniuchi*: Bilinear estimates in BMO and the Navier-Stokes equations. *Math. Z.* 235 (2000), 173–194.
- [9] *P. G. Lemarié-Rieusset*: *Recent Developments in the Navier-Stokes Problem*. Chapman & Hall/CRC, Boca Raton, 2002.
- [10] *P. G. Lemarié-Rieusset, S. Gala*: Multipliers between Sobolev spaces and fractional differentiation. *J. Math. Anal. Appl.* 322 (2006), 1030–1054.
- [11] *J. Leray*: Sur le mouvement d'un liquide visqueux emplissant l'espace. *Acta Math.* 63 (1934), 193–248. (In French.)
- [12] *F. Murat*: Compacité par compensation. *Ann. Sc. Norm. Sup. Pisa, Cl. Sci.* 5 (1978), 489–507. (In French.)
- [13] *K. Masuda*: Weak solutions of Navier-Stokes equations. *Tôhoku Math. J.* 36 (1984), 623–646.
- [14] *J. Serrin*: On the interior regularity of weak solutions of the Navier-Stokes equations. *Arch. Ration. Mech. Anal.* 9 (1962), 187–195.
- [15] *E. M. Stein*: *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton Univ. Press, Princeton, 1993.
- [16] *E. M. Stein, G. Weiss*: *Introduction to Fourier Analysis on Euclidean spaces*. Princeton Mathematical series. Princeton University Press, Princeton, 1971.
- [17] *L. Tartar*: Compensated compactness and applications to partial differential equations. *Nonlinear Analysis and Mechanics: Heriot-Watt Symposium, Vol. 4, Edinburgh 1979*. *Res. Notes Math.* 39 (1979), 136–212.
- [18] *R. Temam*: *Navier-Stokes Equations. Theory and Numerical Analysis*. North-Holland, Amsterdam, 1977.
- [19] *M. E. Taylor*: Analysis on Morrey spaces and applications to Navier-Stokes and other evolution equations. *Commun. Partial Differ. Equations* 17 (1992), 1407–1456.

Author's address: *S. Gala*, University of Mostaganem, Department of Mathematics, Box 227, Mostaganem (27000), Algeria, e-mail: sadek.gala@gmail.com.