

Honghai Li; Jiongsheng Li

On potentially nilpotent double star sign patterns

Czechoslovak Mathematical Journal, Vol. 59 (2009), No. 2, 489–501

Persistent URL: <http://dml.cz/dmlcz/140493>

Terms of use:

© Institute of Mathematics AS CR, 2009

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON POTENTIALLY NILPOTENT DOUBLE STAR SIGN PATTERNS

HONGHAI LI, Nanchang, JIONGSHENG LI, Hefei

(Received September 10, 2007)

Abstract. A matrix \mathcal{A} whose entries come from the set $\{+, -, 0\}$ is called a *sign pattern matrix*, or *sign pattern*. A sign pattern is said to be potentially nilpotent if it has a nilpotent realization. In this paper, the characterization problem for some potentially nilpotent double star sign patterns is discussed. A class of double star sign patterns, denoted by $\mathcal{DSSP}(m, 2)$, is introduced. We determine all potentially nilpotent sign patterns in $\mathcal{DSSP}(3, 2)$ and $\mathcal{DSSP}(5, 2)$, and prove that one sign pattern in $\mathcal{DSSP}(3, 2)$ is potentially stable.

Keywords: sign pattern, double star, potentially nilpotent, potentially stable

MSC 2010: 05C50, 15A18

1. INTRODUCTION

A matrix \mathcal{A} whose entries come from the set $\{+, -, 0\}$ is called a *sign pattern matrix*, or *sign pattern*. Let $\mathcal{A} = [a_{ij}]$ and $\mathcal{B} = [b_{ij}]$ be $n \times n$ sign patterns. If $b_{ij} = a_{ij}$ whenever $a_{ij} \neq 0$, then \mathcal{B} is a superpattern of \mathcal{A} and \mathcal{A} is a subpattern of \mathcal{B} . If \mathcal{A} is a sign pattern and A is a real matrix for which each entry has the same sign as the corresponding entry of \mathcal{A} , then A is said to be a *realization* of \mathcal{A} , and we write $A \in Q(\mathcal{A})$. The *inertia* of a square matrix A is the ordered triple $i(A) = (i_+(A), i_-(A), i_0(A))$, in which $i_+(A)$, $i_-(A)$ and $i_0(A)$ are the numbers of eigenvalues of A with positive, negative and zero real parts, respectively. The *inertia* of sign pattern \mathcal{A} is the set of ordered triples $i(\mathcal{A}) = \{i(A) : A \in Q(\mathcal{A})\}$. A square matrix A is *stable* if $i(A) = (0, n, 0)$. A pattern \mathcal{A} is *potentially stable* if $(0, n, 0) \in i(\mathcal{A})$. A pattern \mathcal{A} is *inertially arbitrary* if $(r, s, t) \in i(\mathcal{A})$ for every

Supported by youth scientific funds of the Education Department of Jiangxi Province (No. GJJ09460), Natural Science Foundation of Jiangxi Normal University (No.2058), National Natural Science Foundation of China (No.10601001).

nonnegative triple (r, s, t) with $r + s + t = n$. In particular, an $n \times n$ ($n \geq 2$) sign pattern matrix \mathcal{A} is *spectrally arbitrary* if given any monic polynomial $r(x)$ of degree n with real coefficients, there exists $A \in Q(\mathcal{A})$ such that the characteristic polynomial of A is $r(x)$.

In the following discussion, we need some notations about the cycles in a sign pattern, since every real matrix associated with it has the same qualitative cycle structure, and such cycle structure is crucial in studying the eigenvalues. The *directed graph* $D(\mathcal{A})$ associated with a pattern $\mathcal{A} = (a_{ij})$ is the directed graph with vertex set $\{1, 2, \dots, n\}$ and arc (i, j) if $a_{ij} \neq 0$, for all $1 \leq i, j \leq n$. A *simple k -cycle* γ of length $|\gamma| = k$ of \mathcal{A} is a sequence of k arcs $(i_1, i_2), (i_2, i_3) \dots (i_k, i_1)$ in $D(\mathcal{A})$ such that the vertices i_1, i_2, \dots, i_k are distinct. We denote the cycle γ by $(i_1, i_2, \dots, i_k, i_1)$. Write $\prod_{\gamma} = a_{i_1 i_2} a_{i_2 i_3} \dots a_{i_k i_1}$, the *cycle product* of \mathcal{A} associated with a simple k -cycle γ .

It is well known that the determinant of an $n \times n$ matrix A is the sum of all possible terms of the form

$$(1.1) \quad (-1)^{|\gamma_1|-1} \prod_{\gamma_1} (-1)^{|\gamma_2|-1} \prod_{\gamma_2} \dots (-1)^{|\gamma_p|-1} \prod_{\gamma_p}$$

where $\gamma_1, \dots, \gamma_p$ are disjoint simple cycles the sum of whose lengths is equal to n . Similarly, $E_k(A)$, the sum of all $k \times k$ principal minors of A , is equal to the sum of all terms of the form (1.1) where $\gamma_1, \dots, \gamma_p$ are disjoint simple cycles whose length sum equals k . The computation of the characteristic polynomial of A is then expressed in terms of its cycle products as follows:

$$(1.2) \quad P_A(x) = x^n + \sum_{k=1}^n (-1)^k E_k(A) x^{n-k}.$$

An $n \times n$ sign pattern \mathcal{A} is said to be *potentially nilpotent* if there exists $A \in Q(\mathcal{A})$ such that $P_A(x) = x^n$. From (1.2), this is equivalent to finding $A \in \mathcal{A}$ such that $E_k(A) = 0$ for all $1 \leq k \leq n$. Clearly, if \mathcal{A} is spectrally arbitrary, then \mathcal{A} is potentially nilpotent, but not conversely. Two sign patterns \mathcal{A} and \mathcal{B} are *equivalent* if \mathcal{B} may be obtained from \mathcal{A} by some combination of negation, transposition, permutation similarity, and signature similarity. Note that if \mathcal{A} and \mathcal{B} are equivalent, then \mathcal{A} is potentially nilpotent if and only if \mathcal{B} is potentially nilpotent. In [6], Lina Yeh characterized some star sign patterns and linear tree sign patterns that are potentially nilpotent. In [3], Eschenbach and Li obtained a number of qualitative necessary or sufficient conditions for a sign pattern to be potentially nilpotent. In [5], all star sign patterns that are potentially nilpotent are characterized. In this paper, we introduce one class of double star sign patterns which is denoted by $\mathcal{DSSP}(m, 2)$

Corollary 2.1. Let $n \geq 4$. If $\mathcal{A} = [a_{ij}]$ is a spectrally arbitrary double star sign pattern, then $a_{ii} = 0$ for at most one leaf vertex i .

Proposition 2.3. Let $n \geq 4$ and A be a real matrix as in (2.1). If A is nilpotent, then the non-zero entries among a_2, \dots, a_m are distinct and so are the non-zero entries among a_{m+2}, \dots, a_n .

Proof. By Proposition 2.1, the characteristic polynomial of A is

$$P_A(x) = \left(\prod_{j=1}^m (x - a_j) - \sum_{i=2}^m \left(b_i \prod_{\substack{j=2 \\ j \neq i}}^m (x - a_j) \right) \right) \\ \times \left(\prod_{j=m+1}^n (x - a_j) - \sum_{i=m+2}^n \left(b_i \prod_{\substack{j=m+2 \\ j \neq i}}^n (x - a_j) \right) \right) - b_{m+1} \prod_{\substack{j=2 \\ j \neq m+1}}^n (x - a_j).$$

If $a_i = a_j$ for some $2 \leq i \neq j \leq m$, then a_i is a zero of $P_A(x) = x^n$, and consequently $a_i = 0$. Thus, if $a_i \neq 0$, then $a_i \neq a_j$ for all $2 \leq i \neq j \leq m$. Similarly, we may prove that if $a_i \neq 0$, then $a_i \neq a_j$ for all $m+2 \leq i \neq j \leq n$. \square

Proposition 2.4. Let A be a real matrix as in (2.1) with zero main diagonal. Then A is nilpotent if and only if either $\sum_{i=2}^m b_i = 0$ and $\sum_{i=m+1}^n b_i = 0$, or $\sum_{i=m+2}^n b_i = 0$ and $\sum_{i=2}^{m+1} b_i = 0$.

Proof. Consider the matrix A in (2.1) with $a_i = 0$ for $1 \leq i \leq n$. The characteristic polynomial of A is $P_A(x) = x^n + E_2(B)x^{n-2} + E_4(B)x^{n-4}$. Thus A being nilpotent is equivalent to

$$(2.2) \quad - \sum_{i=2}^n b_i = E_2(B) = 0,$$

and

$$(2.3) \quad \sum_{i=2}^m b_i \sum_{i=m+2}^n b_i = E_4(B) = 0.$$

From (2.3) we have either $\sum_{i=2}^m b_i = 0$ or $\sum_{i=m+2}^n b_i = 0$. Taking (2.2) into consideration, we see that A is nilpotent if and only if $\sum_{i=2}^m b_i = 0$ and $\sum_{i=m+1}^n b_i = 0$, or $\sum_{i=m+2}^n b_i = 0$ and $\sum_{i=2}^{m+1} b_i = 0$. \square

Proposition 2.5. *Let A be a real matrix as in (2.1) with the only nonzero entries a_1, a_{m+1} on the main diagonal. Then A is nilpotent if and only if $a_1 = -a_{m+1}$, $b_{m+1} = -a_1^2$ and $\sum_{i=2}^m b_i = \sum_{j=m+2}^n b_j = 0$.*

Proof. For the matrix A in (2.1), if $a_i = 0$ for $2 \leq i \leq m$ and $m+2 \leq i \leq n$, then $P_A(x) = x^n - E_1(A)x^{n-1} + E_2(A)x^{n-2} - E_3(A)x^{n-3} + E_4(A)x^{n-4}$. The matrix A is nilpotent if and only if

$$(2.4) \quad a_1 + a_{m+1} = E_1(A) = 0,$$

$$(2.5) \quad a_1 a_{m+1} - \sum_{i=2}^n b_i = E_2(A) = 0,$$

$$(2.6) \quad -a_1 \sum_{i=m+2}^n b_i - a_{m+1} \sum_{j=2}^m b_j = E_3(A) = 0,$$

and

$$(2.7) \quad \sum_{i=2}^m b_i \sum_{j=m+2}^n b_j = E_4(A) = 0.$$

From (2.7), $\sum_{i=2}^m b_i = 0$ or $\sum_{j=m+2}^n b_j = 0$. So, by (2.6), we have $\sum_{i=2}^m b_i = \sum_{j=m+2}^n b_j = 0$. From (2.4) and (2.5), we have $a_1 = -a_{m+1}$ and $b_{m+1} = -a_1^2$. \square

Now consider the following double star matrix

$$(2.8) \quad \begin{pmatrix} 0 & 1 & \dots & 1 & 1 \\ b_2 & a_2 & & & \\ \vdots & & \ddots & & \\ b_m & & & a_m & \\ b_{m+1} & & & & 0 & 1 \\ & & & & b_{m+2} & 0 \end{pmatrix},$$

where $m \geq 3$. The sign pattern \mathcal{A} determined by a real matrix A with form (2.8) is denoted by $\mathcal{DSSP}(m, 2)$. In the following we shall give a characterization for all potentially nilpotent sign patterns in $\mathcal{DSSP}(m, 2)$ for some small values of m .

Theorem 2.1. *Let $\mathcal{A} = [a_{ij}] \in \mathcal{DSSP}(3, 2)$. Then \mathcal{A} is potentially nilpotent if and only if $a_{22}a_{33} = -$, $a_{21}a_{31} = +$ and $(a_{22}, a_{21}, a_{41}, a_{54})$ is one of the forms (up to equivalence) $(+, -, +, -)$, $(+, +, -, +)$, $(+, -, +, +)$.*

Proof. Suppose $A \in Q(\mathcal{A})$ is parameterized as in (2.8). Then A is nilpotent if and only if

$$\begin{cases} a_2 + a_3 = 0, \\ a_2 a_3 - b_2 - b_3 - b_4 - b_5 = 0, \\ -a_2(b_3 + b_4 + b_5) - a_3(b_2 + b_4 + b_5) = 0, \\ -a_2 a_3(b_4 + b_5) + b_5(b_2 + b_3) = 0, \\ (a_2 b_3 + a_3 b_2) b_5 = 0. \end{cases}$$

By simplification, we have

$$\begin{aligned} a_2 &= -a_3, & b_4 + b_5 &= a_2 a_3 - b_2 - b_3, \\ b_5(b_2 + b_3) &= a_2 a_3(b_4 + b_5), & a_2 b_3 + a_3 b_2 &= 0, \end{aligned}$$

that is,

$$a_2 = -a_3, \quad b_2 = b_3 = \frac{a_2^4}{2(b_5 - a_2^2)}, \quad b_4 = \frac{b_5^2}{a_2^2 - b_5}.$$

We distinguish the following cases.

Case 1: $b_5 > 0$.

Subclass 1: $b_5 > a_2^2$. Then $b_2 > 0$ and $b_4 < 0$.

Subclass 2: $b_5 < a_2^2$. Then $b_2 < 0$ and $b_4 > 0$.

Case 2: If $b_5 < 0$, then $b_2 < 0$ and $b_4 > 0$.

Therefore, if \mathcal{A} is potentially nilpotent, then $a_{22}a_{33} = -$, $a_{21}a_{31} = +$ and $(a_{22}, a_{21}, a_{41}, a_{54})$ must be one of the forms $(+, -, +, -)$, $(+, +, -, +)$, $(+, -, +, +)$.

Conversely, for a sign pattern $\mathcal{A} = [a_{ij}] \in \mathcal{DSSP}(3, 2)$, if $a_{22}a_{33} = -$, $a_{21}a_{31} = +$ and $(a_{22}, a_{21}, a_{41}, a_{54})$ is one of the forms $(+, -, +, -)$, $(+, +, -, +)$, $(+, -, +, +)$, choosing real matrices $A \in Q(\mathcal{A})$ as in (2.8) satisfying $a_2 = -a_3$, $b_2 = b_3$, and $(a_2, b_2, b_4, b_5) = (1, \frac{1}{2}, -4, 2)$, $(1, -1, \frac{1}{2}, \frac{1}{2})$, $(1, -\frac{1}{4}, \frac{1}{2}, -1)$, respectively, it is easy to verify that A is nilpotent, respectively. \square

We next introduce some more notation. Let $X = \{x_j : j \in J\}$ be a set of variables with finite index set J . Let $\Sigma_i(X)$ be the sum of the products of the elements of the i -element subsets of X for each i with $1 \leq i \leq |J|$. For each $j \in J$, let $X_j = X \setminus \{x_j\}$. If $j \notin J$, we assume $X_j = X$. For example,

$$\Sigma_1(X_j) = \sum_{k \in J \setminus \{j\}} x_k, \quad \Sigma_2(X_j) = \sum_{\substack{k, l \in J \setminus \{j\} \\ k < l}} x_k x_l.$$

Theorem 2.2. *No sign pattern $\mathcal{A} \in \mathcal{DSSP}(4, 2)$ is potentially nilpotent.*

Proof. Suppose the matrix A is in $Q(\mathcal{A})$. Then we can assume that A has the form

$$A = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ b_2 & a_2 & & & \\ b_3 & & a_3 & & \\ b_4 & & & a_4 & \\ b_5 & & & & 0 & 1 \\ & & & & b_6 & 0 \end{pmatrix}.$$

Let $X = \{a_2, a_3, a_4\}$. Then we have

$$\left\{ \begin{array}{l} E_1(A) = \Sigma_1(X), \\ E_2(A) = \Sigma_2(X) - \sum_{i=2}^6 b_i, \\ E_3(A) = \Sigma_3(X) - \sum_{i=2}^6 b_i \Sigma_1(X_i), \\ E_4(A) = - \sum_{i=2}^6 b_i \Sigma_2(X_i) + b_6 \sum_{i=2}^4 b_i, \\ E_5(A) = - \sum_{i=5}^6 b_i \Sigma_3(X) + b_6 \sum_{i=2}^4 b_i \Sigma_1(X_i), \\ E_6(A) = b_6 \sum_{i=2}^4 b_i \Sigma_2(X_i). \end{array} \right.$$

If A is nilpotent, then $E_k(A) = 0$ for all k . By $E_1(A) = 0$, $E_3(A) = 0$ and $E_5(A) = 0$, we have

$$\left\{ \begin{array}{l} \Sigma_3(X) = \sum_{i=2}^4 b_i \Sigma_1(X_i), \\ \frac{b_5 + b_6}{b_6} \Sigma_3(X) = \sum_{i=2}^4 b_i \Sigma_1(X_i). \end{array} \right.$$

So $(b_5/b_6)a_2a_3a_4 = 0$, which is a contradiction and thus the asserted conclusion follows. \square

Theorem 2.3. *Let $\mathcal{A} = [a_{ij}] \in \mathcal{DSSP}(5, 2)$. Then \mathcal{A} is potentially nilpotent if and only if, up to equivalence, $a_{22}a_{44} = - = a_{33}a_{55}$, $a_{21}a_{41} = + = a_{31}a_{51}$ and $(a_{22}, a_{33}, a_{21}, a_{31}, a_{61}, a_{76})$ is one of the forms: $(-, -, +, -, -, +)$, $(-, -, -, -, +, +)$, $(-, -, -, +, +, -)$.*

Proof. Consider the matrix $A \in Q(\mathcal{A})$ as in (2.8). Let $X = \{a_2, a_3, a_4, a_5\}$. That A is nilpotent is equivalent to

$$(2.9) \quad \Sigma_1(X) = E_1(A) = 0,$$

$$(2.10) \quad \Sigma_2(X) - \sum_{i=2}^7 b_i = E_2(A) = 0,$$

$$(2.11) \quad \Sigma_3(X) - \sum_{i=2}^7 b_i \Sigma_1(X_i) = E_3(A) = 0,$$

$$(2.12) \quad \Sigma_4(X) - \sum_{i=2}^7 b_i \Sigma_2(X_i) + b_7 \sum_{i=2}^5 b_i = E_4(A) = 0,$$

$$(2.13) \quad -\sum_{i=2}^7 b_i \Sigma_3(X_i) + b_7 \sum_{i=2}^5 b_i \Sigma_1(X_i) = E_5(A) = 0,$$

$$(2.14) \quad -(b_6 + b_7) \Sigma_4(X) + b_7 \sum_{i=2}^5 b_i \Sigma_2(X_i) = E_6(A) = 0,$$

and

$$(2.15) \quad b_7 \sum_{i=2}^5 b_i \Sigma_3(X_i) = E_7(A) = 0.$$

From (2.19), we have

$$(2.16) \quad \sum_{i=2}^5 a_i = 0.$$

From (2.15), we have $\sum_{i=2}^5 b_i \Sigma_3(X_i) = 0$, that is,

$$(2.17) \quad \sum_{i=2}^5 \frac{b_i}{a_i} = 0.$$

From (2.13) and (2.11), we obtain that $b_7 \sum_{i=2}^5 b_i \Sigma_1(X_i) = (b_6 + b_7) \Sigma_3(X)$ and $\Sigma_3(X) = \sum_{i=2}^5 b_i \Sigma_1(X_i)$. Combining these two equations, we have

$$(2.18) \quad \sum_{i=2}^5 b_i a_i = 0,$$

$$(2.19) \quad \sum_{i=2}^5 \frac{1}{a_i} = 0.$$

Since the off-diagonal terms in a double star matrix enter into its characteristic polynomial only as products, $-\mathcal{A}$ is equivalent to the sign pattern obtained from \mathcal{A} by taking the negative of the main diagonal. So we may consider only two cases:

Case (1): $(a_2, a_3, a_4, a_5) = (-, +, +, +)$. From (2.16), we have $a_3 + a_4 + a_5 = -a_2$. But from (2.19), we have $1/a_3 + 1/a_4 + 1/a_5 = 1/(a_3 + a_4 + a_5)$, which is a contradiction.

Case (2): $(a_2, a_3, a_4, a_5) = (-, -, +, +)$. We shall prove that there are two pairs of oppositely signed numbers among a_2, a_3, a_4, a_5 . From (2.9), we have $a_2 + a_4 = -(a_3 + a_5)$. Let $a_2 = -x, a_3 = -y, \mu = a_2 + a_4$, where $y > \mu > -x, y > 0, x > 0$. Then $a_4 = x + \mu, a_5 = y - \mu$. By (2.19), we have $-1/x - 1/y + 1/(x + \mu) + 1/(y - \mu) = 0$, and so $(x + \mu)(y - \mu) = xy$, that is, $\mu = 0$ or $\mu = y - x$. Without loss of generality, we always assume that $a_2 = -a_4, a_3 = -a_5$. Thus from (2.17) and (2.18), we have

$$\begin{cases} \frac{1}{a_2}(b_2 - b_4) + \frac{1}{a_3}(b_3 - b_5) = 0, \\ a_2(b_2 - b_4) + a_3(b_3 - b_5) = 0. \end{cases}$$

Consequently, either $b_2 = b_4$ and $b_3 = b_5$ or $a_2 = a_3$ and $b_2 + b_3 = b_4 + b_5$.

Let us distinguish two cases:

Case 1: $a_2 = a_3$ and $b_2 + b_3 = b_4 + b_5$. In this case, $X = \{a_2, a_3, a_4, a_5\} = \{a_2, a_2, -a_2, -a_2\}$ and

$$\begin{aligned} \Sigma_1(X) &= 0, & \Sigma_2(X) &= -2a_2^2, & \Sigma_3(X) &= 0, & \Sigma_4(X) &= a_2^4, \\ \Sigma_1(X_2) &= -a_2 = \Sigma_1(X_3), & \Sigma_1(X_4) &= a_2 = \Sigma_1(X_5), \\ \Sigma_2(X_2) &= \Sigma_2(X_3) = \Sigma_2(X_4) = \Sigma_2(X_5) = -a_2^2, \\ \Sigma_3(X_2) &= a_2^3 = \Sigma_3(X_3), & \Sigma_3(X_4) &= -a_2^3 = \Sigma_2(X_5). \end{aligned}$$

Solving the equations (2.10), (2.12) and (2.14), we get the following equations

$$\begin{cases} 2a_2^2 + 2(b_2 + b_3) + b_6 + b_7 = 0 \text{ (by (2.10))}, \\ a_2^4 + 2a_2^2(b_2 + b_3) + 2a_2^2(b_6 + b_7) + 2b_7(b_2 + b_3) = 0 \text{ (by (2.12))}, \\ a_2^2(b_6 + b_7) + 2b_7(b_2 + b_3) = 0 \text{ (by (2.14))}. \end{cases}$$

By solving the above equations, we come to a contradiction that $a_2 = 0$.

Case 2: $b_2 = b_4$ and $b_3 = b_5$. In this case it is sufficient to solve the six entries $a_2, a_3, b_2, b_3, b_6, b_7$ to insure the nilpotent matrix A . Now $X = \{a_2, a_3, a_4, a_5\} = \{a_2, a_3, -a_2, -a_3\}$ and

$$\begin{aligned} \Sigma_1(X) &= 0, & \Sigma_2(X) &= -a_2^2 - a_3^2, & \Sigma_3(X) &= 0, & \Sigma_4(X) &= a_2^2 a_3^2, \\ \Sigma_1(X_2) &= -a_2, & \Sigma_1(X_3) &= -a_3, & \Sigma_1(X_4) &= a_2, & \Sigma_1(X_5) &= a_3, \\ \Sigma_2(X_2) &= -a_2^2, & \Sigma_2(X_3) &= -a_2^2, & \Sigma_2(X_4) &= -a_3^2, & \Sigma_2(X_5) &= -a_2^2, \\ \Sigma_3(X_2) &= a_2 a_3^2, & \Sigma_3(X_3) &= a_2^2 a_3, & \Sigma_3(X_4) &= -a_2 a_3^2, & \Sigma_2(X_5) &= -a_2^2 a_3. \end{aligned}$$

By a straightforward calculation of the equations (2.10), (2.12) and (2.14), we get the following linear system of equations

$$\begin{cases} a_2^2 + a_3^2 + 2(b_2 + b_3) + b_6 + b_7 = 0 \text{ (by (2.10))}, \\ a_2^2 a_3^2 + 2(b_2 a_3^2 + b_3 a_2^2) + (a_2^2 + a_3^2)(b_6 + b_7) + 2b_7(b_2 + b_3) = 0 \text{ (by (2.12))}, \\ a_2^2 a_3^2 (b_6 + b_7) + 2b_7(b_2 a_3^2 + b_3 a_2^2) = 0 \text{ (by 2.14)}. \end{cases}$$

It is easy to see that the system is underdetermined. In this case, the first three components b_2, b_3, b_6 can be solved in terms of the last three components a_2, a_3, b_7 , that is

$$\begin{aligned} b_2 &= \frac{a_2^6}{2(b_7 - a_2^2)(a_2^2 - a_3^2)}, \\ b_3 &= \frac{-a_3^6}{2(b_7 - a_3^2)(a_2^2 - a_3^2)}, \\ b_6 &= \frac{-b_7^2}{(b_7 - a_2^2)(b_7 - a_3^2)}. \end{aligned}$$

Without loss of generality, we always assume $a_2^2 > a_3^2$.

Subcase 2.1: $b_7 > 0$.

Subcase 2.1.1: $b_7 > a_2^2 \Rightarrow b_6 < 0, b_2 > 0, b_3 < 0$.

Subcase 2.1.2: $a_3^2 < b_7 < a_2^2 \Rightarrow b_6 > 0, b_2 < 0, b_3 < 0$.

Subcase 2.1.3: $b_7 < a_3^2 \Rightarrow b_6 < 0, b_2 < 0, b_3 > 0$.

Subcase 2.2: $b_7 < 0 \Rightarrow b_6 > 0, b_2 < 0, b_3 > 0$.

Note that the sign pattern in Subcase 2.1.1 is permutation similar to that in Subcase 2.1.3.

Therefore, if \mathcal{A} is potentially nilpotent, then $a_{22}a_{44} = - = a_{33}a_{55}$, $a_{21}a_{41} = + = a_{31}a_{51}$ and $(a_{22}, a_{33}, a_{21}, a_{31}, a_{61}, a_{76})$ must be one of the forms: $(-, -, +, -, -, +)$, $(-, -, -, -, +, +)$, $(-, -, -, +, +, -)$.

The same result may be obtained if $a_2^2 < a_3^2$.

Conversely, for a sign pattern $\mathcal{A} = [a_{ij}] \in \mathcal{DSSP}(5, 2)$, if $a_{22}a_{44} = - = a_{33}a_{55}$, $a_{21}a_{41} = + = a_{31}a_{51}$ and $(a_{22}, a_{33}, a_{21}, a_{31}, a_{61}, a_{76})$ is one of the forms $(-, -, +, -, -, +)$, $(-, -, -, -, +, +)$, $(-, -, -, +, +, -)$, choosing real matrices $A \in Q(\mathcal{A})$ as in (2.8) satisfying $a_2 = -a_4$, $a_3 = -a_5$, $b_2 = b_4$, $b_3 = b_5$, and $(a_2, a_3, b_2, b_3, b_6, b_7) = (-2, -1, \frac{32}{3}, -\frac{1}{24}, -\frac{125}{4}, 5)$, $(-2, -1, -\frac{64}{15}, -\frac{1}{3}, \frac{27}{10}, \frac{3}{2})$, $(-2, -1, -\frac{32}{15}, \frac{1}{12}, \frac{1}{10}, -1)$, respectively, it is not difficult to verify that A is nilpotent, respectively. \square

One can see that with the increasing values of m , the work of giving the proofs becomes more and more arduous. Up to now it is not clear how the main results of the paper can be extended to larger size matrices, that is to say, where $m > 5$. A new method might be needed for these cases.

Remark 2.1. It is known that by the method in [2], which is sometimes called the Nilpotent-Jacobian method, if a nilpotent realization of a sign pattern \mathcal{A} can be found (this is usually not an easy task), for which the Jacobian is nonzero, then every superpattern of \mathcal{A} is spectrally arbitrary. Up to now we conclude that some sign patterns in $\mathcal{DSSP}(m, 2)$ for $m = 3, 5$ are potentially nilpotent. It is natural to ask whether these sign patterns are spectrally arbitrary. Unfortunately, the answer is negative.

In other words, we have

Theorem 2.4. *No sign pattern \mathcal{A} in $\mathcal{DSSP}(3, 2)$ or $\mathcal{DSSP}(5, 2)$ is spectrally arbitrary.*

Proof. Consider sign patterns in $\mathcal{DSSP}(3, 2)$. Without loss of generality, it may be assumed that each realization of sign pattern \mathcal{A} in $\mathcal{DSSP}(3, 2)$ has the form of A in (2.8). The characteristic polynomial of A is

$$\begin{aligned} P_A(x) &= x^5 - (a_2 + a_3)x^4 + (a_2a_3 - b_2 - b_3 - b_4 - b_5)x^3 \\ &\quad + (a_2(b_3 + b_4 + b_5) + a_3(b_2 + b_4 + b_5))x^2 \\ &\quad + (b_5(b_2 + b_3) - a_2a_3(b_4 + b_5))x - (a_2b_3 + a_3b_2)b_5. \end{aligned}$$

If $a_2 + a_3 = 0$ and $(a_2b_3 + a_3b_2)b_5 = 0$, then $a_2(b_3 + b_4 + b_5) + a_3(b_2 + b_4 + b_5) = 0$, so $P_A(x)$ cannot equal $x^5 + \alpha x^2$ for any nonzero α . Thus \mathcal{A} is not spectrally arbitrary. Consequently, $\mathcal{DSSP}(3, 2)$ does not contain sign patterns that are spectrally arbitrary. Similarly, it is easy to verify that $\mathcal{DSSP}(5, 2)$ contains no sign pattern that is spectrally arbitrary. \square

However, for the sign pattern

$$\mathcal{A} = \begin{pmatrix} 0 & + & + & + \\ - & - & & \\ - & & + & \\ + & & & 0 & + \\ & & & - & 0 \end{pmatrix}$$

in $\mathcal{DSSP}(3, 2)$, it can be proved that \mathcal{A} is potentially stable. In [1], Bone presented some constructions of potentially stable sign patterns, one of which is the following:

Construction 1'. Given an $(n - 1) \times (n - 1)$ matrix A^- , we construct an $n \times n$ matrix A as follows:

- (i) let A^- appear as a principal submatrix of A ;
- (ii) A contains a negative n -cycle.

Theorem 2.5. *The sign pattern \mathcal{A} (as above) is potentially stable.*

Proof. We know that \mathcal{A}_4 is potentially stable from the result in [4]. It is easy to see that A contains a negative 5-cycle. \square

Example 2.1. For the matrix

$$A = \begin{pmatrix} 0 & 1 & 1 & 1 & 0 \\ -3.87265 & -2.31427 & 0 & 0 & 0 \\ -9.917 & 0 & 0.123349 & 0 & 0 \\ 0.79188 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -0.720757 & 0 \end{pmatrix} \in Q(\mathcal{A}),$$

the spectrum of A is $\{-0.227038 + 3.43587i, -0.227038 - 3.43587i, -1.73634, -0.000253292 + 0.887001i, -0.000253292 - 0.887001i\}$. So A is a stable matrix and \mathcal{A} is potentially stable.

Acknowledgments.

Authors are grateful to the anonymous referee whose comments have led to a better presentation.

References

- [1] *T. Bone*: Positive feedback may sometimes promote stability. *Linear Algebra Appl.* 51 (1983), 143–151.
- [2] *J. H. Drew, C. R. Johnson, D. D. Olesky, P. van den Driessche*: Spectrally arbitrary patterns. *Linear Algebra Appl.* 308 (2000), 121–137.
- [3] *C. A. Eschenbach, Z. Li*: Potentially nilpotent sign pattern matrices. *Linear Algebra Appl.* 299 (1999), 81–99.
- [4] *C. R. Johnson, T. S. Summers*: The potentially stable tree sign patterns for dimensions less than five. *Linear Algebra Appl.* 126 (1989), 1–13.
- [5] *G. MacGillivray, R. M. Tifenbach, P. van den Driessche*: Spectrally arbitrary star sign patterns. *Linear Algebra Appl.* 400 (2005), 99–119.
- [6] *L. Yeh*: Sign pattern matrices that allow a nilpotent matrix. *Bull. Aust. Math. Soc.* 53 (1996), 189–196.

Authors' addresses: Honghai Li, College of Mathematic and Information Science, Jiangxi Normal University, Nanchang, JiangXi, 330022, P. R. China, e-mail: 1hh@mail.ustc.edu.cn; Jiongsheng Li, Department of Mathematics, University of Science and Technology of China, Hefei, Anhui, 230026, P. R. China, e-mail: lijs@ustc.edu.cn.