

Fatih Nuray

Lacunary weak statistical convergence

Mathematica Bohemica, Vol. 136 (2011), No. 3, 259–268

Persistent URL: <http://dml.cz/dmlcz/141648>

Terms of use:

© Institute of Mathematics AS CR, 2011

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

LACUNARY WEAK STATISTICAL CONVERGENCE

FATİH NURAY, Afyonkarahisar

(Received February 15, 2010)

Abstract. The aim of this work is to generalize lacunary statistical convergence to weak lacunary statistical convergence and \mathcal{I} -convergence to weak \mathcal{I} -convergence. We start by defining weak lacunary statistically convergent and weak lacunary Cauchy sequence. We find a connection between weak lacunary statistical convergence and weak statistical convergence.

Keywords: weak convergence, statistical convergence, lacunary sequence, lacunary statistical convergence

MSC 2010: 40A05, 46A25

1. INTRODUCTION

A number sequence (x_k) is statistically convergent to L provided that for every $\varepsilon > 0$,

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - L| \geq \varepsilon\}| = 0$$

where the vertical bars indicate the number of elements in the enclosed set [2], [11].

By a lacunary sequence we mean an increasing integer sequence $\theta = (k_r)$ such that $k_0 = 0$ and $h_r := k_r - k_{r-1} \rightarrow \infty$ as $r \rightarrow \infty$. Throughout this paper the intervals determined by θ will be denoted by $I_r := (k_{r-1}, k_r]$.

Let θ be a lacunary sequence; the number sequence (x_k) is lacunary statistically convergent to L provided that for every $\varepsilon > 0$,

$$\lim_r \frac{1}{h_r} |\{k \in I_r : |x_k - L| \geq \varepsilon\}| = 0$$

(see [3]). The space N_θ of N_θ -convergent sequences is defined by

$$N_\theta := \left\{ (x_k) : \text{for some } L, \lim_r \frac{1}{h_r} \sum_{k \in I_r} |x_k - L| = 0 \right\}.$$

Let B be a Banach space, let (x_k) be a B -valued sequence, and $x \in B$.

1. The sequence (x_k) is weakly C_1 -convergent to x provided that for any f in the continuous dual B^* of B ,

$$\lim_n \frac{1}{n} \sum_{k=1}^n f(x_k - x) = 0.$$

2. The sequence (x_k) is weakly convergent to x provided that for any f in the continuous dual B^* of B ,

$$\lim_k f(x_k - x) = 0.$$

In this case we write $w\text{-}\lim x_k = x$.

3. The sequence (x_k) is norm statistically convergent to x provided that

$$\delta(\{k: \|x_k - x\| \geq \varepsilon\}) = 0$$

where $\delta(A) = \lim_n n^{-1} |\{k \leq n: k \in A\}|$.

4. The sequence (x_k) is weakly statistically convergent to x provided that for any f in the continuous dual B^* of B , the sequence $(f(x_k - x))$ is statistically convergent to 0 (see [1]).

2. WEAKLY LACUNARY STATISTICALLY CONVERGENT SEQUENCE

Definition 1. Let B be a Banach space, let (x_k) be a B -valued sequence, θ a lacunary sequence and $x \in B$.

1. The sequence (x_k) is norm lacunary statistically convergent to x provided that

$$\delta_r(\{k: \|x_k - x\| \geq \varepsilon\}) = 0$$

where $\delta_r(A) = \lim_r h_r^{-1} |\{k \in I_r: k \in A\}|$.

2. The sequence (x_k) is weakly lacunary statistically convergent to x provided that for any f in the continuous dual B^* of B , the sequence $(f(x_k - x))$ is lacunary statistically convergent to 0.

3. The sequence (x_k) is weakly N_θ -convergent to x provided that, for any f in the continuous dual B^* of B , the sequence $(f(x_k - x))$ is N_θ -convergent to 0.

Let WS and WS_θ denote the sets of all weakly statistically convergent and weakly lacunary statistically convergent sequences, respectively.

3. WEAK LACUNARY STATISTICALLY CAUCHY SEQUENCE

In [5], Fridy and Orhan defined the lacunary statistical Cauchy sequence for a complex number sequence (x_k) as follows:

Let θ be a lacunary sequence. The sequence (x_k) is said to be lacunary statistically Cauchy if there is a subsequence $(x_{k'(r)})$ of x such that $k'(r) \in I_r$ for each r , $\lim x_{k'(r)} = x$ and for every $\varepsilon > 0$

$$\lim_r \frac{1}{h_r} |\{k \in I_r : |x_k - x_{k'(r)}| \geq \varepsilon\}| = 0.$$

Now we will give the definition of the weakly lacunary statistically Cauchy sequence for a B -valued sequence (x_k) .

Definition 2. Let B be a Banach space, (x_k) a B -valued sequence, θ a lacunary sequence and $x \in B$. The sequence (x_k) is weakly lacunary statistically Cauchy if there is a subsequence $(x_{k'(r)})$ of (x_k) such that $k'(r) \in I_r$ for each r , $w\text{-}\lim x_{k'(r)} = x$, and for any f in the continuous dual B^* of B and for every $\varepsilon > 0$

$$\lim_r \frac{1}{h_r} |\{k \in I_r : |f(x_k - x_{k'(r)})| \geq \varepsilon\}| = 0.$$

Theorem 3. A sequence (x_k) is weakly lacunary statistically convergent if and only if (x_k) is a weakly lacunary statistically Cauchy sequence.

Proof. Let (x_k) be a weakly lacunary statistically Cauchy sequence. Then for every $\varepsilon > 0$ we have

$$\begin{aligned} & |\{k \in I_r : |f(x_k - x)| \geq \varepsilon\}| \\ & \leq \left| \left\{ k \in I_r : |f(x_k - x_{k'(r)})| \geq \frac{\varepsilon}{2} \right\} \right| + \left| \left\{ k \in I_r : |f(x_{k'(r)} - x)| \geq \frac{\varepsilon}{2} \right\} \right|, \end{aligned}$$

hence we get that the sequence (x_k) is weakly lacunary statistically convergent. Let (x_k) be weakly lacunary statistically convergent to x and write $M_j = \{k \in \mathbb{N} : |f(x_k - x)| < 1/j\}$ for each $j \in \mathbb{N}$, $M_j \supseteq M_{j+1}$ and $|M_j \cap I_r|/h_r \rightarrow 1$ as $r \rightarrow \infty$. Choose m_1 such that $r \geq m_1$ implies $|M_1 \cap I_r|/h_r > 0$, i.e., $M_1 \cap I_r \neq \emptyset$. Next choose $m_1 < m_2$ such that $r \geq m_2$ implies $M_2 \cap I_r \neq \emptyset$. Then for each r satisfying $m_1 \leq r \leq m_2$, choose $k'(r) \in I_r$ such that $k'(r) \in I_r \cap M_1$. In this way, choose $m_{l+1} > m_l$ such that $r > m_{l+1}$ implies $M_{l+1} \cap I_r \neq \emptyset$. Then for all r satisfying $m_l \leq r < m_{l+1}$, choose $k'(r) \in I_r \cap M_l$, i.e.,

$$|f(x_{k'(r)} - x)| < \frac{1}{l}.$$

Hence we get $k'(r) \in I_r$ for every r , and $w\text{-}\lim x_{k'(r)} = x$. Also, we have, for every $\varepsilon > 0$,

$$\begin{aligned} & \frac{1}{h_r} |\{k \in I_r : |f(x_k - x_{k'(r)})| \geq \varepsilon\}| \\ & \leq \frac{1}{h_r} \left| \left\{ k \in I_r : |f(x_{k(r)} - x)| \geq \frac{\varepsilon}{2} \right\} \right| + \frac{1}{h_r} \left| \left\{ k \in I_r : |f(x_{k'(r)} - x)| \geq \frac{\varepsilon}{2} \right\} \right|, \end{aligned}$$

whence (x_k) is a weakly lacunary statistically Cauchy sequence. □

4. INCLUSION THEOREMS

In this section we first give a theorem that provides the relation between weak N_θ - and weak lacunary statistical convergences. We also study the inclusions between weak statistical convergence and weak lacunary statistical convergence.

Theorem 4. *Let θ be a lacunary sequence; then (x_k) is weakly N_θ -convergent to x if and only if (x_k) is weakly lacunary statistically convergent to x .*

Proof. If $\varepsilon > 0$ and (x_k) is weakly N_θ -convergent to x , we can write

$$\begin{aligned} \lim_r \frac{1}{h_r} \sum_{k \in I_r} |f(x_k - x)| & \geq \lim_r \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |f(x_k - x)| \geq \varepsilon}} |f(x_k - x)| \\ & \geq \varepsilon |\{k \in I_r : |f(x_k - x)| \geq \varepsilon\}|, \end{aligned}$$

so (x_k) is weakly lacunary statistically convergent to x .

Conversely, suppose that (x_k) is weakly lacunary statistically convergent to x . Since $f \in B^*$, f is bounded, say $|f(x_k - x)| \leq K$ for all k . Given $\varepsilon > 0$, we get

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} |f(x_k - x)| & = \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |f(x_k - x)| \geq \varepsilon}} |f(x_k - x)| + \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |f(x_k - x)| < \varepsilon}} |f(x_k - x)| \\ & \leq \frac{K}{h_r} |\{k \in I_r : |f(x_k - x)| \geq \varepsilon\}| + \varepsilon, \end{aligned}$$

so (x_k) is weakly N_θ -convergent to x . □

Theorem 5. For any lacunary sequence θ , $\text{WS-lim } x_k = x$ implies $\text{WS}_\theta\text{-lim } x_k = x$ if and only if $\liminf_r k_r/k_{r-1} > 1$.

Proof. k_r/k_{r-1} will be denoted by q_r . If $\liminf_r q_r > 1$ there exist $\eta > 0$ such that $1 + \eta \leq q_r$ for all sufficiently large r , which implies that

$$\frac{h_r}{k_r} \geq \frac{1}{1 + \eta}.$$

If $x_k \rightarrow x(\text{WS})$, then for every $\varepsilon > 0$ and for sufficiently large r we have

$$\begin{aligned} \frac{1}{k_r} |\{k \leq k_r : |f(x_k - x)| \geq \varepsilon\}| &\geq \frac{1}{k_r} |\{k \in I_r : |f(x_k - x)| \geq \varepsilon\}| \\ &\geq \frac{\eta}{1 + \eta} \frac{1}{h_r} |\{k \in I_r : |f(x_k - x)| \geq \varepsilon\}|; \end{aligned}$$

this proves sufficiency. Conversely, if we suppose that $\liminf_r q_r = 1$, then following the idea in [4], we can find a sequence (x_k) such that $(x_k) \notin \text{WS}_\theta$ but $(x_k) \in \text{WS}$. \square

Theorem 6. For any lacunary sequence θ , $\text{WS}_\theta\text{-lim } x_k = x$ implies $\text{WS-lim } x_k = x$ if and only if $\limsup_r k_r/k_{r-1} < \infty$.

Proof. If $\limsup_r q_r < \infty$, then there is a $K > 0$ such that $q_r < K$ for all r . Suppose that $x_k \rightarrow x(\text{WS}_\theta)$, and let $M_r = |\{k \in I_r : |f(x_k - x)| \geq \varepsilon\}|$. Since $\text{WS}_\theta\text{-lim } x_k = x$, given $\varepsilon > 0$, there is an $r_0 \in \mathbb{N}$ such that $M_r/h_r < \varepsilon$ for all $r > r_0$. Now let $M = \max\{M_r : 1 \leq r \leq r_0\}$ and let n be any integer satisfying $k_{r-1} < n \leq k_r$. Then we can write

$$\begin{aligned} \frac{1}{n} |\{k \leq n : |f(x_k - x)| \geq \varepsilon\}| &\leq \frac{1}{k_{r-1}} |\{k \leq k_r : |f(x_k - x)| \geq \varepsilon\}| \\ &= \frac{1}{k_{r-1}} \{M_1 + M_2 + \dots + M_{r_0} + M_{r_0+1} + \dots + M_r\} \\ &\leq \frac{M}{k_{r-1}} r_0 + \frac{1}{k_{r-1}} \left\{ h_{r_0+1} \frac{M_{r_0+1}}{h_{r_0+1}} + \dots + h_r \frac{M_r}{h_r} \right\} \\ &\leq \frac{r_0 M}{k_{r-1}} + \frac{1}{k_{r-1}} \left(\sup_{r > r_0} \frac{M_r}{h_r} \right) \{h_{r_0+1} + \dots + h_r\} \\ &\leq \frac{r_0 M}{k_{r-1}} + \varepsilon \frac{k_r - k_{r_0}}{k_{r-1}} \\ &\leq \frac{r_0 M}{k_{r-1}} + \varepsilon q_r \\ &\leq \frac{r_0 M}{k_{r-1}} + \varepsilon K, \end{aligned}$$

and the sufficiency follows immediately.

Conversely, if we suppose that $\limsup_r q_r = \infty$, then following the idea in [4], we can find a sequence (x_k) such that $(x_k) \notin \text{WS}$ but $(x_k) \in \text{WS}_\theta$. \square

Combining Theorems 5 and 6 we get

Theorem 7. *Let θ be a lacunary sequence; then $WS = WS_\theta$ if and only if $1 < \liminf_r k_r/k_{r-1} \leq \limsup_r k_r/k_{r-1} < \infty$.*

Theorem 8. *If $x \in WS \cap WS_\theta$, then $WS_\theta\text{-lim } x = WS\text{-lim } x$.*

Proof. Suppose $WS\text{-lim } x = x$ and $WS_\theta\text{-lim } x = y$ and $x \neq y$. For $\varepsilon < \frac{1}{2}|x - y|$ we get

$$\lim_n \frac{1}{n} |\{k \leq n : |f(x_k - y)| \geq \varepsilon\}| = 1.$$

Consider the k_m th term of the weak statistical limit expression $n^{-1}|\{k \leq n : |f(x_k - y)| \geq \varepsilon\}|$:

$$\begin{aligned} (1) \quad \frac{1}{k_m} \left| \left\{ k \in \bigcup_{r=1}^m I_r : |f(x_k - y)| \geq \varepsilon \right\} \right| &= \frac{1}{k_m} \sum_{r=1}^m |\{k \in I_r : |f(x_k - y)| \geq \varepsilon\}| \\ &= \frac{1}{\sum_{r=1}^m h_r} \sum_{r=1}^m h_r t_r, \end{aligned}$$

where $t_r = h_r^{-1}|\{k \in I_r : |f(x_k - y)| \geq \varepsilon\}| \rightarrow 0$ because $WS_\theta\text{-lim } x = y$. Since θ is a lacunary sequence, (1) is a regular weighted mean transform of t_r , and therefore it, too, tends to zero as $m \rightarrow \infty$. Also, since this is a subsequence of $\{n^{-1}|\{k \leq n : |f(x_k - y)| \geq \varepsilon\}|\}$, we infer that

$$\lim_n \frac{1}{n} |\{k \leq n : |f(x_k - y)| \geq \varepsilon\}| \neq 1,$$

and this contradiction shows that we can't have $x \neq y$. □

5. WEAK STRONG ALMOST CONVERGENCE AND WEAK LACUNARY STATISTICAL CONVERGENCE

The idea of almost convergence was introduced by Lorentz [9]. Later Maddox [10] and (independently) Freedman et al. [6] introduced the notion of the strong almost convergence. Now we will introduce the notions of weakly almost convergence and weakly strong almost convergence for sequences in a Banach space.

Definition 9. Let B be a Banach space, (x_k) be a B -valued sequence and let f be in the continuous dual B^* of B . Sequence (x_k) is said to be weakly almost convergent to x if

$$\lim_n \frac{1}{n} \sum_{i=m+1}^{m+n} f(x_i - x) = 0$$

uniformly in m .

Definition 10. Let B be a Banach space, (x_k) be a B -valued sequence and let f be in the continuous dual B^* of B . Sequence (x_k) is said to be weakly strongly almost convergent to x if

$$\lim_n \frac{1}{n} \sum_{i=m+1}^{m+n} |f(x_i - x)| = 0$$

uniformly in m .

Let WN_θ , WS_θ , WAC and $[WAC]$ denote the sets of all weakly N_θ -convergent, all weakly statistically convergent, all weakly almost convergent and all weakly strongly almost convergent sequences, respectively.

Lemma 11. $[WAC] = \bigcap_{\theta \in \mathcal{L}} WN_\theta$.

Proof is similar to the proof of Theorem 3.1 in [6].

Theorem 12. If \mathcal{L} denotes the set of all lacunary sequences, then

$$[WAC] = \bigcap_{\theta \in \mathcal{L}} WS_\theta.$$

Proof. By Lemma 12 and Theorem 4, we have

$$[WAC] = \bigcap_{\theta \in \mathcal{L}} WN_\theta = \bigcap_{\theta \in \mathcal{L}} WS_\theta.$$

□

6. WEAK \mathcal{I} -CONVERGENCE

The concept of the \mathcal{I} -convergence is a generalization of statistical convergence and is based on the notion of the ideal \mathcal{I} of subsets of the set \mathbb{N} of positive integers. A non-void class $\mathcal{I} \subseteq 2^{\mathbb{N}}$ is called an ideal if \mathcal{I} is additive (i.e., $A, B \in \mathcal{I} \Rightarrow A \cup B \in \mathcal{I}$) and hereditary (i.e., $A \in \mathcal{I}$ and $B \subseteq A \Rightarrow B \in \mathcal{I}$).

An ideal \mathcal{I} is said to be non-trivial if $\mathcal{I} \neq 2^{\mathbb{N}}$. A non-trivial ideal \mathcal{I} is said to be admissible if \mathcal{I} contains every finite subset of \mathbb{N} . For any ideal \mathcal{I} there is a filter $\mathcal{F}(\mathcal{I})$ corresponding to \mathcal{I} , given by

$$\mathcal{F}(\mathcal{I}) = \{K \subseteq \mathbb{N} : \mathbb{N} \setminus K \in \mathcal{I}\}.$$

Definition 13. Let B be a Banach space, let (x_k) be a B -valued sequence, and $x \in B$. The sequence (x_k) is norm \mathcal{I} -convergent to x provided that

$$\{k \in \mathbb{N} : \|x_k - x\| \geq \varepsilon\} \in \mathcal{I}.$$

Definition 14. Let B be a Banach space, let f be in the continuous dual B^* of B , let (x_k) be a B -valued sequence, and $x \in B$. The sequence (x_k) is weakly \mathcal{I} -convergent to x provided that

$$\{k \in \mathbb{N} : |f(x_k - x)| \geq \varepsilon\} \in \mathcal{I}.$$

If $\mathcal{I} = \mathcal{I}_{\text{fin}}$ the ideal of all finite subsets of \mathbb{N} , we have the usual weak convergence. Denote by \mathcal{I}_δ the class of all $K \subset \mathbb{N}$ with

$$\delta(K) = \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : k \in K\}| = 0,$$

then \mathcal{I}_δ is a non-trivial admissible ideal, and the \mathcal{I}_δ -convergence coincides with the weak statistical convergence.

Denote by \mathcal{I}_θ the class of all $K \subset \mathbb{N}$ with

$$\delta_r(K) = \lim_{r \rightarrow \infty} \frac{1}{h_r} |\{k \in I_r : k \in K\}| = 0,$$

then \mathcal{I}_θ is a non-trivial admissible ideal, \mathcal{I}_θ -convergence coincides with the weak lacunary statistical convergence.

Definition 15. Let B be a Banach space, (x_k) a B -valued sequence and let f be in the continuous dual B^* of B , and $x \in B$. The sequence (x_k) is weakly \mathcal{I}^* -convergent to x if and only if there exists a set $M = \{m_1 < m_2 < \dots < m_k < \dots\} \subseteq \mathbb{N}$, $M \in \mathcal{F}(\mathcal{I})$ such that $\lim_k f(x_{m_k} - x) = 0$.

Let \mathcal{WI} and \mathcal{WI}^* denote the sets of all weakly \mathcal{I} -convergent and all weakly \mathcal{I}^* -convergent sequences, respectively.

Theorem 16. Let \mathcal{I} be an admissible ideal. If $\mathcal{WI}^*\text{-lim } x_k = x$, then $\mathcal{WI}\text{-lim } x_k = x$.

Proof. By assumption there is a set $L \in \mathcal{I}$ such that for $M = \mathbb{N} \setminus L = \{m_1 < m_2 < \dots < m_k < \dots\}$ we have

$$(2) \quad \lim_k f(x_{m_k} - x) = 0.$$

Let $\varepsilon > 0$. By (2), there exists $k_0 \in \mathbb{N}$ such that $|f(x_{m_k} - x)| < \varepsilon$ for each $k > k_0$. Then since \mathcal{I} is admissible, we get

$$\{k \in \mathbb{N} : |f(x_{m_k} - x)| \geq \varepsilon\} \subset L \cup \{m_1 < m_2 < \dots < m_{k_0}\} \in \mathcal{I}.$$

□

Definition 17 (see [8]). An admissible ideal $\mathcal{I} \subset 2^{\mathbb{N}}$ is said to satisfy the condition (AP) if for every countable family of mutually disjoint sets $\{A_1, A_2, \dots\}$ belonging to \mathcal{I} there exists a countable family of sets $\{B_1, B_2, \dots\}$ such that $A_j \triangle B_j$ is a finite set for $j \in \mathbb{N}$ and $B = \bigcup_{j=1}^{\infty} B_j \in \mathcal{I}$.

Theorem 18. Let $\mathcal{I} \subset 2^{\mathbb{N}}$ be an admissible ideal. If the ideal \mathcal{I} has property (AP), then for an arbitrary sequence $(x_k) \in X$, $\mathcal{WI}\text{-lim}(x_k) = x$ implies $\mathcal{WI}^*\text{-lim}(x_k) = x$.

Proof. Suppose that \mathcal{I} satisfies condition (AP). Let $\mathcal{WI}\text{-lim}(x_k) = x$. Then $\{k \in \mathbb{N} : |f(x_{k_m} - x)| \geq \varepsilon\} \in \mathcal{I}$ for $\varepsilon > 0$. Put $A_1 = \{k \in \mathbb{N} : |f(x_{k_m} - x)| \geq 1\}$ and $A_k = \{k \in \mathbb{N} : 1/k \leq |f(x_{k_m} - x)| \leq 1/(k+1)\}$ for $k \geq 2, k \in \mathbb{N}$. Obviously $A_i \cap A_j = \varnothing$ for $i \neq j$. By condition (AP) there exists a sequence of sets $(B_k)_{k \in \mathbb{N}}$ such that $A_j \triangle B_j$ are finite sets for $j \in \mathbb{N}$ and $B = \bigcup_{j=1}^{\infty} B_j \in \mathcal{I}$. It is sufficient to prove that for $M = \mathbb{N} \setminus B$ we have

$$(3) \quad \lim_{\substack{k \rightarrow \infty \\ k \in M}} f(x_k - x) = 0.$$

Let $\xi > 0$. Choose $k \in \mathbb{N}$ such that $1/(k+1) < \xi$. Then $\{k \in \mathbb{N} : |f(x_k - x)| \geq \xi\} \subset \bigcup_{j=1}^{n+1} A_j$. Since $A_j \triangle B_j, j = 1, 2, \dots, n+1$ are finite sets there exists $k_0 \in \mathbb{N}$ such that

$$(4) \quad \bigcup_{j=1}^{n+1} B_j \cap \{k \in \mathbb{N} : k > k_0\} = \bigcup_{j=1}^{n+1} A_j \cap \{k \in \mathbb{N} : k > k_0\}.$$

If $k > k_0$ and $k \notin B$, then $k \notin \bigcup_{j=1}^{n+1} B_j$ and by (4), $k \notin \bigcup_{j=1}^{n+1} A_j$. But then $|f(x_k - x)| < 1/(k+1) < \xi$; so (3) holds. \square

7. WEAK \mathcal{I} -LIMIT POINTS AND WEAK \mathcal{I} -CLUSTER POINTS

Definition 19. Let B be a Banach space, (x_k) a B -valued sequence, let f be in the continuous dual B^* of B and $x \in B$.

(a) An element $x \in X$ is said to be a weak \mathcal{I} -limit point of (x_k) provided that there exists a set $M = \{m_1 < m_2 < \dots < m_k < \dots\} \subseteq \mathbb{N}$ such that $M \notin \mathcal{I}$ and

$$\lim_{k \rightarrow \infty, k \in M} f(x_k - x) = 0.$$

(b) An element $x \in X$ is said to be a weak \mathcal{I} -cluster point of (x_k) if and only if for each $\varepsilon > 0$ we have $\{k \in \mathbb{N} : |f(x_k - x)| < \varepsilon\} \notin \mathcal{I}$.

Let $WI(\Lambda_x)$ and $WI(\Gamma_x)$ denote the sets of all WI -limit and WI -cluster points of x , respectively.

Theorem 20. *Let \mathcal{I} be an admissible ideal. Then for each sequence $(x_k) \in B$ we have $WI(\Lambda_x) \subset WI(\Gamma_x)$.*

Proof. Let $x \in WI(\Lambda_x)$. Then there exists a set $M = \{m_1 < m_2 < \dots < m_k < \dots\} \notin \mathcal{I}$ such that

$$(5) \quad \lim_{k \rightarrow \infty} f(x_{m_k} - x) = 0.$$

Take $\vartheta > 0$. According to (5) there exists $k_0 \in \mathbb{N}$ such that for $k > k_0$ we have $|f(x_{m_k} - x)| < \vartheta$. Hence $\{k \in \mathbb{N}: |f(x_k - x)| < \vartheta\} \supset M \setminus \{m_1, m_2, \dots, m_{k_0}\}$ and $\{k \in \mathbb{N}: |f(x_k - x)| < \vartheta\} \notin \mathcal{I}$, which means that $x \in WI(\Gamma_x)$. \square

Acknowledgement. The author is grateful to the referee for his kind remarks which improved the presentation of the paper.

References

- [1] *J. Connor, M. Ganichev, V. Kadets:* A characterization of Banach spaces with separable duals via weak statistical convergence. *J. Math. Anal. Appl.* *244* (2000), 251–261.
- [2] *J. A. Fridy:* On statistical convergence. *Analysis* *5* (1985), 301–313.
- [3] *J. A. Fridy:* Statistical limit points. *Proc. Am. Math. Soc.* *118* (1993), 1187–1192.
- [4] *J. A. Fridy, C. Orhan:* Lacunary statistical convergence. *Pac. J. Math.* *160* (1993), 43–51.
- [5] *J. A. Fridy, C. Orhan:* Lacunary statistical summability. *J. Math. Anal. Appl.* *173* (1993), 497–504.
- [6] *A. R. Freedman, J. J. Sember, M. Raphael:* Some Cesaro type summability spaces. *Proc. London Math. Soc., III. Ser.* *37* (1978), 508–520.
- [7] *A. R. Freedman, J. J. Sember:* Densities and summability. *Pac. J. Math.* *95* (1981), 293–305.
- [8] *P. Kostyrko, T. Šalát, W. Wilczyński:* \mathcal{I} -convergence. *Real Anal. Exch.* *26* (2000/2001), 669–686.
- [9] *G. G. Lorentz:* A contribution to the theory of divergent sequences. *Acta Math.* *80* (1948), 167–190.
- [10] *I. J. Maddox:* A new type of convergence. *Math. Proc. Camb. Philos. Soc.* *83* (1978), 61–64.
- [11] *I. J. Schoenberg:* The integrability of certain functions and related summability methods. *Am. Math. Mon.* *66* (1959), 361–375.

Author's address: Fatih Nuray, Department of Mathematics, Afyon Kocatepe University, Afyonkarahisar, Turkey, e-mail: fnuray@aku.edu.tr.