

# Applications of Mathematics

---

Jamol I. Baltaev; Milan Kučera; Martin Väth

A variational approach to bifurcation in reaction-diffusion systems with Signorini type boundary conditions

*Applications of Mathematics*, Vol. 57 (2012), No. 2, 143–165

Persistent URL: <http://dml.cz/dmlcz/142033>

## Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A VARIATIONAL APPROACH TO BIFURCATION IN  
REACTION-DIFFUSION SYSTEMS WITH SIGNORINI TYPE  
BOUNDARY CONDITIONS\*

JAMOL I. BALTAEV, Khorezm, MILAN KUČERA, Praha,  
MARTIN VÁTH, Praha

(Received August 24, 2010)

*Abstract.* We consider a simple reaction-diffusion system exhibiting Turing’s diffusion driven instability if supplemented with classical homogeneous mixed boundary conditions. We consider the case when the Neumann boundary condition is replaced by a unilateral condition of Signorini type on a part of the boundary and show the existence and location of bifurcation of stationary spatially non-homogeneous solutions. The nonsymmetric problem is reformulated as a single variational inequality with a potential operator, and a variational approach is used in a certain non-direct way.

*Keywords:* reaction-diffusion system, unilateral condition, variational inequality, local bifurcation, variational approach, spatial patterns

*MSC 2010:* 35B32, 35K57, 35J50, 35J57, 47J20

## 1. INTRODUCTION

Let  $\Omega \subset \mathbb{R}^m$  be a bounded domain with Lipschitzian boundary  $\partial\Omega$ , and let  $\Gamma_D$ ,  $\Gamma_N$ ,  $\Gamma_U$  be pairwise disjoint parts of  $\partial\Omega$ ,

$$(1.1) \quad \text{mes } \Gamma_D > 0, \quad \text{mes } \Gamma_U > 0, \quad \text{mes}(\partial\Omega \setminus (\Gamma_D \cup \Gamma_N \cup \Gamma_U)) = 0$$

(the  $(m - 1)$ -dimensional Lebesgue measure). Our goal is to show on the basis of a simple variational approach the existence and location of bifurcations of nontrivial

---

\*The research has been supported by the Academy of Sciences of the Czech Republic under the Grant IAA100190805 of the GAAV and the Institutional Research Plan AV0Z10190503.

solutions of the system

$$(1.2) \quad \begin{aligned} d_1 \Delta u + b_{11}u + b_{12}v + n(u) &= 0 \\ d_2 \Delta v + b_{21}u + b_{22}v &= 0 \end{aligned} \quad \text{in } \Omega$$

with unilateral boundary conditions, i.e.

$$(1.3) \quad \begin{cases} u = v = 0 \text{ on } \Gamma_D, & \frac{\partial u}{\partial \nu} = 0 \text{ on } \Gamma_N, & \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega \setminus \Gamma_D, \\ u \geq 0, & \frac{\partial u}{\partial \nu} \geq 0, & u \cdot \frac{\partial u}{\partial \nu} = 0 \text{ on } \Gamma_U. \end{cases}$$

Simultaneously, similar results will be proved also for a “dual” problem with a non-linearity  $n$  in the second equation and unilateral conditions for  $v$  instead of  $u$ . See Section 5 for further examples. It will be always assumed that the constant real matrix  $B = (b_{ij})$  satisfies

$$(1.4) \quad \begin{aligned} b_{11} > 0 > b_{22}, & \quad b_{12}b_{21} < 0, \\ \det B = b_{11}b_{22} - b_{12}b_{21} > 0, & \quad b_{11} + b_{22} < 0, \end{aligned}$$

$n$  will be a function satisfying

$$(1.5) \quad n(0) = n'(0) = 0,$$

and  $(d_1, d_2) \in \mathbb{R}_+^2 := (0, \infty) \times (0, \infty)$  will be real parameters (diffusion coefficients).

In order to explain the sense of our results and include it in the framework of the previous research, we must start our exposition with a more general reaction-diffusion system

$$(1.6) \quad \begin{aligned} u_t &= d_1 \Delta u + b_{11}u + b_{12}v + n_1(u, v) \\ v_t &= d_2 \Delta v + b_{21}u + b_{22}v + n_2(u, v) \end{aligned} \quad \text{in } (0, \infty) \times \Omega,$$

first with the classical mixed boundary conditions

$$(1.7) \quad u = v = 0 \text{ on } \Gamma_D, \quad \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega \setminus \Gamma_D.$$

Here,  $n_j(0, 0) = (\partial n_j / \partial u)(0, 0) = (\partial n_j / \partial v)(0, 0) = 0$ . Clearly,  $(0, 0)$  is a solution of (1.6) with (1.3) as well as with (1.7).

In terms of models of chemical reactions, the first line in (1.4) means that our system is of an activator-inhibitor type (the case  $b_{12} < 0 < b_{21}$ ) or of a positive feedback (substrate-depletion) type. See e.g. [4], [11], [16]. In the first case,  $u$  and  $v$  describe

the concentration of the activator and inhibitor, respectively. It is well known that under the assumption (1.4), the trivial solution of the problem without any diffusion (i.e. of ODE's,  $d_1 = d_2 = 0$ ) is stable but the trivial solution of the problem (1.6), (1.7) is stable only for parameters  $d_1, d_2$  from a certain subset  $D_S$  (the domain of stability) of  $\mathbb{R}_+^2$  and unstable for  $d_1, d_2 \in D_U = \mathbb{R}_+^2 \setminus \overline{D_S}$  (the domain of instability). See Proposition 2.1, Remark 2.2 and Fig. 1 for details. Moreover, stationary but spatially nonhomogeneous solutions bifurcate at the border  $C_E$  between  $D_S$  and  $D_U$  (see e.g. [15], [17]). Such solutions describe spatial patterns in mathematical models in biology (see e.g. [4], [11], [16]). In fact, in applications there is usually a positive (not zero) spatially constant stationary solution  $\bar{u}, \bar{v}$  and spatially nonhomogeneous stationary solutions bifurcate from  $\bar{u}, \bar{v}$ . However, this basic solution can be shifted to zero, which is done in our system. Let us note that any nontrivial solution of (1.6) with boundary conditions (1.7) or (1.3) is spatially nonhomogeneous due to the Dirichlet conditions on a part of the boundary.

In a series of papers (e.g. [1], [2], [6], [9], [12], [18]), an influence of unilateral conditions to this bifurcation was studied. (Usually only systems of activator-inhibitor type were discussed but in fact only the assumption (1.4) was used, i.e. the results were true also for systems of positive feedback type.) Roughly speaking, it was proved that if unilateral conditions are prescribed for the inhibitor  $v$  then, if an eigenfunction of the Laplacian satisfies a certain sign condition, bifurcation occurs even in the domain  $D_S$ , while if unilateral conditions are prescribed for the activator  $u$  then bifurcation is excluded in  $D_S$  and in some situations it is excluded even in  $C_E$ . However, the existence of a bifurcation in the last case has not been proved up to now, and is shown only in the current paper for the particular case  $n_2 = 0$ .

The goal of this paper is two-fold. One goal is to prove existence of bifurcation in  $D_U$  for the particular case of a nonlinearity only in the first equation and unilateral conditions for  $u$ . The other goal is to prove bifurcation in the domain  $D_S$  for the case of a nonlinearity only in the second equation and unilateral conditions for  $v$  even if all eigenfunctions of the Laplacian fail to satisfy the mentioned sign condition demanded in all previous papers. Both results are obtained by the same approach in a somewhat dual manner.

All previous results concerning bifurcation for our system with unilateral conditions for the inhibitor were based on topological methods. Namely, either on a certain homotopical joining of the variational inequality to the equation by a system of penalty problems and a transfer of the information about the existence of small nontrivial solutions from the equation to the inequality (e.g. [2], [6]), or on the direct use of the Leray-Schauder degree (a jump of the degree implies bifurcation, e.g. [18], [9]). As we already mentioned, for the proof it was always essential that certain eigenfunctions of the Laplacian satisfy a certain sign condition (in general, it must

be in the interior or in a certain pseudo-interior of the cone related to the corresponding variational inequality), see e.g. [1], [9]. In the present variational approach we need no such assumption, cf. Remark 5.2. The price for the use of the variational approach is that we consider only some particular situations, and that we get no information concerning the character of the set of bifurcating solutions. However, in a particular case of situations like in Examples 5.2, 5.3, our method combined with the results [8], [19] can give a bifurcation for the case of general  $n_1, n_2$  (Remark 3.2), and even the direction of bifurcation can be described.

In Section 2 we give an abstract formulation of our problem and summarize some facts necessary for the formulation of main results, which are given in an abstract form in Section 3 (Theorems 3.1 and 3.2). The proofs of the abstract results are given in Section 4. Section 5 is devoted to applications to unilateral boundary value problems; the first example covers the boundary conditions (1.3).

For the proof of existence of a bifurcation in  $D_U$ , we transfer for fixed  $d_2 > 0$  the problem to a single variational inequality and prove by a variational approach that there is a bifurcation for a suitable  $d_1$ . The proof of existence of a bifurcation in  $D_S$  works in a dual manner by interchanging the roles of  $d_1$  and  $d_2$ .

The idea to transfer our nonsymmetric problem for fixed  $d_2$  to a single variational inequality comes already from [12], but up to now it has been always used only for the proof of nonexistence of critical points (and consequently, nonexistence of bifurcation). The dual approach, i.e. to transfer the nonsymmetric problem for fixed  $d_1$  to a single variational inequality, was used in [5], also for the proof of nonexistence of bifurcation.

Unilateral boundary conditions can describe a certain regulation, e.g. by a unilateral membrane. Interpretation of the boundary conditions (1.3) (even in a more general form) and of unilateral conditions from Examples 5.2, 5.3 (the last section) is described e.g. in [1] and in Remark 5.3, respectively.

## 2. MOTIVATION OF ABSTRACT FORMULATION, GENERAL REMARKS

Let us assume that  $n$  is a continuous function satisfying (1.5) and that there exists  $c \in \mathbb{R}$  such that

$$(2.1) \quad |n(u)| \leq c(1 + |u|)^{q-1}$$

with some  $q > 2$  or  $2 < q < 2m/(m-2)$  in the case  $m \leq 2$  or  $m > 2$ , respectively (in the case  $m = 1$ , one can even formally put  $q = \infty$  and does not need to require (2.1)).

Let us introduce the real Hilbert space

$$(2.2) \quad \mathbb{H} = \{\varphi \in W^{1,2}(\Omega): \varphi = 0 \text{ on } \Gamma_D \text{ in the sense of traces}\}$$

and the closed convex cone  $K$  with its vertex at the origin in  $\mathbb{H}$ ,

$$K = \{\varphi \in H: \varphi \geq 0 \text{ on } \Gamma_U \text{ in the sense of traces}\}.$$

We equip  $\mathbb{H}$  with the scalar product

$$(2.3) \quad \langle u, \varphi \rangle = \int_{\Omega} \nabla u(x) \cdot \nabla \varphi(x) \, dx \quad (u, \varphi \in \mathbb{H})$$

and the corresponding norm  $\|\varphi\|^2 = \langle \varphi, \varphi \rangle$  which is equivalent to the usual Sobolev norm under the assumption (1.1), see e.g. [10]. Set

$$\langle U, W \rangle = \langle u, w \rangle + \langle v, z \rangle, \quad \|U\|^2 = \|u\|^2 + \|v\|^2 \text{ for } U = (u, v), \quad W = (w, z) \in \mathbb{H} \times \mathbb{H}.$$

Let us define operators  $A, N: \mathbb{H} \rightarrow \mathbb{H}$  by

$$(2.4) \quad \langle Au, \varphi \rangle = \int_{\Omega} u(x)\varphi(x) \, dx \quad \text{for all } u, \varphi \in \mathbb{H},$$

$$(2.5) \quad \langle N(u), \varphi \rangle = \int_{\Omega} n(u(x))\varphi(x) \, dx \quad \text{for all } u, \varphi \in \mathbb{H}.$$

It follows from the compactness of the embedding  $\mathbb{H} \hookrightarrow L^q(\Omega)$  and the continuity of the Nemyckij operator of  $L^q(\Omega)$  into  $L^{q^*}(\Omega)$ ,  $1/q + 1/q^* = 1$  (see e.g. [10]) that under the assumption (2.1)

$$(2.6) \quad A \text{ is linear, symmetric, positive and compact,}$$

$$(2.7) \quad N \text{ is nonlinear, continuous and compact.}$$

Furthermore, under the conditions (1.5) and (2.1)

$$(2.8) \quad N \text{ is Fréchet differentiable at } 0, \quad N(0) = 0, \quad N'(0) = 0,$$

see e.g. [3]. Moreover, let us introduce the functional  $G_N: \mathbb{H} \rightarrow \mathbb{R}$  by

$$G_N(u) = \int_{\Omega} \int_0^{u(x)} n(s) \, ds \, dx.$$

Under the assumptions (2.1), this functional is well defined, Fréchet differentiable and we have

$$(2.9) \quad G'_N(u) = N(u),$$

i.e.  $G_N$  is a potential of the operator  $N$ .

Now, we introduce a weak solution of the problem (1.2), (1.3) or (1.2), (1.7) as a couple  $(u, v)$  satisfying the variational inequality

$$(2.10) \quad \begin{cases} u \in K, v \in \mathbb{H}, \\ \langle d_1 u - b_{11} A u - b_{12} A v - N(u), \varphi - u \rangle \geq 0 \quad \text{for all } \varphi \in K, \\ d_2 v - b_{21} A u - b_{22} A v = 0, \end{cases}$$

or the equations

$$(2.11) \quad \begin{cases} u, v \in \mathbb{H}, \\ d_1 u - b_{11} A u - b_{12} A v - N(u) = 0, \\ d_2 v - b_{21} A u - b_{22} A v = 0, \end{cases}$$

respectively. The weak formulation of the linearized system

$$(2.12) \quad d_1 \Delta u + b_{11} u + b_{12} v = 0, \quad d_2 \Delta v + b_{21} u + b_{22} v = 0 \quad \text{in } \Omega$$

with (1.3) or with (1.7) is

$$(2.13) \quad \begin{cases} u \in K, v \in \mathbb{H}, \\ \langle d_1 u - b_{11} A u - b_{12} A v, \varphi - u \rangle \geq 0 \quad \text{for all } \varphi \in K, \\ d_2 v - b_{21} A u - b_{22} A v = 0, \end{cases}$$

or

$$(2.14) \quad \begin{cases} u, v \in \mathbb{H}, \\ d_1 u - b_{11} A u - b_{12} A v = 0, \\ d_2 v - b_{21} A u - b_{22} A v = 0, \end{cases}$$

respectively.

In parallel, we will consider the following dual situation with the variational inequality for  $v$  and nonlinearity dependent only on  $v$ , that means

$$(2.15) \quad \begin{cases} u \in \mathbb{H}, v \in K, \\ d_1 u - b_{11} A u - b_{12} A v = 0, \\ \langle d_2 v - b_{21} A u - b_{22} A v - N(v), \varphi - u \rangle \geq 0 \quad \text{for all } \varphi \in K, \end{cases}$$

the corresponding system of equations

$$(2.16) \quad \begin{cases} u, v \in \mathbb{H}, \\ d_1 u - b_{11} A u - b_{12} A v = 0, \\ d_2 v - b_{21} A u - b_{22} A v - N(v) = 0, \end{cases}$$

and the variational inequality with linearized operators

$$(2.17) \quad \begin{cases} u \in \mathbb{H}, v \in K, \\ d_1 u - b_{11} A u - b_{12} A v = 0, \\ \langle d_2 v - b_{21} A u - b_{22} A v, \varphi - u \rangle \geq 0 \quad \text{for all } \varphi \in K. \end{cases}$$

In fact, we will formulate our main results in the abstract form for the variational inequalities (2.10) and (2.15) in a general real Hilbert space with a closed convex cone  $K$  with its vertex at the origin in  $\mathbb{H}$ , and with operators  $A, N$  satisfying (2.6), (2.7), (2.8), with  $N$  having a potential  $G_N$ , i.e. (2.9) holds.

In the sequel, with the exception of the last section,  $\mathbb{H}, K, A, N$  have this general meaning if nothing else is mentioned.

In some of the papers mentioned in Section 1, bifurcations with respect to  $(d_1, d_2)$  along general curves in  $\mathbb{R}_+^2$  were studied. Our variational approach enables us to consider only particular cases of bifurcation in the direction  $d_1$  or in the direction  $d_2$  with fixed  $d_2$  or  $d_1$ , respectively.

**Definition 2.1.** A parameter  $d_1$  is a *bifurcation point* of (2.10) or (2.11) with fixed  $d_2 = d_2^0$  if in any neighborhood of  $(d_1, 0, 0)$  in  $\mathbb{R} \times \mathbb{H} \times \mathbb{H}$  there is  $(\tilde{d}_1, u, v)$  with  $(u, v) \neq (0, 0)$  such that  $(\tilde{d}_1, d_2^0, u, v)$  satisfies (2.10) or (2.11), respectively.

A parameter  $d_2$  is a *bifurcation point* of (2.15) or (2.16) with fixed  $d_1 = d_1^0$  if in any neighborhood of  $(d_2, 0, 0)$  in  $\mathbb{R} \times \mathbb{H} \times \mathbb{H}$  there is  $(\tilde{d}_2, u, v)$  with  $(u, v) \neq (0, 0)$  such that  $(d_1^0, \tilde{d}_2, u, v)$  satisfies (2.15) or (2.16), respectively.

By a bifurcation point of the problem (1.2), (1.3) we mean always a bifurcation point of the problem (2.10) with  $A, N$  from (2.4), (2.5). Analogously for the other problems considered.

A *critical point* of the problem (2.13), (2.17) or (2.14) is a parameter  $d = (d_1, d_2) \in \mathbb{R}_+^2$  for which the corresponding system has a solution  $(u, v) \neq (0, 0)$ . By a critical point of the problem (2.12), (1.3) we mean always a critical point of the problem (2.13) with  $A, N$  from (2.4), (2.5). Analogously for the other problems considered.

**Remark 2.1.** If  $d_1$  is a bifurcation point of (2.10) with fixed  $d_2 = d_2^0$  or  $d_2$  is a bifurcation point of (2.15) with fixed  $d_1 = d_1^0$  then  $(d_1, d_2^0)$  or  $(d_1^0, d_2)$  is a critical point of (2.13) or (2.17), respectively, cf. e.g. [2]. Of course, analogously for (2.11) and (2.16).

**Remark 2.2.** Let us consider the eigenvalue problem

$$(2.18) \quad \begin{aligned} d_1 \Delta u + b_{11} u + b_{12} v &= \lambda u \\ d_2 \Delta v + b_{21} u + b_{22} v &= \lambda v \end{aligned} \quad \text{in } \Omega$$



with the boundary conditions (1.7). Let us recall that if  $\operatorname{Re} \lambda \leq -\varepsilon < 0$  for all eigenvalues of the problem (2.18), (1.7) then the trivial solution of (1.6), (1.7) is linearly stable, and if there is at least one eigenvalue of (2.18), (1.7) satisfying  $\operatorname{Re} \lambda > 0$  then the trivial solution of (1.6), (1.7) is linearly unstable (see e.g. [20]). Hence, the definition of the domain  $D_S$  and  $D_U$  of stability and instability below (related to the classical problem (1.6), (1.7)) is natural due to Proposition 2.1 below.

**Notation 2.1.** Let us denote by  $0 < \kappa_1 < \kappa_2 \leq \kappa_3 \leq \dots$  the characteristic values (i.e. reciprocals of eigenvalues) of the operator  $A$ , counted according to their multiplicity. Furthermore, let  $e_j$  ( $j = 1, 2, \dots$ ) be a corresponding orthonormal system of eigenvectors. With each  $\kappa_j$ , we associate the hyperbola

$$\tilde{C}_j := \left\{ d = (d_1, d_2) \in \mathbb{R}^2 : d_2 = \frac{b_{12}b_{21}/\kappa_j^2}{d_1 - b_{11}/\kappa_j} + \frac{b_{22}}{\kappa_j} \right\}$$

and denote by  $C_j$  the part of  $\tilde{C}_j$  lying in the positive quadrant  $\mathbb{R}_+^2$ , i.e.

$$C_j := \left\{ d = (d_1, d_2) \in \mathbb{R}_+^2 : d_2 = \frac{b_{12}b_{21}/\kappa_j^2}{d_1 - b_{11}/\kappa_j} + \frac{b_{22}}{\kappa_j} \right\}.$$

We denote by  $C_E$  the envelope of  $C_j$  ( $j = 1, 2, \dots$ ) and define the domain of stability

$$D_S := \{d \in \mathbb{R}_+^2 : d \text{ lies to the right from } C_E, \text{ i.e. from all } C_j, j = 1, 2, \dots\}$$

and the domain of instability

$$D_U := \{d \in \mathbb{R}_+^2 : d \text{ lies to the left from } C_E, \text{ i.e. from at least one } C_j\}$$

(see Fig. 1).




Figure 1. The system of hyperbolas  $C_j$ , their asymptotes  $x_j$ , their intersection  $y_j$  with the axis  $d_2$ , domains of stability  $D_S$  (to the right from the envelope  $C_E$ ) and instability  $D_U$  (to the left from  $C_E$ ).

For any  $j = 1, 2, \dots$ , we will denote by  $y_j := \det B / \kappa_j b_{11}$  the  $d_2$ -coordinate of the intersection point of  $C_j$  with the  $d_2$ -axis, that means

$$y_j \text{ is the positive real with } (0, y_j) \in C_j,$$

and by  $x_j := b_{11} / \kappa_j$  the  $d_1$ -coordinate of the vertical asymptote of  $C_j$ .

For  $(d_1, d_2) \in \bigcup_j \tilde{C}_j$ , it will be convenient to define the index set

$$I(d_1, d_2) = \{j: (d_1, d_2) \in \tilde{C}_j\}.$$

Note that two hyperbolas  $C_j, C_k$  are either identical or intersect at exactly one point, and no more than two different hyperbolas intersect at any point. Hence, concerning the number of elements  $|I(d_1, d_2)|$  there are essentially only two cases: If  $(d_1, d_2) \in C_j$  lies on only one hyperbola, then  $|I(d_1, d_2)|$  is the multiplicity of the characteristic value  $\kappa_j$ . If  $(d_1, d_2)$  lies on the intersection points of two hyperbolas  $C_j \neq C_k$  then  $|I(d_1, d_2)|$  is the sum of the multiplicities of  $\kappa_j$  and of  $\kappa_k$ .

**Remark 2.3.** In the case of the operator  $A$  from (2.4),  $\kappa_j$  is a characteristic value of  $A$  if and only if it is an eigenvalue of the boundary value problem

$$(2.19) \quad \begin{cases} -\Delta u = \kappa_j u & \text{in } \Omega, \\ u = 0 \text{ on } \Gamma_D, \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega \setminus \Gamma_D, \end{cases}$$

and the corresponding eigenvectors of  $A$  coincide with the eigenfunctions of (2.19).

The weak formulation of the eigenvalue problem (2.18), (1.7) is

$$(2.20) \quad \begin{aligned} d_1 u - b_{11} A u - b_{12} A v &= \lambda A u, \\ d_2 v - b_{21} A u - b_{22} A v &= \lambda A v \end{aligned}$$

with the operator  $A$  defined by (2.4). In this particular case the eigenvalues and the corresponding eigenfunctions of (2.20) and of (2.18), (1.7) coincide.

**Proposition 2.1.** *Let  $\mathbb{H}$  be a real Hilbert space and let  $A: \mathbb{H} \rightarrow \mathbb{H}$  be an operator satisfying (2.6). Assume that (1.4) is fulfilled. Then  $\bigcup_{j=1}^{\infty} \tilde{C}_j$  is the set of all critical points of the problem (2.14), and for  $(d_1, d_2) \in \bigcup_{j=1}^{\infty} \tilde{C}_j$  we have*

$$\begin{aligned} & \left\{ \begin{pmatrix} u \\ v \end{pmatrix} \in \mathbb{H} \times \mathbb{H}: (2.14) \text{ is fulfilled} \right\} \\ &= \text{span} \left\{ \begin{pmatrix} e_j \\ b_{21}(d_2 \kappa_j - b_{22})^{-1} e_j \end{pmatrix} : j \in I(d_1, d_2) \right\}. \end{aligned}$$

If  $d \in D_S$  then there is  $\varepsilon > 0$  such that  $\operatorname{Re} \lambda < -\varepsilon$  for all eigenvalues of (2.20) and if  $d \in D_U$  then there exists at least one positive eigenvalue of (2.20).

**Proof.** For a particular case of the reaction-diffusion system in one space dimension see e.g. [15], [17], for the general case see e.g. [2], [9].  $\square$

### 3. GENERAL RESULTS

In this section we will consider a general real Hilbert space  $\mathbb{H}$  with the scalar product  $\langle \cdot, \cdot \rangle$  and a closed convex cone  $K$  with its vertex at the origin in  $\mathbb{H}$ . We will discuss the variational inequalities (2.10) and (2.15) with general operators  $A, N: \mathbb{H} \rightarrow \mathbb{H}$  satisfying (2.6), (2.7), (2.8) with  $N$  having a potential  $G_N$ , i.e. (2.9) holds. The condition (1.4) will be always assumed. The proofs of the results described here will be given in Section 4.

**Theorem 3.1.** *Let  $d_2^0 > 0$  be such that there are  $\xi_j \in \mathbb{R}$  with*

$$(3.1) \quad \sum_{j=1}^{\infty} \xi_j e_j \in K \quad \text{and} \quad \sum_{j=1}^{\infty} \frac{d_2^0 b_{11} - \kappa_j^{-1} \det B}{d_2^0 \kappa_j - b_{22}} \xi_j^2 > 0.$$

Then the value

$$(3.2) \quad d_1^0 := \max \left\{ \sum_{j=1}^{\infty} \frac{d_2^0 b_{11} - \kappa_j^{-1} \det B}{d_2^0 \kappa_j - b_{22}} \xi_j^2 \middle/ \sum_{j=1}^{\infty} \xi_j^2 : (\xi_j) \in \ell_2 \setminus \{0\} \text{ and } \sum_{j=1}^{\infty} \xi_j e_j \in K \right\}$$

is the largest bifurcation point of the problem (2.10) with fixed  $d_2 = d_2^0$ . Moreover,  $(d_1^0, d_2^0) \in D_U \cup C_E$ , and the case  $(d_1^0, d_2^0) \in C_E$  occurs only if there is  $(u, v) \neq (0, 0)$  satisfying (2.14) with  $(d_1, d_2) = (d_1^0, d_2^0)$  and  $u \in K$ . In the latter case, these  $(u, v)$  are exactly the nonzero solutions of (2.13).

Let us emphasize that in the case  $(d_1^0, d_2^0) \in C_E$  in Theorem 3.1 it can happen that the bifurcating solutions of (2.10) are not solutions of (2.11) because the bifurcating solutions of (2.11) (if they exist) need not satisfy  $u \in K$ .

For the proof in Section 4 we will introduce an operator  $S_{2, d_2^0}$  (see Proposition 4.2) such that the assumption (3.1) is equivalent to the existence of  $u \in K$  satisfying  $\langle S_{2, d_2^0} u, u \rangle > 0$ , and (3.2) means

$$(3.3) \quad d_1^0 = \max_{u \in K \setminus \{0\}} \frac{\langle S_{2, d_2^0} u, u \rangle}{\|u\|^2}.$$

**Corollary 3.1.** *Let  $d_2^0 > 0$  be such that the assumptions of Theorem 3.1 are fulfilled. Let  $d_1^{\max} > 0$  denote the unique number with  $(d_1^{\max}, d_2^0) \in C_E$ . If  $(d_1, d_2) = (d_1^{\max}, d_2^0)$  satisfies*

$$(3.4) \quad \sum_{j \in I(d_1, d_2)} \alpha_j e_j \notin K \quad \text{for all nontrivial } (\alpha_j)_{j \in I(d_1, d_2)},$$

then  $(d_1, d_2^0) \in D_U$  for all bifurcation points  $d_1$  of (2.10) with fixed  $d_2 = d_2^0$ .

*Proof.* The assumption (3.4) means in view of Proposition 2.1 that for  $(d_1, d_2) = (d_1^{\max}, d_2^0)$  there is no solution  $(u, v) \neq (0, 0)$  of (2.14) with  $u \in K$ . Hence, it follows from the last assertion of Theorem 3.1 that the case  $d_1^0 = d_1^{\max}$  is excluded, and therefore  $d_1^0 < d_1^{\max}$ .  $\square$

Theorem 3.1 implies in particular that problem (2.10) has at least one bifurcation point with fixed  $d_2 = d_2^0$  for every  $d_2^0 > y_1$ :

*Remark 3.1.* Hypothesis (3.1) is satisfied for  $d_2^0 > y_1$  if there is  $u \in K \setminus \{0\}$ . More generally, condition (3.1) is satisfied for  $d_2^0 > y_{j_0}$  if

$$(3.5) \quad \text{there is } u \in K \setminus \{0\} \quad \text{with } u \perp \{e_j : j < j_0\}.$$

Indeed, we can write  $u$  uniquely in the form  $u = \sum_{j=1}^{\infty} \xi_j e_j$ , and in view of  $u \perp \{e_j : j < j_0\}$ , we must have  $\xi_j = 0$  for  $j < j_0$ . The coefficients

$$c_j(d_2) := \frac{d_2 b_{11} - \kappa_j^{-1} \det B}{d_2 \kappa_j - b_{22}}$$

in (3.1) are strictly increasing in  $d_2$  on  $[0, \infty)$ , because  $c'_j(d) > 0$  for  $d > 0$  under the assumption (1.4), and  $y_{j_0} \geq y_j$  for  $j \geq j_0$ . Hence,

$$c_j(d_2^0) > c_j(y_{j_0}) \geq c_j(y_j) = 0 \quad \text{for } j \geq j_0.$$

The assumption (3.5) as well as the conditions given below will be verified in concrete examples in Section 5.

*Remark 3.2.* In particular, if  $N = 0$  then Theorem 3.1 yields the existence of a critical point of problem (2.13). In the case of particular variational inequalities (e.g. when the cone  $K$  is given by a finite number of isolated obstacles as in Examples 5.2, 5.3 in the last section), such a critical point is also a bifurcation point even for our system with general  $N_1, N_2$  (not only  $N_2 = 0$ ) if certain simplicity assumptions are fulfilled. Moreover, bifurcating solutions form a smooth branch if  $N_1, N_2$  are smooth, see [19]. In this case also the bifurcation direction can be described, see [8].

The following result is the “dual” version of Theorem 3.1 for problem (2.15).

**Theorem 3.2.** *Let  $d_1^0 > 0$  satisfy  $d_1^0 \notin \{x_j : j = 1, 2, \dots\}$ . Assume that there are  $\xi_j \in \mathbb{R}$  with*

$$(3.6) \quad \sum_{j=1}^{\infty} \xi_j e_j \in K \quad \text{and} \quad \sum_{j=1}^{\infty} \frac{\kappa_j^{-1} \det B - d_1^0 b_{22}}{b_{11} - d_1^0 \kappa_j} \xi_j^2 > 0.$$

Then the value

$$(3.7) \quad d_2^0 := \max \left\{ \sum_{j=1}^{\infty} \frac{\kappa_j^{-1} \det B - d_1^0 b_{22}}{b_{11} - d_1^0 \kappa_j} \xi_j^2 \middle/ \sum_{j=1}^{\infty} \xi_j^2 : (\xi_j) \in \ell_2 \setminus \{0\} \text{ and } \sum_{j=1}^{\infty} \xi_j e_j \in K \right\}$$

is the largest bifurcation point of problem (2.15) with fixed  $d_1 = d_1^0$ . Moreover,  $d_1^0 < x_1$  and  $d_2^0 \leq d_2^{\max} := \max \left\{ d_2 > 0 : (d_1^0, d_2) \in \bigcup_j C_j \right\}$ , and the case  $d_2^0 = d_2^{\max}$  occurs only if there is a solution  $(u, v) \neq (0, 0)$  of (2.14) with  $(d_1, d_2) = (d_1^0, d_2^0)$  and  $v \in K$ . In the latter case, these  $(u, v)$  are exactly the nonzero solutions of (2.17).

**Corollary 3.2.** *Let  $d_1^0 > 0$  be such that the assumptions of Theorem 3.2 are fulfilled (and thus  $d_1^0 < x_1$ ). Let  $d_2^{\max} = \max \left\{ d_2 : (d_1^0, d_2) \in \bigcup_j C_j \right\}$ . If (3.4) holds with  $(d_1, d_2) = (d_1^0, d_2^{\max})$ , then  $d_2^0 < d_2^{\max}$  in Theorem 3.2.*

*In particular, if  $d_1^0 \geq x_2$  and (3.4) holds with  $(d_1, d_2) = (d_1^0, d_2^{\max})$ , then  $(d_1^0, d_2^0) \in D_S$ .*

**Remark 3.3.** Hypothesis (3.6) is satisfied for  $d_1^0 \in (0, x_{j_0})$  if

$$(3.8) \quad \text{there is } u \in K \setminus \{0\} \text{ with } u \in \text{span}\{e_j : j = 1, \dots, j_0\}.$$

Indeed, writing  $u = \sum_{j=1}^{\infty} \xi_j e_j$  with  $\xi_j = 0$  for  $j > j_0$ , it suffices to observe that the coefficients in (3.6) are positive for  $j \leq j_0$ , because the nominator is positive due to the assumption (1.4), and the denominator is positive in view of  $d_1^0 < x_{j_0} \leq x_j = \kappa_j^{-1} b_{11}$ .

Remark 3.3 is “dual” to Remark 3.1, nevertheless condition (3.8) is much more restrictive than (3.5) because the orthogonal complement to  $\{e_j : j < j_0\}$  is infinite-dimensional while the space  $\text{span}\{e_j : j \leq j_0\}$  has only a finite dimension. In particular, while Remark 3.1 implies that Theorem 3.1 applies for at least some  $d_2^0 > 0$  (if  $K \neq \{0\}$ ), Remark 3.3 does not give an analogous consequence for Theorem 3.2. However, this consequence is also true for Theorem 3.2, but it requires a different

argument. Indeed, if there is  $u \in K \setminus \{0\}$  then the completeness of the basis  $(e_j)$  implies that there is at least one  $j_0$  with  $\langle u, e_{j_0} \rangle \neq 0$ . For these  $j_0$ , we can use the following observation:

**Remark 3.4.** Let  $j_0 \in \{1, 2, \dots\}$  be such that

$$(3.9) \quad \text{there is } u \in K \text{ with } \langle u, e_{j_0} \rangle \neq 0.$$

Then there is  $\varepsilon_{j_0} > 0$  such that hypothesis (3.6) holds for every  $d_1^0 \in [x_{j_0} - \varepsilon_{j_0}, x_{j_0})$ .

Indeed, writing  $u = \sum_{j=1}^{\infty} \xi_j e_j$ , we have  $\xi_{j_0} = \langle u, e_{j_0} \rangle \neq 0$ . Put  $I := \{j : \kappa_j = \kappa_{j_0}\}$ .

The corresponding series in (3.6) has the form  $S_1(d_1) + S_2(d_1)$  with

$$S_1(d_1) := \sum_{j \in I} \frac{\kappa_j^{-1} \det B - d_1 b_{22}}{b_{11} - d_1 \kappa_j} \xi_j^2, \quad S_2(d_1) := \sum_{\substack{j=1 \\ j \notin I}}^{\infty} \frac{\kappa_j^{-1} \det B - d_1 b_{22}}{b_{11} - d_1 \kappa_j} \xi_j^2.$$

Now observe that  $S_2(d_1)$  remains bounded for  $d_1$  close to  $\kappa_{j_0}^{-1} b_{11} = x_{j_0}$ , and that  $S_1(d_1) \rightarrow \infty$  as  $d_1$  approaches  $x_{j_0}$  from the left due to (1.4). Hence, for all  $d_1 < x_{j_0}$  sufficiently close to  $x_{j_0}$  the series in (3.6) is strictly positive.

#### 4. PROOF OF THE MAIN RESULTS

The proofs are based on the following well-known variational principle. Given a mapping  $G: \mathbb{H} \rightarrow \mathbb{H}$  with  $G(0) = 0$ , we call  $\lambda_0 \in \mathbb{R}$  a bifurcation point of the variational inequality

$$(4.1) \quad u \in K, \quad \langle \lambda u - G(u), \varphi - u \rangle \geq 0 \text{ for all } \varphi \in K,$$

if every neighborhood of  $(\lambda_0, 0) \in \mathbb{R} \times \mathbb{H}$  contains  $(\lambda, u)$  satisfying (4.1) with  $u \neq 0$ . If  $S: \mathbb{H} \rightarrow \mathbb{H}$  is linear, we say that  $\lambda$  is an eigenvalue of

$$(4.2) \quad u \in K, \quad \langle \lambda u - Su, \varphi - u \rangle \geq 0 \text{ for all } \varphi \in K,$$

if (4.2) has a solution  $u \neq 0$ ; we call each such  $u$  a corresponding eigenvector.

**Proposition 4.1.** *Let  $G: \mathbb{H} \rightarrow \mathbb{H}$  be a compact potential operator satisfying  $G(0) = 0$ . Suppose that  $G$  is Fréchet differentiable at 0,  $S := G'(0)$ , and that there is  $u_0 \in K$  with  $\langle Su_0, u_0 \rangle > 0$ . Then the maximum*

$$(4.3) \quad \lambda_0 := \max_{u \in K \setminus \{0\}} \frac{\langle Su, u \rangle}{\|u\|^2}$$

*exists and is the largest bifurcation point of (4.1) and the largest eigenvalue of (4.2). Moreover, the eigenvectors of (4.2) are exactly those  $u$  for which the maximum in (4.3) is assumed.*

*Proof.* The result follows from [14, Theorem 1] or [23, Theorem 64.A], where it is formulated in terms of bilinear forms. Let us verify the assumptions of the last theorem. Let  $g$  be a potential of  $G$  with  $g(0) = 0$ . The compactness of  $g' = G$  implies the weak sequential continuity of  $g$  and the compactness of  $S$ , i.e. also the compactness of the bilinear form  $b(u, u) := \langle Su, u \rangle$  (see e.g. [22, Corollary 41.9] and [21, Corollary 21.33]). Since  $g$  is twice Fréchet differentiable at 0 with  $g(0) = 0$  and  $g'(0) = 0$ , we conclude from [23, Corollary 64.4] that all hypotheses of [23, Theorem 64.A] are satisfied with  $F(u) = \|u\|^2$  and  $a(u, v) := \langle u, v \rangle$  (and  $G$  in place of  $g$ ). The latter implies that the maximum in (4.3) exists and is the largest bifurcation point of (4.1) and the largest eigenvalue of (4.2). The last assertion of Proposition 4.1 follows from the last statement of [14, Theorem 1 (c)].  $\square$

The crucial hypothesis of Proposition 4.1 is of course that the operator  $g$  needs to have a potential, which is not the case for the operators occurring in (2.10) and (2.15) if we interpret the equation in an obvious manner on the product space  $\mathbb{H} \times \mathbb{H}$ . However, we can equivalently rewrite the equations in the space  $\mathbb{H}$  in the required form when we consider only  $d_1$  or only  $d_2$  as a bifurcation parameter (cf. [12] for this idea).

**Proposition 4.2.** *For fixed  $d_2 > 0$  the problem (2.10) is equivalent to the problem*

$$(4.4) \quad u \in K, \quad \langle d_1 u - S_{2,d_2} u - N(u), \varphi - u \rangle \geq 0 \quad \text{for all } \varphi \in K,$$

$$(4.5) \quad v = (d_2 I - b_{22} A)^{-1} b_{21} A u,$$

with the operator

$$S_{2,d_2} := b_{11} A u + b_{12} A (d_2 I - b_{22} A)^{-1} b_{21} A.$$

*Proof.* Since  $d_2/b_{22} < 0$  by (1.4) and the operator  $A$  has only positive eigenvalues, the inverse in (4.5) exists. Hence, we can uniquely solve the last equation of (2.10) for  $v$ , which is equivalent to (4.5). Inserting this value into the inequality of (2.10), we obtain (4.4).  $\square$

Interchanging the roles of  $d_1$  and  $d_2$  in Proposition 4.2, we can similarly rewrite the problem (2.15):

**Proposition 4.3.** *For fixed  $d_1 > 0$  with  $d_1 \notin \{x_j : j = 1, 2, \dots\}$  the problem (2.15) is equivalent to the problem*

$$(4.6) \quad v \in K, \quad \langle d_2 v - S_{1,d_1} v - N(v), \varphi - v \rangle \geq 0 \quad \text{for all } \varphi \in K,$$

$$(4.7) \quad u = (d_1 I - b_{11} A)^{-1} b_{12} A v,$$

with the operator

$$S_{1,d_1} := b_{21}A(d_1I - b_{11}A)^{-1}b_{12}A + b_{22}A.$$

**P r o o f.** Since  $d_1 \neq x_j = b_{11}\kappa_j^{-1}$ , we have  $d_1 \neq b_{11}\lambda$  for every eigenvalue  $\lambda$  of  $A$ , and so the inverse in the last equation exists. Hence, we can solve the first equation in (2.15) for  $u$  and obtain (4.7). Inserting this into the second equation, we obtain (4.6).  $\square$

The same calculation shows the corresponding statements for (2.14):

**Proposition 4.4.** *With the notation of Propositions 4.2, 4.3 the problem (2.14) is equivalent to the equation  $S_{2,d_2}u = d_1u$  with (4.5), and also equivalent to  $S_{1,d_1}u = d_2u$  with (4.7).*

We recall the well-known variational characterization of the largest eigenvalue of a symmetric compact operator, and the characterization of the corresponding eigenvectors.

**Proposition 4.5.** *Let  $S: \mathbb{H} \rightarrow \mathbb{H}$  be linear, symmetric, and compact. Then the largest eigenvalue of  $S$  is given by*

$$(4.8) \quad \lambda_0 := \max_{u \in \mathbb{H} \setminus \{0\}} \frac{\langle Su, u \rangle}{\|u\|^2},$$

and this maximum is attained exactly in all the corresponding eigenvectors.

**Corollary 4.1.** *If the number  $\lambda_0$  of Proposition 4.1 is simultaneously the largest eigenvalue of the operator  $S$ , then there is  $u_0 \in K \setminus \{0\}$  with  $Su_0 = \lambda_0u_0$ .*

**P r o o f.** If  $u_0$  is an eigenvector of (4.2) corresponding to  $\lambda_0$  then the maximum in (4.3) is attained at  $u_0$  by Proposition 4.1. If  $\lambda_0$  is the largest eigenvalue of the operator  $S$  then Proposition 4.5 implies that also the maximum in (4.8) is assumed at  $u_0$  and  $Su_0 = \lambda_0u_0$ .  $\square$

**P r o o f** of Theorem 3.1. In view of Proposition 4.2, we only have to study bifurcation points of the problem (4.4) (with fixed  $d_2 = d_2^0$ ). However, this problem has exactly the form (4.1) with  $\lambda = d_1$ ,  $G(u) = S_{2,d_2}u + N(u)$ . Using operator calculus, we can write  $S_{2,d_2^0} = f_{2,d_2^0}(A)$ , where the function  $f_{2,d_2^0}$  is defined on the spectrum of  $A$  by

$$f_{2,d_2^0}(\lambda) := b_{11}\lambda + \frac{b_{12}b_{21}\lambda^2}{d_2^0 - b_{22}\lambda} = \frac{d_2^0b_{11}\lambda - (\det B)\lambda^2}{d_2^0 - b_{22}\lambda}.$$



Note that  $G$  has a potential, since  $S := S_{2,d_2^0} = f_{2,d_2^0}(A)$  is symmetric and since  $N$  has a potential  $G_N$  by hypothesis. Moreover, (2.8) implies  $G(0) = 0$  and  $G'(0) = S$ .

Every  $u \in K$  can be uniquely written in the form  $u = \sum_{j=1}^{\infty} \xi_j e_j$  with  $\xi_j \in \mathbb{R}$ . By Parseval's identity, we have  $(\xi_j) \in \ell_2$ ,  $\|u\|^2 = \sum_{j=1}^{\infty} \xi_j^2$ , and by the rules of calculus for symmetric operators, we have

$$\langle S_{2,d_2^0} u, u \rangle = \langle f_{2,d_2^0}(A)u, u \rangle = \left\langle \sum_{j=1}^{\infty} f_{2,d_2^0}(\kappa_j^{-1}) \xi_j e_j, \sum_{i=1}^{\infty} \xi_i e_i \right\rangle = \sum_{j=1}^{\infty} f_{2,d_2^0}(\kappa_j^{-1}) \xi_j^2,$$

which is exactly the series occurring in (3.1) and (3.2). In particular, (3.1) means that  $\langle S_{2,d_2^0} u_0, u_0 \rangle > 0$  for  $u_0 = \sum_{j=1}^{\infty} \xi_j e_j$ , and (3.2) means exactly (3.3). Hence,  $d_1^0$  is the largest bifurcation point of (2.10) with fixed  $d_2 = d_2^0$  by Proposition 4.1.

Propositions 2.1 and 4.4 imply that the eigenvalues of  $S = S_{2,d_2^0}$  are exactly those  $d_1$  with  $(d_1, d_2^0) \in \bigcup_j C_j$ . Hence, putting  $d_1^{\max} = \max\{d_1 \in \mathbb{R} : (d_1, d_2^0) \in \bigcup_j \tilde{C}_j\}$ , we have  $d_1^0 \leq d_1^{\max}$ , i.e.  $(d_1^0, d_2^0) \in D_U \cup C_E$ , and the equality  $d_1^0 = d_1^{\max}$  (i.e.  $(d_1^0, d_2^0) \in C_E$ ) holds if and only if  $d_1^0$  is the maximal eigenvalue of  $S$ . Corollary 4.1 and Proposition 4.2 used for  $N = 0$  imply that if this is true then all solutions of (2.13) (with  $(d_1, d_2) = (d_1^0, d_2^0)$ ) satisfy also (2.14), and the last assertions of Theorem 3.1 follow.  $\square$

**Proof of Theorem 3.2.** Using operator calculus, we can write  $S_{1,d_1^0} = f_{1,d_1^0}(A)$  with the function  $f_{1,d_1^0}$  defined on the spectrum of  $A$  by

$$f_{1,d_1^0}(\lambda) := \frac{b_{12}b_{21}\lambda^2}{d_1^0 - b_{11}\lambda} + b_{22}\lambda = \frac{(\det B)\lambda^2 - d_1^0 b_{22}\lambda}{b_{11}\lambda - d_1^0}.$$

Then the proof is analogous to the proof of Theorem 3.1 by applying Proposition 4.1 to the problem (4.6), and observing that for  $u = \sum_{j=1}^{\infty} \xi_j e_j \in K$  the expression

$$\langle S_{1,d_1^0} u, u \rangle = \sum_{j=1}^{\infty} f_{1,d_1^0}(\kappa_j^{-1}) \xi_j^2$$

is exactly the series occurring in (3.6) and (3.7). By an analogous reasoning as in the proof of Theorem 3.1, we obtain  $d_2^0 \leq \tilde{d}_2^{\max} := \max\{d_2 \in \mathbb{R} : (d_1^0, d_2) \in \bigcup_j \tilde{C}_j\}$  and that  $\tilde{d}_2^{\max}$  is the maximal eigenvalue of  $S := S_{1,d_1^0}$ . Since  $d_2^0 > 0$ , we must have

$\tilde{d}_2^{\max} > 0$ . The operator  $S_{1,d_1^0}$  has only negative eigenvalues if  $d_1^0 \geq x_1$  due to the form of the hyperbolas  $C_n$  and Proposition 4.4, which in turn implies  $d_1^0 < x_1$  and  $\tilde{d}_2^{\max} = \max\{d_2 > 0: (d_1^0, d_2) \in \bigcup_j C_j\} = d_2^{\max}$ . The proof of the last assertion is again similar to that of the last assertion of Theorem 3.1.  $\square$

## 5. APPLICATION TO UNILATERAL BOUNDARY VALUE PROBLEMS

Throughout this section, we consider a domain  $\Omega$  as in the introduction, a function  $n$  satisfying (1.5) and (2.1), the Hilbert space  $\mathbb{H}$  from (2.2) with the scalar product (2.3) and the corresponding norm  $\|\cdot\|$ , and the operators defined by (2.4), (2.5). The condition (1.4) will be always assumed. The problem (2.11), (2.14), or (2.20) is a weak formulation of (1.2), (2.12), or (2.18), respectively, with boundary conditions (1.7). Similarly, (2.16) is the weak formulation of the problem

$$(5.1) \quad \begin{aligned} d_1 \Delta u + b_{11}u + b_{12}v &= 0 \\ d_2 \Delta v + b_{21}u + b_{22}v + n(v) &= 0 \end{aligned} \quad \text{in } \Omega$$

with boundary conditions (1.7). The characteristic values and eigenvectors  $\kappa_j$  and  $e_j$  of the operator  $A$  are the eigenvalues and eigenfunctions of the problem (2.19), see Section 2. Hence, Proposition 2.1 remains valid if we replace (2.14) by (2.12), (1.7), and (2.20) by (2.18), (1.7). Speaking about solutions of boundary value problems we have always in mind weak solutions.

We will apply our abstract results to concrete choices of the cone  $K$  corresponding to different unilateral boundary conditions. As we have already mentioned in Section 2, the conditions (2.6), (2.7), (2.8) are automatically fulfilled and need not be repeated.

**Example 5.1.** Suppose that  $\Gamma_D, \Gamma_N, \Gamma_U^1, \Gamma_U^2$  are pairwise disjoint nonempty parts of  $\partial\Omega$ ,

$$\text{mes } \Gamma_D > 0, \quad \text{mes}(\Gamma_U^1 \cup \Gamma_U^2) > 0, \quad \text{mes}(\partial\Omega \setminus (\Gamma_D \cup \Gamma_N \cup \Gamma_U^1 \cup \Gamma_U^2)) = 0.$$

We consider the system (1.2) with unilateral boundary conditions

$$(5.2) \quad \begin{cases} u = v = 0 & \text{on } \Gamma_D, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \Gamma_N, \quad \frac{\partial v}{\partial \nu} = 0 & \text{on } \partial\Omega \setminus \Gamma_D, \\ u \geq 0, \quad \frac{\partial u}{\partial \nu} \geq 0, \quad u \cdot \frac{\partial u}{\partial \nu} = 0 & \text{on } \Gamma_U^1, \\ u \leq 0, \quad \frac{\partial u}{\partial \nu} \leq 0, \quad u \cdot \frac{\partial u}{\partial \nu} = 0 & \text{on } \Gamma_U^2, \end{cases}$$

or the system (5.1) with unilateral boundary conditions

$$(5.3) \quad \begin{cases} u = v = 0 & \text{on } \Gamma_D, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \setminus \Gamma_D, \quad \frac{\partial v}{\partial \nu} = 0 & \text{on } \Gamma_N, \\ v \geq 0, \quad \frac{\partial v}{\partial \nu} \geq 0, \quad v \cdot \frac{\partial v}{\partial \nu} = 0 & \text{on } \Gamma_U^1, \\ v \leq 0, \quad \frac{\partial v}{\partial \nu} \leq 0, \quad v \cdot \frac{\partial v}{\partial \nu} = 0 & \text{on } \Gamma_U^2. \end{cases}$$

It is natural to introduce a weak solution of (1.2), (5.2) or (5.1), (5.3) as a solution of (2.10) or (2.15), respectively, with the operators  $A$ ,  $N$  from (2.4), (2.5) and with the closed convex cone

$$K = \{\varphi \in \mathbb{H} : \varphi \geq 0 \text{ on } \Gamma_U^1, \varphi \leq 0 \text{ on } \Gamma_U^2 \text{ in the sense of traces}\}.$$

Hypotheses (3.5) or (3.8) mean in this example that

$$(5.4) \quad \text{there is a nontrivial } u = \sum_{j=j_0}^{\infty} \xi_j e_j \text{ with } u \geq 0 \text{ on } \Gamma_U^1 \text{ and } u \leq 0 \text{ on } \Gamma_U^2$$

or

$$(5.5) \quad \text{there is a nontrivial } u = \sum_{j=1}^{j_0} \xi_j e_j \text{ with } u \geq 0 \text{ on } \Gamma_U^1 \text{ and } u \leq 0 \text{ on } \Gamma_U^2,$$

respectively. Hypothesis (3.9) means that

$$(5.6) \quad \text{there is } u \in \mathbb{H} \text{ with } \langle u, e_{j_0} \rangle \neq 0 \text{ such that } u \geq 0 \text{ on } \Gamma_U^1, \text{ and } u \leq 0 \text{ on } \Gamma_U^2.$$

The assumption (3.4) is fulfilled if

$$(5.7) \quad \sum_{j \in I(d_1, d_2)} \alpha_j e_j \text{ changes sign in } \Gamma_U^1 \text{ or in } \Gamma_U^2$$

or is nonzero with constant sign on a set  $\Gamma$

$$\text{with } \text{mes}(\Gamma \cap \Gamma_U^i) > 0 \text{ for } i = 1, 2$$

for all nontrivial  $(\alpha_j)_{j \in I(d_1, d_2)}$ .

Hence, the following assertion follows by using Theorem 3.1, Corollary 3.1, and Remark 3.1.

**Theorem 5.1.** *Let  $d_2^0 > y_{j_0}$  where  $j_0$  is such that (5.4) holds. Then the problem (1.2), (5.2) has a bifurcation point with fixed  $d_2 = d_2^0$ . Moreover, if additionally for the unique  $d_1^{\max} > 0$  with  $(d_1^{\max}, d_2^0) \in C_E$  the hypothesis (5.7) is satisfied with  $(d_1, d_2) = (d_1^{\max}, d_2^0)$ , then  $(d_1, d_2^0) \in D_U$  for every bifurcation point  $d_1$  of (1.2), (5.2) with fixed  $d_2 = d_2^0$ .*

In general, it can happen that the largest bifurcating point in the situation of the last theorem is  $d_1^0$  with  $(d_1^0, d_2^0) \in C_E$  but the corresponding bifurcating solutions of (1.2), (5.2) are not solutions of (1.2), (1.7) because the bifurcating solutions of (1.2), (1.7) (if they exist) need not satisfy  $u \in K$  (cf. the text after Theorem 3.1).

Theorem 3.2 implies in view of Remark 3.3 or 3.4 the following results.

**Theorem 5.2.** *Let  $j_0$  be such that (5.5) holds. Then for any  $d_1^0 \in (0, x_{j_0})$  the problem (5.1), (5.3) has a bifurcation point with fixed  $d_1 = d_1^0$ .*

**Theorem 5.3.** *Let  $j_0$  be such that (5.6) holds. Then there is  $\varepsilon_{j_0} > 0$  such that for all  $d_1^0 \in [x_{j_0} - \varepsilon_{j_0}, x_{j_0})$  the problem (5.1), (5.3) has a bifurcation point with fixed  $d_1 = d_1^0$ .*

For  $j_0 = 1$ , we get the following modification which needs a proof.

**Theorem 5.4.** *If  $\text{mes } \Gamma_U^k > 0$  for both  $k = 1, 2$  then there is  $\varepsilon > 0$  such that for all  $d_1^0 \in [x_1 - \varepsilon, x_1)$  the problem (5.1), (5.3) has a bifurcation point with fixed  $d_1 = d_1^0$ , and  $(d_1^0, d_2) \in D_S$  for every such bifurcation point  $d_2$ .*

*Proof.* Since the eigenfunction  $e_1$  is simple and does not change sign, we have (5.7) for all  $(d_1, d_2)$  with  $I(d_1, d_2) = \{1\}$ . In particular, (3.4) holds if  $(d_1, d_2) = (d_1^0, d_2^{\max})$  with  $d_1^0 \in (x_1 - \varepsilon, x_1)$  ( $\varepsilon$  small) and  $d_2^{\max} > 0$  being the unique number with  $(d_1^0, d_2^{\max}) \in C_1$ . There is a function  $u \in \mathbb{H}$  with  $\langle u, e_1 \rangle \neq 0$ ,  $u \geq 0$  on  $\Gamma_U^1$  and  $u \leq 0$  on  $\Gamma_U^2$ . This  $u$  satisfies (5.6) with  $j_0 = 1$ , and so Remark 3.4 implies that the hypotheses of Theorem 3.1 and of Corollary 3.2 are satisfied. Our assertion follows.  $\square$

**Remark 5.1.** If  $\text{mes } \Gamma_U^1 = 0$  or if  $\text{mes } \Gamma_U^2 = 0$  then, since  $e_1$  does not change sign, the hypothesis (5.7) and thus (3.4) is not fulfilled for  $(d_1, d_2) \in C_1$ .

**Remark 5.2.** All previously known results about the existence of a bifurcation point of (5.1), (5.3) in  $D_S$  have had the hypothesis that

$$\text{there is } u = \sum_{j \in I(d_1, d_2)} \xi_j e_j \text{ with } u > 0 \text{ on } \Gamma_U^1 \text{ and } u < 0 \text{ on } \Gamma_U^2$$

for certain  $(d_1, d_2) \in \bigcup_{j=1}^{\infty} C_j$ . Theorem 5.4 applies in particular if this hypothesis is violated for all  $(d_1, d_2) \in \bigcup_{j=1}^{\infty} C_j$ .

**Example 5.2.** Let  $I_+$  and  $I_-$  be two finite sets of indices, at least one of them nonempty. Let  $\Gamma_D$ ,  $\Gamma_N$ , and  $\Gamma_i$  ( $i \in I_+ \cup I_-$ ) be pairwise disjoint parts of  $\partial\Omega$  such that

$$\text{mes } \Gamma_D > 0, \quad \text{mes } \Gamma_i > 0, \quad \text{mes} \left( \partial\Omega \setminus \left( \Gamma_D \cup \Gamma_N \cup \bigcup_{i \in I_+ \cup I_-} \Gamma_i \right) \right) = 0.$$

We consider the system (1.2) with unilateral boundary conditions

$$(5.8) \quad \begin{cases} u = v = 0 & \text{on } \Gamma_D, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \Gamma_N, \quad \frac{\partial v}{\partial \nu} = 0 & \text{on } \partial\Omega \setminus \Gamma_D, \\ \int_{\Gamma_i} u \, dx \geq 0, \quad \frac{\partial u}{\partial \nu} = \text{const} \geq 0, \quad \int_{\Gamma_i} u \, dx \frac{\partial u}{\partial \nu} = 0 & \text{on } \Gamma_i, \quad i \in I_+, \\ \int_{\Gamma_i} u \, dx \leq 0, \quad \frac{\partial u}{\partial \nu} = \text{const} \leq 0, \quad \int_{\Gamma_i} u \, dx \frac{\partial u}{\partial \nu} = 0 & \text{on } \Gamma_i, \quad i \in I_-, \end{cases}$$

or the system (5.1) with unilateral boundary conditions

$$(5.9) \quad \begin{cases} u = v = 0 & \text{on } \Gamma_D, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \setminus \Gamma_D, \quad \frac{\partial v}{\partial \nu} = 0 & \text{on } \Gamma_N, \\ \int_{\Gamma_i} v \, dx \geq 0, \quad \frac{\partial v}{\partial \nu} = \text{const} \geq 0, \quad \int_{\Gamma_i} v \, dx \frac{\partial v}{\partial \nu} = 0 & \text{on } \Gamma_i, \quad i \in I_+, \\ \int_{\Gamma_i} v \, dx \leq 0, \quad \frac{\partial v}{\partial \nu} = \text{const} \leq 0, \quad \int_{\Gamma_i} v \, dx \frac{\partial v}{\partial \nu} = 0 & \text{on } \Gamma_i, \quad i \in I_-. \end{cases}$$

It is natural to introduce a weak solution of (1.2), (5.8) or (5.1), (5.9) as a solution of (2.10) or (2.15), respectively, with the operators  $A$ ,  $N$  from (2.4), (2.5), and with the closed convex cone

$$K = \left\{ \varphi \in \mathbb{H} : \int_{\Gamma_i} \varphi \, dx \geq 0 \text{ for all } i \in I_+, \int_{\Gamma_i} \varphi \, dx \leq 0 \text{ for all } i \in I_- \right\}.$$

More precisely,  $u, v$  is a solution of (2.10) if and only if  $u, v \in \mathbb{H}$ ,  $\Delta u, \Delta v \in L_2(\Omega)$ , the equation (1.2) holds a.e. in  $\Omega$  and (5.8) is fulfilled; an analogous statement holds for (2.15). See [7, Observation 5.2] for details.

Hypotheses (3.5) or (3.8) mean in this example that

$$\text{there is a nontrivial } u = \sum_{j=j_0}^{\infty} \xi_j e_j \text{ such that } \sum_{j=j_0}^{\infty} \xi_j \int_{\Gamma_i} e_j \, dx \text{ is} \\ \text{nonnegative for all } i \in I_+ \text{ and nonpositive for all } i \in I_-,$$

or

there is a nontrivial  $u = \sum_{j=1}^{j_0} \xi_j e_j$  such that  $\sum_{j=1}^{j_0} \xi_j \int_{\Gamma_i} e_j dx$  is

nonnegative for all  $i \in I_+$  and nonpositive for all  $i \in I_-$ ,

respectively. Hypothesis (3.9) means

there is  $u \in \mathbb{H}$  with  $\langle u, e_{j_0} \rangle \neq 0$  such that  $\int_{\Gamma_i} u dx$  is

nonnegative for all  $i \in I_+$  and nonpositive for all  $i \in I_-$ .

The condition (3.4) is fulfilled if

for any nontrivial  $(\alpha_j)_{j \in I(d_1, d_2)}$  there are

$i_1, i_2$  lying both in  $I_+$  or both in  $I_-$  such that

$$(5.10) \quad \operatorname{sgn} \sum_{j \in I(d_1, d_2)} \alpha_j \int_{\Gamma_{i_1}} e_j dx = -\operatorname{sgn} \sum_{j \in I(d_1, d_2)} \alpha_j \int_{\Gamma_{i_2}} e_j dx$$

or there are  $i_1 \in I_+$ ,  $i_2 \in I_-$  such that

$$\operatorname{sgn} \sum_{j \in I(d_1, d_2)} \alpha_j \int_{\Gamma_{i_1}} e_j dx = \operatorname{sgn} \sum_{j \in I(d_1, d_2)} \alpha_j \int_{\Gamma_{i_2}} e_j dx.$$

If  $I_+ \neq \emptyset \neq I_-$  then (5.10) automatically holds for  $(d_1, d_2) \in C_1 \setminus \bigcup_{j=2}^{\infty} C_j$  due to the positivity and simplicity of the first eigenfunction  $e_1$  of (2.19).

Summarizing these facts to obtain assertions concerning all situations mentioned on the basis of the abstract results of Section 3 is left to the reader.

**Example 5.3.** Analogously to Example 5.1 or 5.2, we can consider also a regulation in the interior of the domain  $\Omega$  described by the closed convex cone

$$K = \{ \varphi \in \mathbb{H} : \varphi \geq 0 \text{ on } G_U^1, \varphi \leq 0 \text{ on } G_U^2 \}$$

or

$$K = \left\{ \varphi \in \mathbb{H} : \int_{G_i} \varphi dx \geq 0 \text{ for all } i \in I_+, \int_{G_i} \varphi dx \leq 0 \text{ for all } i \in I_- \right\},$$

where  $G_U^1, G_U^2$  or  $G_i$  ( $i \in I_+ \cup I_-$ ) are pairwise disjoint subsets of  $\Omega$ . It is straightforward to change the respective conditions (5.4)–(5.10) to these situations.

**Remark 5.3.** Let us suppose that the system (1.2) describes a coexistence of two populations with densities  $u, v$ . Let us consider a given nontrivial solution  $(u, v)$  of (1.2), (5.8) corresponding to some  $(d_1, d_2)$ . Hence,  $(u, v)$  is a nontrivial equilibrium of the original evolution system (1.6) with (5.8). Then the integral  $\int_{\Gamma_j} u \, dx$  determines the amount of the species  $u$  which can be removed (e.g. harvested) in the region  $\Gamma_j$  to keep the equilibrium. It is assumed that the amount harvested is the same at all places of  $\Gamma_j$ . Analogously for the integral  $\int_{G_j} u \, dx$  from Example 5.3. The boundary conditions from Example 5.1 can describe unilateral membranes, see e.g. [1].

**Remark 5.4.** Example 5.2 and the second case of Example 5.3 fit into the theory developed in [8], [19]. It follows that if the critical point  $(d_1, d_2)$  obtained from Theorem 3.1 or 3.2 used for the particular case  $n = 0$  is such that the nontrivial solution of the problem (2.12), (5.8) or (2.12), (5.9) is unique up to a positive multiple and satisfies certain activity conditions then a smooth branch of nontrivial solutions of the problem

$$(5.11) \quad \begin{aligned} \sigma_1(s)\Delta u + b_{11}u + b_{12}v + n_1(u, v) &= 0, \\ \sigma_2(s)\Delta v + b_{21}u + b_{22}v + n_2(u, v) &= 0 \end{aligned}$$

with the boundary conditions (5.8) or (5.9), respectively, bifurcates at a point  $s_B$  with  $(\sigma_1(s_B), \sigma_2(s_B)) = (d_1, d_2)$ , where  $\sigma$  is a smooth curve in  $\mathbb{R}_+^2$  containing  $(d_1, d_2)$ . The bifurcation direction for such situations is described in [8]. Cf. also [13] where an application of the result [19] is given to (5.11), (5.9).

#### References

- [1] *J. I. Baltaev, M. Kučera*: Global bifurcation for quasivariational inequalities of reaction-diffusion type. *J. Math. Anal. Appl.* *345* (2008), 917–928.
- [2] *P. Drábek, M. Kučera, M. Míková*: Bifurcation points of reaction-diffusion systems with unilateral conditions. *Czech. Math. J.* *35* (1985), 639–660.
- [3] *P. Drábek, A. Kufner, F. Nicolosi*: Quasilinear Elliptic Equations with Degenerations and Singularities. Walter de Gruyter, Berlin, 1997.
- [4] *L. Edelstein-Keshet*: *Mathematical Models in Biology*. McGraw-Hill, Boston, 1988.
- [5] *J. Eisner*: Critical and bifurcation points of reaction-diffusion systems with conditions given by inclusions. *Nonlinear Anal., Theory Methods Appl.* *46* (2001), 69–90.
- [6] *J. Eisner, M. Kučera*: Spatial patterning in reaction-diffusion systems with nonstandard boundary conditions. *Fields Institute Communications* *25* (2000), 239–256.
- [7] *J. Eisner, M. Kučera, L. Recke*: Smooth continuation of solutions and eigenvalues for variational inequalities based on the implicit function theorem. *J. Math. Anal. Appl.* *274* (2002), 159–180.
- [8] *J. Eisner, M. Kučera, L. Recke*: Direction and stability of bifurcating branches for variational inequalities. *J. Math. Anal. Appl.* *301* (2005), 276–294.

- [9] *J. Eisner, M. Kučera, M. Váth*: Global bifurcation for a reaction-diffusion system with inclusions. *J. Anal. Anwend.* *28* (2009), 373–409.
- [10] *S. Fučík, A. Kufner*: *Nonlinear Differential Equations*. Elsevier, Amsterdam-Oxford-New York, 1980.
- [11] *D. S. Jones, B. D. Sleeman*: *Differential Equations and Mathematical Biology*. Chapman & Hall/CRC, Boca Raton, 2003.
- [12] *M. Kučera*: Reaction-diffusion systems: Stabilizing effect of conditions described by quasivariational inequalities. *Czech. Math. J.* *47* (1997), 469–486.
- [13] *M. Kučera, L. Recke, J. Eisner*: Smooth bifurcation for variational inequalities and reaction-diffusion systems. *Progresses in Analysis* (J. G. W. Begehr, R. P. Gilbert, M. W. Wong, eds.). World Scientific, Singapore-New Jersey-London-Hong Kong, 2001, pp. 1125–1133.
- [14] *E. Miersemann*: Verzweigungsprobleme für Variationsungleichungen. *Math. Nachr.* *65* (1975), 187–209. (In German.)
- [15] *M. Mimura, Y. Nishiura, M. Yamaguti*: Some diffusive prey and predator systems and their bifurcation problems. *Ann. New York Acad. Sci.* *316* (1979), 490–510.
- [16] *J. D. Murray*: *Mathematical Biology*, 2nd ed. Springer, Berlin, 1993.
- [17] *Y. Nishiura*: Global structure of bifurcating solutions of some reaction-diffusion systems and their stability problem. *Proceedings of the 5th Int. Symp. Computing Methods in Applied Sciences and Engineering*, Versailles, France, 1981 (R. Glowinski, J. L. Lions, eds.). North-Holland, Amsterdam-New York-Oxford, 1982.
- [18] *P. Quittner*: Bifurcation points and eigenvalues of inequalities of reaction-diffusion type. *J. Reine Angew. Math.* *380* (1987), 1–13.
- [19] *L. Recke, J. Eisner, M. Kučera*: Smooth bifurcation for variational inequalities based on the implicit function theorem. *J. Math. Anal. Appl.* *275* (2002), 615–641.
- [20] *J. Smoller*: *Shock Waves and Reaction-Diffusion Equations*. Springer, New York-Heidelberg-Berlin, 1983.
- [21] *E. Zeidler*: *Nonlinear Functional Analysis and Its Applications*, vol. II/A. Springer, New York-Berlin-Heidelberg, 1990.
- [22] *E. Zeidler*: *Nonlinear Functional Analysis and Its Applications*, vol. III. Springer, New York-Berlin-Heidelberg, 1985.
- [23] *E. Zeidler*: *Nonlinear Functional Analysis and Its Applications*, vol. IV. Springer, New York-Berlin-Heidelberg, 1988.

*Authors' addresses:* *J. I. Baltaev*, Department of Mathematics, Urgench State University, Khorezm, Uzbekistan, e-mail: [jamol\\_b@yahoo.com](mailto:jamol_b@yahoo.com); *M. Kučera*, Institute of Mathematics of the Academy of Sciences of the Czech Republic, Žitná 25, 115 67 Prague 1, Czech Republic, e-mail: [kucera@math.cas.cz](mailto:kucera@math.cas.cz); *M. Váth*, Institute of Mathematics of the Academy of Sciences of the Czech Republic, Žitná 25, 115 67 Praha 1, Czech Republic, current address: Free University of Berlin, Dept. of Mathematics (WE1), Arnimallee 3, D-14195 Berlin, Germany, e-mail: [vaeth@mathematik.uni-wuerzburg.de](mailto:vaeth@mathematik.uni-wuerzburg.de).