

M. Sambasiva Rao

δ -ideals in pseudo-complemented distributive lattices

Archivum Mathematicum, Vol. 48 (2012), No. 2, 97--105

Persistent URL: <http://dml.cz/dmlcz/142823>

Terms of use:

© Masaryk University, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

δ -IDEALS IN PSEUDO-COMPLEMENTED DISTRIBUTIVE LATTICES

M. SAMBASIVA RAO

ABSTRACT. The concept of δ -ideals is introduced in a pseudo-complemented distributive lattice and some properties of these ideals are studied. Stone lattices are characterized in terms of δ -ideals. A set of equivalent conditions is obtained to characterize a Boolean algebra in terms of δ -ideals. Finally, some properties of δ -ideals are studied with respect to homomorphisms and filter congruences.

INTRODUCTION

The theory of pseudo-complements was introduced and extensively studied in semi-lattices and particularly in distributive lattices by Orrin Frink [4] and G. Birkhoff [2]. Later the problem of characterizing Stone lattices has been studied by several authors like R. Balbes [1], O. Frink [4], G. Grätzer [5] etc.

In this paper, the concept of δ -ideals is introduced in a distributive lattice in terms of pseudo-complementation and filters. Some properties of these δ -ideals are studied and then proved that the set of all δ -ideals can be made into a complete distributive lattice. We derive a set of equivalent conditions for the class of all δ -ideals to become a sublattice to the lattice of all ideals, which leads to a characterization of Stone lattices. A set of equivalent conditions are established for every prime ideal to become a δ -ideal which leads to a characterization of a Boolean algebra. Finally, the set of δ -ideals of a pseudo-complemented distributive lattice is characterized in terms of filter congruences.

1. PRELIMINARIES

In this section, we recall certain definitions and important results taken from [6] for the ready reference to the reader.

Definition 1.1 ([6]). For any element a of a distributive lattice L , the pseudo-complement a^* of a is an element satisfying the following property for all $x \in L$:

$$a \wedge x = 0 \Leftrightarrow a^* \wedge x = x \Leftrightarrow x \leq a^* .$$

2010 *Mathematics Subject Classification*: primary 06D99; secondary 06D15.

Key words and phrases: pseudo-complemented distributive lattice, dense element, closed element, δ -ideal, Stone lattice, congruence.

Received July 10, 2011. Editor J. Rosický.

DOI: 10.5817/AM2012-2-97

A distributive lattice L in which every element has a pseudo-complement is called a pseudo-complemented distributive lattice. Throughout this paper L stands for a pseudo-complemented distributive lattice $(L, \vee, \wedge, *, 0, 1)$.

Theorem 1.2 ([6]). *For any two elements a, b of a pseudo-complemented distributive lattice, we have the following:*

- (1) $0^{**} = 0$,
- (2) $a \wedge a^* = 0$,
- (3) $a \leq b$ implies $b^* \leq a^*$,
- (4) $a \leq a^{**}$,
- (5) $a^{***} = a^*$,
- (6) $(a \vee b)^* = a^* \wedge b^*$,
- (7) $(a \wedge b)^{**} = a^{**} \wedge b^{**}$.

An element x of a pseudo-complemented lattice L is called dense [6] if $x^* = 0$ and the set $D(L)$ of all dense element of L forms a filter of L .

Definition 1.3 ([1]). A pseudo-complemented distributive lattice L is called a Stone lattice if, for all $x \in L$, it satisfies the property: $x^* \vee x^{**} = 1$.

Theorem 1.4 ([6]). *Let I be an ideal and F a filter of a distributive lattice L such that $I \cap F = \emptyset$. Then there exists a prime ideal P such that $I \subseteq P$ and $P \cap F = \emptyset$.*

A prime ideal P of a distributive lattice L is called a minimal prime ideal [8] if there exists no prime ideal Q such that $Q \subset P$. A prime ideal P is minimal if and only if to each $x \in P$ there exists $y \notin P$ such that $x \wedge y = 0$.

2. δ -IDEALS

In this section, the concept of δ -ideals is introduced in a pseudo-complemented distributive lattice. Stone lattice and Boolean algebras are characterized in terms of δ -ideals. Finally, δ -ideals are characterized in terms of congruences.

Definition 2.1. Let L be a pseudo-complemented distributive lattice. Then for any filter F of L , define the set $\delta(F)$ as follows:

$$\delta(F) = \{x \in L \mid x^* \in F\}.$$

In the following, some basic properties of $\delta(F)$ can be observed.

Lemma 2.2. *Let L be a pseudo-complemented distributive lattice. Then for any filter F of L , $\delta(F)$ is an ideal of L .*

Proof. Since $0^* \in F$, we get that $0 \in \delta(F)$. Let $x, y \in \delta(F)$. Then $x^*, y^* \in F$. Hence $(x \vee y)^* = x^* \wedge y^* \in F$. Again, let $x \in \delta(F)$ and $r \in L$. Then $x^* \in F$. Hence $(x \wedge r)^* = (x \wedge r)^{***} = (x^{**} \wedge r^{**})^* = (x^* \vee r^*)^{**} \in F$ (because $x^* \vee r^* \in F$). Hence we get that $x \wedge r \in \delta(F)$. Therefore $\delta(F)$ is an ideal in L . \square

Lemma 2.3. *Let L be a pseudo-complemented distributive lattice. For any two filters F, G of L , we have the following:*

- (1) $F \cap \delta(F) = \emptyset$,
- (2) $x \in \delta(F) \Rightarrow x^{**} \in \delta(F)$,
- (3) $F = L$ if and only if $\delta(F) = L$,
- (4) $F \subseteq G \Rightarrow \delta(F) \subseteq \delta(G)$,
- (5) $\delta(F \cap G) = \delta(F) \cap \delta(G)$.

Proof. (1) Suppose $x \in F \cap \delta(F)$. Then $x \in F$ and $x^* \in F$. Since F is a filter, we get $0 = x^* \wedge x \in F$, which is a contradiction. Therefore $F \cap \delta(F) = \emptyset$.

(2) Since $x^{***} = x^*$, it is clear.

(3) Assume that $F = L$. Then we have $0^{**} = 0 \in F$. Hence $0^* \in \delta(F)$. Therefore $\delta(F) = L$. Converse is an easy reverse of the above.


(4) Suppose $F \subseteq G$. Let $x \in \delta(F)$. Then $x^* \in F \subseteq G$. Therefore $x \in \delta(G)$.

(5) Clearly $\delta(F \cap G) \subseteq \delta(F) \cap \delta(G)$. Conversely, let $x \in \delta(F) \cap \delta(G)$. Then $x^* \in F \cap G$. Hence $x \in \delta(F \cap G)$. Therefore $\delta(F) \cap \delta(G) \subseteq \delta(F \cap G)$. □

The concept of δ -ideals is now introduced in the following.

Definition 2.4. Let L be a pseudo-complemented distributive lattice. An ideal I of L is called a δ -ideal if $I = \delta(F)$ for some filter F of L .

Example 2.5. Consider the distributive lattice $L = \{0, a, b, c, 1\}$ whose Hasse diagram is given in the following figure:



Consider $I = \{0, a\}$ and $F = \{b, c, 1\}$.

Clearly I is an ideal and F a filter of L .

Now $\delta(F) = \{x \mid x^* \in F\} = \{0, a\}$.

Therefore I is a δ -ideal of L .

But $J = \{0, a, b, c\}$ is not a δ -ideal of L .

Suppose $J = \delta(F)$ for some filter F

Then $0 = c^* \in F$. Hence $F = L$, which yields that $J = \delta(F) = L$.

The following lemmas produce some more examples for δ -ideals.

Lemma 2.6. For each $x \in L$, (x^*) is a δ -ideal of L .

Proof. It is enough to show that $(x^*) = \delta([x])$. Let $a \in (x^*)$. Then $a \wedge x = 0$ and hence $a^* \wedge x = x \in [x]$. Thus $a^* \in [x]$. Therefore $a \in \delta([x])$. Conversely, suppose that $a \in \delta([x])$. Then $a^* \wedge x = x$ and hence $a \wedge x = a \wedge a^* \wedge x = 0$. Thus $a \wedge x^* = a$, which yields that $a \in (x^*)$. Therefore (x^*) is a δ -ideal of L . □

Lemma 2.7. Every prime ideal without dense element is a δ -ideal.

Proof. Let P be a prime ideal with out dense element. Let $x \in P$. Then clearly $x \wedge x^* = 0$ and $x \vee x^*$ is dense. Hence $x \vee x^* \notin P$. Thus we get $x^* \notin P$, which yields that $x^* \in L - P$. Thus $x \in \delta(L - P)$. Therefore $P \subseteq \delta(L - P)$. Conversely, let $x \in \delta(L - P)$. Then $x^* \in L - P$, which implies that $x^* \notin P$. Hence $x \in P$. Thus $P = \delta(L - P)$. Therefore P is a δ -ideal of L . □

Corollary 2.8. *Every minimal prime ideal is a δ -ideal.*

Proof. Let P be a minimal prime ideal of L . Suppose $x \in P \cap D(L)$. Then there exists $y \notin P$ such that $x \wedge y = 0$. Hence $y \leq x^* = 0 \in P$, which is a contradiction. Thus $P \cap D(L) = \emptyset$. Therefore, by above lemma, P is a δ -ideal. \square

In the following, a simple property of δ -ideals is observed.

Lemma 2.9. *A proper δ -ideal contains no dense element.*

Proof. Let $\delta(F)$ be a proper δ -ideal of L . Suppose $x \in \delta(F) \cap D(L)$. Then we get $0 = x^* \in F$, which is a contradiction. Therefore $\delta(F) \cap D(L) = \emptyset$. \square

Let us denote the set of all δ -ideals of L by $\mathcal{I}^\delta(L)$. Then by Example 2.5, it can be observed that $\mathcal{I}^\delta(L)$ is not a sublattice of $\mathcal{I}(L)$ of all ideals of L . Consider $F = \{b, c, 1\}$ and $G = \{a, c, 1\}$. Clearly F and G are filters of L . Now $\delta(F) = \{0, a\}$ and $\delta(G) = \{0, b\}$. But $\delta(F) \vee \delta(G) = \{0, a, b, c\}$ is not a δ -ideal of L , because $c \in \delta(F) \vee \delta(G)$ is a dense element. However, in the following theorem, we prove that $\mathcal{I}^\delta(L)$ forms a complete distributive lattice.

Theorem 2.10. *Let L be a pseudo-complemented distributive lattice. Then the set $\mathcal{I}^\delta(L)$ forms a complete distributive lattice on its own.*

Proof. For any two filters F, G of L , define two binary operations \cap and \sqcup as follows:

$$\delta(F) \cap \delta(G) = \delta(F \cap G) \quad \text{and} \quad \delta(F) \sqcup \delta(G) = \delta(F \vee G).$$

It is clear that $\delta(F \cap G)$ is the infimum of $\delta(F)$ and $\delta(G)$ in $\mathcal{I}^\delta(L)$. Also $\delta(F) \sqcup \delta(G)$ is a δ -ideal of L . Clearly $\delta(F), \delta(G) \subseteq \delta(F \vee G) = \delta(F) \sqcup \delta(G)$. Let $\delta(H)$ be a δ -ideal of L such that $\delta(F) \subseteq \delta(H)$ and $\delta(G) \subseteq \delta(H)$, where H is a filter of L . Now we claim that $\delta(F \vee G) \subseteq \delta(H)$. Let $x \in \delta(F \vee G)$. Then $x^* \in F \vee G$. Hence $x^* = f \wedge g$ for some $f \in F$ and $g \in G$. Since $f \in F$ and $g \in G$, we can get $f^* \in \delta(F) \subseteq \delta(H)$ and $g^* \in \delta(G) \subseteq \delta(H)$. Now

$$\begin{aligned} f^* \in \delta(H), g^* \in \delta(H) &\Rightarrow f^* \vee g^* \in \delta(H) \\ &\Rightarrow (f^* \vee g^*)^{**} \in \delta(H) \\ &\Rightarrow (f^{**} \wedge g^{**})^* \in \delta(H) \\ &\Rightarrow x^{**} \in \delta(H) \\ &\Rightarrow x \in \delta(H) \end{aligned}$$

Hence $\delta(F) \sqcup \delta(G) = \delta(F \vee G)$ is the supremum of both $\delta(F)$ and $\delta(G)$ in $\mathcal{I}^\delta(L)$. Therefore $\langle \mathcal{I}^\delta(L), \cap, \sqcup \rangle$ is a lattice. Distributivity of δ -ideals can be easily followed by using the above operations of $\mathcal{I}^\delta(L)$.

It is clear that $\mathcal{I}^\delta(L)$ is a partially ordered set with respect to set-inclusion. Then by the extension of the property of Lemma 2.3(5), we can obtain that $\mathcal{I}^\delta(L)$ is a complete lattice. Therefore $\mathcal{I}^\delta(L)$ is a complete distributive lattice. \square

From Lemma 2.6, we have already observed that each (x^*) (for $x \in L$) is a δ -ideal of L . Now let us denote that $\mathcal{A}^*(L) = \{(x^*) \mid x \in L\}$. Then, in the following theorem, it is proved that $\mathcal{A}^*(L)$ is a Boolean algebra.

Theorem 2.11. *For any pseudo-complemented distributive lattice L , $\mathcal{A}^*(L)$ is a sublattice of the lattice $\mathcal{I}^\delta(L)$ of all δ -ideals of L and hence is a Boolean algebra. Moreover, the mapping $x \mapsto (x^*]$ is a dual homomorphism from L onto $\mathcal{A}^*(L)$.*

Proof. Let $(x^*], (y^*] \in \mathcal{A}^*(L)$ for some $x, y \in L$. Then clearly $(x^*] \cap (y^*] \in \mathcal{A}^*(L)$. Again, $(x^*] \sqcup (y^*] = \delta([x]) \sqcup \delta([y]) = \delta([x] \vee [y]) = \delta([x \wedge y]) = ((x \wedge y)^*) \in \mathcal{A}^*(L)$. Hence $\mathcal{A}^*(L)$ is a sublattice of $\mathcal{I}^\delta(L)$ and hence a distributive lattice. Clearly (0^{**}) and $(0^*]$ are the least and greatest elements of $\mathcal{A}^*(L)$. Now for any $x \in L$, $(x^*] \cap (x^{**}) = (0)$ and $(x^*] \sqcup (x^{**}) = \delta([x]) \sqcup \delta([x^*]) = \delta([x] \vee [x^*]) = \delta([x \wedge x^*]) = \delta([0]) = \delta(L) = L$. Hence (x^{**}) is the complement of $(x^*]$ in $\mathcal{A}^*(L)$. Therefore $\langle \mathcal{A}^*(L), \sqcup, \cap \rangle$ is a bounded distributive lattice in which every element is complemented. The remaining part can be easily observed. \square

It was already observed that $\mathcal{I}^\delta(L)$ is not a sublattice of the ideal lattice $\mathcal{I}(L)$. However, we establish some equivalent conditions for $\mathcal{I}^\delta(L)$ to become a sublattice of $\mathcal{I}(L)$, which leads to a characterization of Stone lattices. For this, we need the following lemma.

Lemma 2.12. *Every proper δ -ideal is contained in a minimal prime ideal.*

Proof. Let I be a proper δ -ideal of L . Then $I = \delta(F)$ for some filter F of L . Clearly $\delta(F) \cap D(L) = \emptyset$. Then there exists a prime ideal P of L such that $\delta(F) \subseteq P$ and $P \cap D(L) = \emptyset$. Let $x \in P$. We have always $x \wedge x^* = 0$. Suppose $x^* \in P$. Then $x \vee x^* \in P \cap D(L)$, which is a contradiction. Thus P is a minimal prime ideal of L . \square

Since the minimal prime ideals are precisely the complements of maximal filter of L , the following corollary is a direct consequence.

Corollary 2.13. *The minimal prime ideals of a pseudo-complemented distributive lattice L are maximal elements of the complete lattice $\mathcal{I}^\delta(L)$.*

We now characterize Stone lattices in terms of δ -ideals.

Theorem 2.14. *Let L be a pseudo-complemented distributive lattice. Then the following are equivalent:*

- (1) L is a Stone lattice,
- (2) For any $x, y \in L$, $(x \wedge y)^* = x^* \vee y^*$,
- (3) For any two filters F, G of L , $\delta(F) \vee \delta(G) = \delta(F \vee G)$,
- (4) $\mathcal{I}^\delta(L)$ is a sublattice of $\mathcal{I}(L)$.

Proof. (1) \Rightarrow (2): It is obtained in Lemma 3 of [6] pp.113.

(2) \Rightarrow (3): Assume the condition (2). Let F, G be two filters of L . We have always $\delta(F) \vee \delta(G) \subseteq \delta(F \vee G)$. Conversely, let $x \in \delta(F \vee G)$. Then

$$\begin{aligned} x^* \in F \vee G &\Rightarrow x^* = f \wedge g \quad \text{for some } f \in F, g \in G \\ &\Rightarrow x^{**} = (f \wedge g)^* \\ &\Rightarrow x^{**} = f^* \vee g^* \\ &\Rightarrow x^{**} = f^* \vee g^* \in \delta(F) \vee \delta(G) \quad \text{since } f^{**} \in F, g^{**} \in G \\ &\Rightarrow x \in \delta(F) \vee \delta(G). \end{aligned}$$

Hence $\delta(F \vee G) \subseteq \delta(F) \vee \delta(G)$. Therefore $\delta(F) \vee \delta(G) = \delta(F \vee G)$.

(3) \Rightarrow (4): It is obvious.

(4) \Rightarrow (1): Assume that $\mathcal{I}^\delta(L)$ is a sublattice of $\mathcal{I}(L)$. Let $x \in L$. By Lemma 2.6, $(x^*]$ and $(x^{**}]$ are both δ -ideals of L . Suppose $x^* \vee x^{**} \neq 1$. Then by condition (4), $(x^*] \vee (x^{**}]$ is a proper δ -ideal of L . Hence there exists a minimal prime ideal P such that $(x^*] \vee (x^{**}] \subseteq P$. Since P is minimal, we get that $x^{**} \notin P$, which is a contradiction. Therefore L is a Stone lattice. \square

In the following theorem, a set of equivalent conditions are obtained for every prime ideal of L to become a δ -ideal which in turn leads to establish some equivalent conditions for a pseudo-complemented distributive lattice to become a Boolean algebra. Let us recall that an element x is called closed [6] if $x^{**} = x$.

Theorem 2.15. *Let L be a pseudo-complemented distributive lattice. Then the following conditions are equivalent:*

- (1) L is a Boolean algebra,
- (2) Every element of L is closed,
- (3) Every principal ideal is a δ -ideal,
- (4) For any ideal I , $x \in I$ implies $x^{**} \in I$,
- (5) For any proper ideal I , $I \cap D(L) = \emptyset$,
- (6) For any prime ideal P , $P \cap D(L) = \emptyset$,
- (7) Every prime ideal is a minimal prime ideal,
- (8) Every prime ideal is a δ -ideal,
- (9) For any $x, y \in L$, $x^* = y^*$ implies $x = y$,
- (10) $D(L)$ is a singleton set.

Proof. (1) \Rightarrow (2): Assume that L is a Boolean algebra. Then clearly L has a unique dense element, precisely the greatest element. Let $x \in L$. Then $x^* \wedge x = 0 = x^* \wedge x^{**}$. Also $x^* \vee x, x^* \vee x^{**} \in D(L)$. Hence $x^* \vee x = x^* \vee x^{**}$. By the cancelation property of L , we get $x = x^{**}$. Therefore every element of L is closed.

(2) \Rightarrow (3): Let I be a principal ideal of L . Then $I = (x]$ for some $x \in L$. Then by condition (2), $x = x^{**}$. Now, $(x) = (x^{**}) = \delta([x^*])$. Therefore $(x]$ is a δ -ideal.

(3) \Rightarrow (4): Let I be a proper ideal of L . Let $x \in I$. Then $(x) = \delta(F)$ for some filter F of L . Hence we get $x^{***} = x^* \in F$. Therefore $x^{**} \in \delta(F) = (x) \subseteq I$.

(4) \Rightarrow (5): Let I be a proper ideal of L . Suppose $x \in I \cap D(L)$. Then $x^{**} \in I$ and $x^* = 0$. Therefore $1 = 0^* = x^{**} \in I$, which is a contradiction.

(5) \Rightarrow (6): It is clear.

(6) \Rightarrow (7): Let P be a prime ideal of L such that $P \cap D(L) = \emptyset$. Let $x \in P$. Then clearly $x \wedge x^* = 0$ and $x \vee x^* \in D(L)$. Hence $x \vee x^* \notin P$. Thus $x^* \notin P$. Therefore P is a minimal prime ideal of L .

(7) \Rightarrow (8): Let P be a minimal prime ideal of L . Then clearly $L - P$ is a filter of L . Let $x \in P$. Since P is minimal, there exists $y \notin P$ such that $x \wedge y = 0$. Hence $x^* \wedge y = y$, which implies that $x^* \notin P$. Thus $x^* \in L - P$, which yields $x \in \delta(L - P)$. Conversely, let $x \in \delta(L - P)$. Then we get $x^* \notin P$. Hence we have $x \in P$. Thus

$P = \delta(L - P)$ and therefore P is a δ -ideal of L .

(8) \Rightarrow (9): Assume that every prime ideal of L is a δ -ideal. Let $x, y \in L$ be such that $x^* = y^*$. Suppose $x \neq y$. Then there exists a prime ideal P of L such that $x \in P$ and $y \notin P$. By hypothesis, P is a δ -ideal of L . Hence $P = \delta(F)$ for some filter F of L . Since $x \in P = \delta(F)$, we get $y^* = x^* \in F$. Hence $y \in \delta(F) = P$, which is a contradiction. Therefore $x = y$.

(9) \Rightarrow (10): Suppose a, b be two elements of $D(L)$. Then $a^* = 0 = b^*$. Hence $a = b$. Therefore $D(L)$ is a singleton set.

(10) \Rightarrow (1): Assume that $D(L) = \{d\}$ is a singleton set. Let $x \in L$. We have always $x \vee x^* \in D(L)$. Therefore $x \wedge x^* = 0$ and $x \vee x^* = d$. This true for all $x \in L$. Also $0 \leq x \leq x \vee x^* = d$. Therefore L is a bounded distributive lattice in which every element is complemented. \square

We now prove that the homomorphic image of a δ -ideal is again a δ -ideal. By a homomorphism [6] on a bounded lattice, we mean a homomorphism which preserves 0 and 1. We now start our observation with the following fact.

Unlike in rings, if an onto homomorphism of a distributive lattice L into another lattice L' such that $\ker f = \{x \in L \mid f(x) = 0\} = \{0\}$, then f need not be an isomorphism. For this, we consider two chains $L = \{0, a, 1\}$ and $L' = \{0', 1'\}$. Now, define a mapping $f: L \rightarrow L'$ by $f(0) = 0'$ and $f(a) = f(1) = 1'$. Then clearly f is a homomorphism from L into L' and also f is onto. Also $\text{Ker } f = \{0\}$. But f is not one-one. Hence f is not an isomorphism.

Lemma 2.16. *Let L and L' be two pseudo-complemented distributive lattices with pseudo-complementation $*$ and $f: L \rightarrow L'$ an onto homomorphism. If $\text{Ker } f = \{0\}$, then $f(x^*) = \{f(x)\}^*$ for all $x \in L$.*

Proof. We have always $f(x) \wedge f(x^*) = f(x \wedge x^*) = f(0) = 0$. Suppose $f(x) \wedge f(t) = 0$ for some $t \in L$. Then $f(x \wedge t) = 0$ and hence $x \wedge t \in \ker f = \{0\}$. Thus $x \wedge t = 0$. Hence $x^* \wedge t = t$, which yields $f(x^*) \wedge f(t) = f(x^* \wedge t) = f(t)$. Therefore $f(x^*)$ is the pseudo-complement of $f(x)$ in L' . \square

In the following, we prove that the image of a δ -ideal of L under the above homomorphism is again a δ -ideal.

Theorem 2.17. *Let L, L' be two pseudo-complemented distributive lattices with pseudo-complementation $*$ and $f: L \rightarrow L'$ an onto homomorphism such that $\text{Ker } f = \{0\}$. If I is a δ -ideal of L , then $f(I)$ is a δ -ideal of L' .*

Proof. Let I be a δ -ideal of L . Then $I = \delta(G)$ for some filter G of L . Since the homomorphism f preserves 1, we can get that $f(G)$ is a filter in L' . Now, it is enough to show that $f\{\delta(G)\} = \delta\{f(G)\}$. Let $a \in f\{\delta(G)\}$. Then $a = f(x)$ for some $x \in \delta(G)$. Hence $x^* \in G$. Now $f(x) \wedge f(x^*) = f(x \wedge x^*) = f(0) = 0$. Hence $\{f(x)\}^* \wedge f(x^*) = f(x^*) \in f(G)$. Thus $\{f(x)\}^* \in f(G)$. Therefore $a = f(x) \in \delta\{f(G)\}$. Therefore $f\{\delta(G)\} \subseteq \delta\{f(G)\}$. Conversely, let $y \in \delta\{f(G)\}$. Since f is on-to, there exists $x \in L$ such that $y = f(x)$. Then $\{f(x)\}^* \in f(G)$. Hence

$\{f(x)\}^* = f(a)$ for some $a \in G$. Now

$$\begin{aligned} f(x) \wedge \{f(x)\}^* = 0 &\Rightarrow f(x) \wedge f(a) = 0 \\ &\Rightarrow f(x \wedge a) = 0 \\ &\Rightarrow x \wedge a \in \text{Ker } f = \{0\} \\ &\Rightarrow x^* \wedge a = a \in G \\ &\Rightarrow x^* \in G \\ &\Rightarrow x \in \delta(G) \\ &\Rightarrow y = f(x) \in f\{\delta(G)\}. \end{aligned}$$

Thus $\delta\{f(G)\} \subseteq f\{\delta(G)\}$. Therefore $\delta\{f(G)\} = f\{\delta(G)\}$. \square

We now characterize δ -ideals in terms of congruences. For this, we consider a well known filter congruence introduced by T. P. Speed [7].

Theorem 2.18 ([7]). *For any filter F of L , define a relation $\theta(F)$ as follows:*

$$(a, b) \in \theta(F) \Leftrightarrow a \wedge f = b \wedge f \text{ for some } f \in F.$$

Then $\theta(F)$ is a congruence relation on L .

Lemma 2.19. *Let L be a pseudo-complemented distributive lattice. Then for any ideal I of L , $F_I = \{x \in L \mid x^* \wedge a^* = 0 \text{ for some } a \in I\}$ is a filter of L .*

Proof. Clearly $0^* \in F_I$. Let $x, y \in F_I$. Then $x^* \wedge a^* = 0$ and $y^* \wedge b^* = 0$ for some $a, b \in I$. Hence $x^{**} \wedge a^* = a^*$ and $y^{**} \wedge b^* = b^*$. Now $(x \wedge y)^{**} \wedge (a \vee b)^* = x^{**} \wedge y^{**} \wedge a^* \wedge b^* = a^* \wedge b^*$. Thus $(x \wedge y)^* \wedge (a \vee b)^* = 0$. Therefore $x \wedge y \in F_I$. Again, let $x \in F_I$ and $s \in L$. Then $x^* \wedge a^* = 0$ for some $a \in I$. Now $(x \vee s)^* \wedge a^* \leq x^* \wedge a^* = 0$. Thus $x \vee s \in F_I$. Therefore F_I is a filter of L . \square

Theorem 2.20. *For any ideal I of a pseudo-complemented distributive lattice L , the following conditions are equivalent:*

- (1) I is a δ -ideal,
- (2) $I = \text{Ker } \theta(F_I)$,
- (3) $I = \text{Ker } \theta(F)$ for some filter F of L .

Proof. (1) \Rightarrow (2): Assume that I is a δ -ideal of L . Then $I = \delta(F)$ for some filter F of L . Let $x \in I$. Since $x^{**} \wedge x^* = 0$, we can get $x^* \in F_I$. Since $x \wedge x^* = 0$ and $x^* \in F_I$, we thus get $x \in \text{Ker } \theta(F_I)$. Therefore $I \subseteq \text{Ker } \theta(F_I)$. Conversely, let $x \in \text{Ker } \theta(F_I)$. Then $(x, 0) \in \theta(F_I)$. Thus $x \wedge f = 0$ for some $f \in F_I$. Since $f \in F_I$, we get that $f^* \wedge a^* = 0$ for some $a \in I$. Hence $x \leq f^* \leq a^{**} \in \delta(F) = I$. Therefore $I = \text{Ker } \theta(F_I)$.

(2) \Rightarrow (3): It is clear.

(3) \Rightarrow (1): Assume that $I = \text{Ker } \theta(F)$ for some filter F of L . Let $x \in I = \text{Ker } \theta(F)$. Then $x \wedge f = 0$ for some $f \in F$. Hence $x^* \wedge f = f \in F$. Thus $x^* \in F$, which yields that $x \in \delta(F)$. Therefore $I \subseteq \delta(F)$. Conversely, let $x \in \delta(F)$. Then $x^* \in F$. Since $x \wedge x^* = 0$ and $x^* \in F$, we get $(x, 0) \in \theta(F)$. Thus $x \in \text{Ker } \theta(F) = I$. Therefore I is a δ -ideal of L . \square

Acknowledgement. The author is grateful to the referee for his valuable comments and suggestions for the improvement of the paper.

REFERENCES

- [1] Balbes, R., Horn, A., *Stone lattices*, Duke Math. J. **37** (1970), 537–545.
- [2] Birkhoff, G., *Lattice Theory*, Colloquium Publications, vol. 25, Amer. Math. Soc., New York, 1948.
- [3] Cornish, W. H., *Congruences on distributive pseudocomplemented lattices*, Bull. Austral. Math. Soc. **8** (1973), 161–179.
- [4] Frink, O., *Pseudo-complements in semi-lattices*, Duke Math. J. **29** (1962), 505–514.
- [5] Grätzer, G., *A generalization on Stone's representation theorem for Boolean algebras*, Duke Math. J. **30** (1963), 469–474.
- [6] Grätzer, G., *General lattice theory*, Academic Press, New York, San Francisco, 1978.
- [7] Speed, T. P., *Two congruences on distributive lattices*, Bull. Soc. Roy. Sci. Liège **38** (3–4) (1969), 86–95.
- [8] Speed, T. P., *Spaces of ideals of distributive lattices II. Minimal prime ideals*, J. Austral. Math. Soc. **18** (1974), 54–72.

DEPARTMENT OF MATHEMATICS,
MVGR COLLEGE OF ENGINEERING, CHINTALVALASA,
VIZIANAGARAM, ANDHRA PRADESH, INDIA-535005
E-mail: mssraomaths35@rediffmail.com