

Huaning Liu; Jing Gao

Generalized Knopp identities for homogeneous Hardy sums and Cochrane-Hardy sums

Czechoslovak Mathematical Journal, Vol. 62 (2012), No. 4, 1147–1159

Persistent URL: <http://dml.cz/dmlcz/143050>

Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

GENERALIZED KNOPP IDENTITIES FOR HOMOGENEOUS
HARDY SUMS AND COCHRANE-HARDY SUMS

HUANING LIU, JING GAO, Xi'an

(Received September 27, 2011)

Abstract. Let q, h, a, b be integers with $q > 0$. The classical and the homogeneous Dedekind sums are defined by

$$s(h, q) = \sum_{j=1}^q \left(\left(\frac{j}{q} \right) \right) \left(\left(\frac{hj}{q} \right) \right), \quad s(a, b, q) = \sum_{j=1}^q \left(\left(\frac{aj}{q} \right) \right) \left(\left(\frac{bj}{q} \right) \right),$$

respectively, where

$$((x)) = \begin{cases} x - [x] - \frac{1}{2}, & \text{if } x \text{ is not an integer;} \\ 0, & \text{if } x \text{ is an integer.} \end{cases}$$

The Knopp identities for the classical and the homogeneous Dedekind sum were the following:

$$\sum_{d|n} \sum_{r=1}^d s\left(\frac{n}{d}a + rq, dq\right) = \sigma(n)s(a, q),$$

$$\sum_{d|n} \sum_{r_1=1}^d \sum_{r_2=1}^d s\left(\frac{n}{d}a + r_1q, \frac{n}{d}b + r_2q, dq\right) = n\sigma(n)s(a, b, q),$$

where $\sigma(n) = \sum_{d|n} d$.

In this paper generalized homogeneous Hardy sums and Cochrane-Hardy sums are defined, and their arithmetic properties are studied. Generalized Knopp identities for homogeneous Hardy sums and Cochrane-Hardy sums are given.

Keywords: Dedekind sum, Cochrane sum, Knopp identity

MSC 2010: 11F20

Supported by the National Natural Science Foundation of China under Grant No. 10901128, the Specialized Research Fund for the Doctoral Program of Higher Education of China under Grant No. 20090201120061, and the Fundamental Research Funds for the Central University.

1. INTRODUCTION

For a positive integer q and an arbitrary integer h , the classical Dedekind sum is defined by

$$s(h, q) = \sum_{j=1}^q \left(\left(\frac{j}{q} \right) \right) \left(\left(\frac{hj}{q} \right) \right),$$

where

$$\left(\left(x \right) \right) = \begin{cases} x - [x] - \frac{1}{2}, & \text{if } x \text{ is not an integer;} \\ 0, & \text{if } x \text{ is an integer.} \end{cases}$$

The sum $s(h, q)$ plays an important role in the transformation theory of the Dedekind η function (see [9] and Chapter 3 of [1] for details).

In [6], Knopp derived the following arithmetical identity

$$\sum_{d|n} \sum_{r=1}^d s\left(\frac{n}{d}a + rq, dq\right) = \sigma(n)s(a, q),$$

where $\sigma(n) = \sum_{d|n} d$. Many authors have given elementary proofs for the Knopp identity, for example, Goldberg [4], Parson [7] and Zheng [15].

According to [5], the homogeneous Dedekind sum is defined by

$$s(a, b, q) = \sum_{j=1}^q \left(\left(\frac{aj}{q} \right) \right) \left(\left(\frac{bj}{q} \right) \right).$$

Zheng [16] extended the Knopp identity to homogeneous Dedekind sums as follows:

$$\sum_{d|n} \sum_{r_1=1}^d \sum_{r_2=1}^d s\left(\frac{n}{d}a + r_1q, \frac{n}{d}b + r_2q, dq\right) = n\sigma(n)s(a, b, q).$$

Moreover, let $B_r(x)$ be the Bernoulli polynomials given by

$$\frac{ze^{xz}}{e^z - 1} = \sum_{r=0}^{\infty} B_r(x) \frac{z^r}{r!} \quad (|z| < 2\pi)$$

and

$$P_r(x) = \begin{cases} B_r(x - [x]), & \text{if } x \text{ is not an integer;} \\ 0, & \text{if } x \text{ is an integer.} \end{cases}$$

Clearly $P_1(x) = \left(\left(x \right) \right)$. He defined the generalized homogeneous Dedekind sum by

$$s_{\alpha, \beta}(a, b, q) = \sum_{j=1}^q P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right),$$

and proved the generalized Knopp identity for $s_{\alpha,\beta}(a, b, q)$ as follows:

$$(1.1) \quad \sum_{d|n} d^{\alpha+\beta-2} \sum_{r_1=1}^d \sum_{r_2=1}^d s_{\alpha,\beta} \left(\frac{n}{d}a + r_1q, \frac{n}{d}b + r_2q, dq \right) = n\sigma_{\alpha+\beta-1}(n)s_{\alpha,\beta}(a, b, q),$$

where $\sigma_r(n) = \sum_{d|n} d^r$.

In [2], Berndt studied the following Hardy sums:

$$\begin{aligned} S(h, k) &= \sum_{j=1}^{k-1} (-1)^{j+1+[hj/k]}, & s_1(h, k) &= \sum_{j=1}^k (-1)^{[hj/k]} \left(\left(\frac{j}{k} \right) \right), \\ s_2(h, k) &= \sum_{j=1}^k (-1)^j \left(\left(\frac{j}{k} \right) \right) \left(\left(\frac{hj}{k} \right) \right), & s_3(h, k) &= \sum_{j=1}^k (-1)^j \left(\left(\frac{hj}{k} \right) \right), \\ s_4(h, k) &= \sum_{j=1}^{k-1} (-1)^{[hj/k]}, & s_5(h, k) &= \sum_{j=1}^k (-1)^{j+[hj/k]} \left(\left(\frac{j}{k} \right) \right), \end{aligned}$$

which are related to the classical Dedekind sums, and obtained some arithmetic properties (see [3]). Sitaramachandrarao [10] and Pettet [8] expressed the Hardy sums in terms of the Dedekind sums as follows:

$$\begin{aligned} S(h, k) &= 8s(h, 2k) + 8s(2h, k) - 20s(h, k), & \text{if } h+k \text{ is odd;} \\ s_1(h, k) &= 2s(h, k) - 4s(h, 2k), & \text{if } h \text{ is even;} \\ s_2(h, k) &= -s(h, k) + 2s(2h, k), & \text{if } k \text{ is even;} \\ s_3(h, k) &= 2s(h, k) - 4s(2h, k), & \text{if } k \text{ is odd;} \\ s_4(h, k) &= -4s(h, k) + 8s(h, 2k), & \text{if } h \text{ is odd;} \\ s_5(h, k) &= -10s(h, k) + 4s(2h, k) + 4s(h, 2k), & \text{if } h+k \text{ is even.} \end{aligned}$$

Naturally, one might ask whether there is an analogous class of Hardy sums for the generalized homogeneous Dedekind sums? If yes, then what results can be expected?

We define the generalized homogeneous Hardy sums as follows:

$$\begin{aligned} s_{\beta}^{(1)}(a, b, q) &= \sum_{j=1}^q (-1)^{[aj/q]} P_{\beta} \left(\frac{bj}{q} \right), & \beta &\equiv 1 \pmod{2}; \\ s_{\alpha,\beta}^{(2)}(a, b, q) &= \sum_{j=1}^q (-1)^j P_{\alpha} \left(\frac{aj}{q} \right) P_{\beta} \left(\frac{bj}{q} \right), & \alpha &\equiv \beta \pmod{2}; \\ s_{\beta}^{(5)}(a, b, q) &= \sum_{j=1}^q (-1)^{j+[aj/q]} P_{\beta} \left(\frac{bj}{q} \right), & \beta &\equiv 1 \pmod{2}. \end{aligned}$$

We will express the generalized homogeneous Hardy sums in terms of the generalized homogeneous Dedekind sums in Section 2, and give the generalized Knopp identities for the homogeneous Hardy sums in Section 3. Next we study the properties of the generalized homogeneous Cochrane-Hardy sums. Finally in Section 5, the generalized Knopp identities for the generalized homogeneous Cochrane-Hardy sums are given.

2. THE GENERALIZED HOMOGENEOUS HARDY SUMS

In this section, we shall express the generalized homogeneous Hardy sums in terms of the generalized homogeneous Dedekind sums. First we need the following lemmas.

Lemma 2.1. *If m , a and q are integers with $m \geq 0$ and $q > 0$, then for any real x , we have*

$$\sum_{r=1}^q P_m \left(x + \frac{ar}{q} \right) = (a, q)^m q^{1-m} P_m \left(\frac{qx}{(a, q)} \right).$$

Proof. This is Lemma 3 of [16]. □

Lemma 2.2. *For any non-negative integers α and β , and any positive integer k , we have*

- (i) $s_{\alpha, \beta}(ak, b, qk) = (b, k)^\beta k^{1-\beta} s_{\alpha, \beta}(a, b/(b, k), q)$;
- (ii) $s_{\alpha, \beta}(a, bk, qk) = (a, k)^\alpha k^{1-\alpha} s_{\alpha, \beta}(a/(a, k), b, q)$;
- (iii) $s_{\alpha, \beta}(ak, bk, qk) = k s_{\alpha, \beta}(a, b, q)$.

Proof. This is Lemma 4 of [16]. □

From Lemma 2.1 and Lemma 2.2 we can get the following:

Lemma 2.3. *For any prime p , we have*

$$\sum_{r=0}^{p-1} s_{\alpha, \beta}(a + rq, b, pq) = p^{1-\alpha} s_{\alpha, \beta}(ap, b, pq) - p^{1-\alpha} s_{\alpha, \beta}(ap, b, q) + p s_{\alpha, \beta}(a, b, q).$$

For the special case $p = 2$, from Lemma 2.2 and Lemma 2.3 we immediately get

$$\begin{aligned} s_{\alpha, \beta}(a + q, b, 2q) &= 2^{2-\alpha-\beta} (b, 2)^\beta s_{\alpha, \beta}(a, b/(b, 2), q) - 2^{1-\alpha} s_{\alpha, \beta}(2a, b, q) \\ &\quad + 2 s_{\alpha, \beta}(a, b, q) - s_{\alpha, \beta}(a, b, 2q). \end{aligned}$$

Proof. By Lemma 2.1 we easily get

$$\begin{aligned}
 \sum_{r=0}^{p-1} s_{\alpha,\beta}(a+rq, b, pq) &= \sum_{r=0}^{p-1} \sum_{j=1}^{pq} P_{\alpha}\left(\frac{(a+rq)j}{pq}\right) P_{\beta}\left(\frac{bj}{pq}\right) \\
 &= \sum_{j=1}^{pq} P_{\beta}\left(\frac{bj}{pq}\right) \sum_{r=0}^{p-1} P_{\alpha}\left(\frac{aj}{pq} + \frac{rj}{p}\right) \\
 &= p^{1-\alpha} \sum_{j=1}^{pq} (j, p)^{\alpha} P_{\alpha}\left(\frac{aj}{(j, p)q}\right) P_{\beta}\left(\frac{bj}{pq}\right) \\
 &= p^{1-\alpha} \sum_{\substack{j=1 \\ (j, p)=1}}^{pq} P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{pq}\right) + p \sum_{\substack{j=1 \\ (j, p)=p}}^{pq} P_{\alpha}\left(\frac{aj}{pq}\right) P_{\beta}\left(\frac{bj}{pq}\right) \\
 &= p^{1-\alpha} \left[\sum_{j=1}^{pq} P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{pq}\right) - \sum_{j=1}^q P_{\alpha}\left(\frac{apj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) \right] \\
 &\quad + p \sum_{j=1}^q P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) \\
 &= p^{1-\alpha} s_{\alpha,\beta}(ap, b, pq) - p^{1-\alpha} s_{\alpha,\beta}(ap, b, q) + p s_{\alpha,\beta}(a, b, q).
 \end{aligned}$$

This proves Lemma 2.3. □

Now we shall prove the following:

Theorem 2.1. *Let $(ab, q) = 1$. Then we have*

$$(2.1) \quad s_{\beta}^{(1)}(a, b, q) = 2s_{1,\beta}(a, b, q) - 2^{\beta+1}(b, 2)^{-\beta} s_{1,\beta}(a, b(b, 2), 2q),$$

if a is even;

$$(2.2) \quad s_{\alpha,\beta}^{(2)}(a, b, q) = 2^{\beta}(b, 2)^{-\beta} s_{\alpha,\beta}(2a, b(b, 2), q) - s_{\alpha,\beta}(a, b, q),$$

if q is even;

$$(2.3) \quad s_{\beta}^{(5)}(a, b, q) = -2s_{1,\beta}(a, b, q) - 2^{\beta+2}(b, 2)^{-\beta} s_{1,\beta}(a, b(b, 2), q)$$

$$+ 2^{\beta+1}(b, 2)^{-\beta} s_{1,\beta}(2a, b(b, 2), q)$$

$$+ 2^{\beta+1}(b, 2)^{-\beta} s_{1,\beta}(a, b(b, 2), 2q),$$

if $a + q$ is even.

Moreover, each one of

$$(2.4) \quad s_{\beta}^{(1)}(a, b, q) \quad (a \text{ odd}), \quad s_{\alpha,\beta}^{(2)}(a, b, q) \quad (q \text{ odd}), \quad s_{\beta}^{(5)}(a, b, q) \quad (a + q \text{ odd})$$

is zero.

Proof. By (5.8) of [10] we know that

$$(2.5) \quad (-1)^{[x]} = 2\binom{x}{1} - 4\binom{x}{2}, \quad \text{if } x \text{ is not an integer.}$$

Thus for any even number a , by Lemma 2.2 we have

$$\begin{aligned} s_{\beta}^{(1)}(a, b, q) &= \sum_{j=1}^q (-1)^{[aj/q]} P_{\beta}\left(\frac{bj}{q}\right) = 2 \sum_{j=1}^{q-1} \left(\binom{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) - 4 \sum_{j=1}^{q-1} \left(\binom{aj}{2q}\right) P_{\beta}\left(\frac{bj}{q}\right) \\ &= 2s_{1,\beta}(a, b, q) - 4s_{1,\beta}\left(\frac{a}{2}, b, q\right) \\ &= 2s_{1,\beta}(a, b, q) - 2^{\beta+1}(b, 2)^{-\beta} s_{1,\beta}(a, b(b, 2), 2q). \end{aligned}$$

This proves (2.1).

If q is even, then by Lemma 2.2 we also have

$$\begin{aligned} s_{\alpha,\beta}^{(2)}(a, b, q) &= \sum_{j=1}^q (-1)^j P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) \\ &= 2 \sum_{\substack{j=1 \\ 2|j}}^q P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) - \sum_{j=1}^q P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) \\ &= 2 \sum_{j=1}^{q/2} P_{\alpha}\left(\frac{2aj}{q}\right) P_{\beta}\left(\frac{2bj}{q}\right) - s_{\alpha,\beta}(a, b, q) \\ &= 2s_{\alpha,\beta}\left(a, b, \frac{q}{2}\right) - s_{\alpha,\beta}(a, b, q) \\ &= 2^{\beta}(b, 2)^{-\beta} s_{\alpha,\beta}(2a, b(b, 2), q) - s_{\alpha,\beta}(a, b, q). \end{aligned}$$

This proves (2.2).

If $a + q$ is even, from Lemma 2.2 and Lemma 2.3 we get

$$\begin{aligned} s_{\beta}^{(5)}(a, b, q) &= \sum_{j=1}^q (-1)^{j+[aj/q]} P_{\beta}\left(\frac{bj}{q}\right) = \sum_{j=1}^q (-1)^{[j(a+q)/q]} P_{\beta}\left(\frac{bj}{q}\right) \\ &= \sum_{j=1}^{q-1} \left[2\binom{j(a+q)}{q} - 4\binom{j(a+q)}{2q} \right] P_{\beta}\left(\frac{bj}{q}\right) \\ &= 2s_{1,\beta}(a, b, q) - 4s_{1,\beta}\left(\frac{a+q}{2}, b, q\right) \\ &= 2s_{1,\beta}(a, b, q) - 2^{\beta+1}(b, 2)^{-\beta} s_{1,\beta}(a+q, b(b, 2), 2q) \\ &= -2s_{1,\beta}(a, b, q) - 2^{\beta+2}(b, 2)^{-\beta} s_{1,\beta}(a, b(b, 2), q) \\ &\quad + 2^{\beta+1}(b, 2)^{-\beta} s_{1,\beta}(2a, b(b, 2), q) + 2^{\beta+1}(b, 2)^{-\beta} s_{1,\beta}(a, b(b, 2), 2q). \end{aligned}$$

This proves (2.3).

To prove (2.4), noting that

$$(2.6) \quad P_m(-x) = (-1)^m P_m(x),$$

we have from the definition of $s_\beta^{(1)}(a, b, q)$

$$\begin{aligned} s_\beta^{(1)}(a, b, q) &= \sum_{j=1}^{q-1} (-1)^{[aj/q]} P_\beta\left(\frac{bj}{q}\right) = \sum_{k=1}^{q-1} (-1)^{[a(q-k)/q]} P_\beta\left(\frac{b(q-k)}{q}\right) \\ &= \sum_{k=1}^{q-1} (-1)^{a+[-ak/q]} P_\beta\left(-\frac{bk}{q}\right) = \sum_{k=1}^{q-1} (-1)^{a-[ak/q]-1} P_\beta\left(-\frac{bk}{q}\right) \\ &= (-1)^{a+\beta-1} \sum_{k=1}^{q-1} (-1)^{[ak/q]} P_\beta\left(\frac{bk}{q}\right) = (-1)^a \sum_{k=1}^{q-1} (-1)^{[ak/q]} P_\beta\left(\frac{bk}{q}\right) \\ &= (-1)^a s_\beta^{(1)}(a, b, q). \end{aligned}$$

Thus $s_\beta^{(1)}(a, b, q) = 0$ if a is odd. The proofs of the remaining assertions in (2.4) are similar. \square

Remark. Taking $\alpha = \beta = b = 1$ in Theorem 2.1, we immediately get (1.3), (1.4) and (1.7).

3. GENERALIZED KNOPP IDENTITIES FOR THE GENERALIZED HOMOGENEOUS HARDY SUMS

In this section, we shall give the generalized Knopp identities for the generalized homogeneous Hardy sums.

Theorem 3.1. *Let n be a positive odd integer, and $(a, q) = 1$. Then we have*

$$(3.1) \quad \begin{aligned} \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d s_\beta^{(1)}\left(\frac{n}{d}a + 2r_1q, \frac{n}{d}b + r_2q, dq\right) \\ = n\sigma_\beta(n) s_\beta^{(1)}(a, b, q), \quad \text{if } a \text{ is even;} \end{aligned}$$

$$(3.2) \quad \begin{aligned} \sum_{d|n} d^{\alpha+\beta-2} \sum_{r_1=1}^d \sum_{r_2=1}^d s_{\alpha,\beta}^{(2)}\left(\frac{n}{d}a + r_1q, \frac{n}{d}b + r_2q, dq\right) \\ = n\sigma_{\alpha+\beta-1}(n) s_{\alpha,\beta}^{(2)}(a, b, q), \quad \text{if } q \text{ is even;} \end{aligned}$$

$$(3.3) \quad \begin{aligned} \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d s_\beta^{(5)}\left(\frac{n}{d}a + 2r_1q, \frac{n}{d}b + r_2q, dq\right) \\ = n\sigma_\beta(n) s_\beta^{(5)}(a, b, q), \quad \text{if } a + q \text{ is even.} \end{aligned}$$

Proof. We only prove (3.1). The other identities can be deduced similarly. By Theorem 2.1 and (1.1) we have

$$\begin{aligned}
& \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d s_{\beta}^{(1)}\left(\frac{n}{d}a + 2r_1q, \frac{n}{d}b + r_2q, dq\right) \\
&= \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d \left[2s_{1,\beta}\left(\frac{n}{d}a + 2r_1q, \frac{n}{d}b + r_2q, dq\right) \right. \\
&\quad \left. - 4s_{1,\beta}\left(\frac{n}{d} \cdot \frac{a}{2} + r_1q, \frac{n}{d}b + r_2q, dq\right) \right] \\
&= 2 \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d s_{1,\beta}\left(\frac{n}{d}a + r_1q, \frac{n}{d}b + r_2q, dq\right) \\
&\quad - 4 \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d s_{1,\beta}\left(\frac{n}{d} \cdot \frac{a}{2} + r_1q, \frac{n}{d}b + r_2q, dq\right) \\
&= 2n\sigma_{\beta}(n)s_{1,\beta}(a, b, q) - 4n\sigma_{\beta}(n)s_{1,\beta}\left(\frac{a}{2}, b, q\right) = n\sigma_{\beta}(n)s_{\beta}^{(1)}(a, b, q).
\end{aligned}$$

This completes the proof of Theorem 3.1. □

4. GENERALIZED HOMOGENEOUS COCHRANE SUMS AND COCHRANE-HARDY SUMS

For a positive integer q and an arbitrary integer h , the Cochrane sums are defined by

$$c(h, q) = \sum'_{a=1}^q \left(\left(\frac{a}{q} \right) \right) \left(\left(\frac{\bar{a}h}{q} \right) \right),$$

where \sum'_a denotes the summation over all a such that $(a, q) = 1$, and \bar{a} is defined by the equation $a\bar{a} \equiv 1 \pmod{q}$. Wenpeng Zhang and Yuan Yi [14] gave the following upper bound estimate

$$|c(h, q)| \ll \sqrt{q}d(q) \ln^2 q$$

and the mean value estimate

$$\sum_{h=1}^{p-1} c^2(h, p) = \frac{5}{144}p^2 + O\left(p \exp\left(\frac{4 \ln p}{\ln \ln p}\right)\right),$$

where $d(q)$ is the divisor function, and $\exp(y) = e^y$.

In [11], Wenpeng Zhang found that there are some relationships between $c(h, q)$ and the Kloosterman sum

$$K(m, n; q) = \sum_{a=1}^q{}' e\left(\frac{ma + n\bar{a}}{q}\right),$$

where $e(y) = e^{2\pi iy}$. For example, if q is a square-full number, he obtained

$$\sum_{h=1}^q{}' c(h, q)K(h, 1; q) = -\frac{1}{2\pi^2}q\varphi(q) + O\left(q \exp\left(\frac{3 \ln q}{\ln \ln q}\right)\right),$$

where $\varphi(q)$ is the Euler function. For general integer $q \geq 3$, he obtained in [12] the asymptotic formula

$$(4.1) \quad \sum_{h=1}^q{}' c(h, q)K(h, 1; q) = -\frac{1}{2\pi^2}q\varphi(q) \prod_{p||q} \left(1 - \frac{1}{p(p-1)}\right) + O(q^{\frac{3}{2}+\varepsilon}),$$

where ε is any fixed positive number. Later he and the author [13] proved that the error term in (4.1) is $O(q^{1+\varepsilon})$.

We turn now to the study of the generalized homogeneous Cochrane sums and Cochrane-Hardy sums. First we give the following definitions:

$$\begin{aligned} c_{\alpha, \beta}(a, b, q) &= \sum_{j=1}^q{}' P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{b\bar{j}}{q}\right), & \alpha &\equiv \beta \pmod{2}; \\ c_{\beta}^{(1)}(a, b, q) &= \sum_{j=1}^q{}' (-1)^{[aj/q]} P_{\beta}\left(\frac{b\bar{j}}{q}\right), & \beta &\equiv 1 \pmod{2}; \\ c_{\alpha, \beta}^{(2)}(a, b, q) &= \sum_{j=1}^q{}' (-1)^j P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{b\bar{j}}{q}\right), & \alpha &\equiv \beta \pmod{2}; \\ c_{\beta}^{(5)}(a, b, q) &= \sum_{j=1}^q{}' (-1)^{j+[aj/q]} P_{\beta}\left(\frac{b\bar{j}}{q}\right), & \beta &\equiv 1 \pmod{2}. \end{aligned}$$

Now we express the generalized homogeneous Cochrane-Hardy sums in terms of generalized homogeneous Cochrane sums.

Theorem 4.1. *Let $(ab, q) = 1$. Then we have*

$$(4.2) \quad c_{\beta}^{(1)}(a, b, q) = 2c_{1,\beta}(a, b, q) - 4c_{1,\beta}\left(\frac{a}{2}, b, q\right), \quad \text{if } a \text{ is even;}$$

$$(4.3) \quad c_{\alpha,\beta}^{(2)}(a, b, q) = -c_{\alpha,\beta}(a, b, q), \quad \text{if } q \text{ is even;}$$

$$(4.4) \quad c_{\beta}^{(5)}(a, b, q) = 2c_{1,\beta}(a, b, q) - 4c_{1,\beta}\left(\frac{a+q}{2}, b, q\right), \quad \text{if } a+q \text{ is even.}$$

Moreover, each one of

$$(4.5) \quad c_{\beta}^{(1)}(a, b, q) \quad (a \text{ odd}), \quad c_{\alpha,\beta}^{(2)}(a, b, q) \quad (q \text{ odd}), \quad c_{\beta}^{(5)}(a, b, q) \quad (a+q \text{ odd})$$

is zero.

Proof. For any even number a , by (2.5) we have

$$\begin{aligned} c_{\beta}^{(1)}(a, b, q) &= \sum_{j=1}^q (-1)^{[aj/q]} P_{\beta}\left(\frac{bj}{q}\right) \\ &= 2 \sum_{j=1}^q \left(\left(\frac{aj}{q}\right)\right) P_{\beta}\left(\frac{bj}{q}\right) - 4 \sum_{j=1}^q \left(\left(\frac{aj}{2q}\right)\right) P_{\beta}\left(\frac{bj}{q}\right) \\ &= 2c_{1,\beta}(a, b, q) - 4c_{1,\beta}\left(\frac{a}{2}, b, q\right). \end{aligned}$$

This proves (4.2).

If q is even, then $(j, q) = 1$ only if j is odd. Therefore

$$\begin{aligned} c_{\alpha,\beta}^{(2)}(a, b, q) &= \sum_{j=1}^q (-1)^j P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) \\ &= - \sum_{j=1}^q P_{\alpha}\left(\frac{aj}{q}\right) P_{\beta}\left(\frac{bj}{q}\right) = -c_{\alpha,\beta}(a, b, q). \end{aligned}$$

This proves (4.3).

If $a+q$ is even, by (2.5) we get

$$\begin{aligned} c_{\beta}^{(5)}(a, b, q) &= \sum_{j=1}^q (-1)^{j+[aj/q]} P_{\beta}\left(\frac{bj}{q}\right) = \sum_{j=1}^q (-1)^{[j(a+q)/q]} P_{\beta}\left(\frac{bj}{q}\right) \\ &= \sum_{j=1}^q \left[2 \left(\left(\frac{j(a+q)}{q}\right)\right) - 4 \left(\left(\frac{j(a+q)}{2q}\right)\right) \right] P_{\beta}\left(\frac{bj}{q}\right) \\ &= 2c_{1,\beta}(a, b, q) - 4c_{1,\beta}\left(\frac{a+q}{2}, b, q\right). \end{aligned}$$

This proves (4.4).

To prove (4.5), from (2.6) and the definition of $c_\beta^{(1)}(a, b, q)$ we have

$$\begin{aligned} c_\beta^{(1)}(a, b, q) &= \sum_{j=1}^q (-1)^{[aj/q]} P_\beta\left(\frac{b\bar{j}}{q}\right) = \sum_{j=1}^q (-1)^{[a(q-j)/q]} P_\beta\left(\frac{b\overline{q-j}}{q}\right) \\ &= \sum_{j=1}^q (-1)^{a-[aj/q]-1} P_\beta\left(-\frac{b\bar{j}}{q}\right) = (-1)^{a+\beta-1} \sum_{j=1}^q (-1)^{[aj/q]} P_\beta\left(\frac{b\bar{j}}{q}\right) \\ &= (-1)^a c_\beta^{(1)}(a, b, q). \end{aligned}$$

Thus $c_\beta^{(1)}(a, b, q) = 0$ if a is odd. The proofs of the remaining assertions in (4.5) are similar. \square

5. GENERALIZED KNOPP IDENTITIES FOR THE GENERALIZED HOMOGENEOUS COCHRANE SUMS AND COCHRANE-HARDY SUMS

First we prove the generalized Knopp identity for the generalized homogeneous Cochrane sums as follows:

Theorem 5.1. *Let $(abn, q) = 1$. Then we have*

$$\sum_{d|n} d^{\alpha+\beta-2} \sum_{r_1=1}^d \sum_{r_2=1}^d c_{\alpha,\beta}(a+r_1q, b+r_2q, dq) = nc_{\alpha,\beta}(a, b, q).$$

Proof. By Lemma 2.1 we get

$$\begin{aligned} (5.1) \quad & \sum_{d|n} d^{\alpha+\beta-2} \sum_{r_1=1}^d \sum_{r_2=1}^d c_{\alpha,\beta}(a+r_1q, b+r_2q, dq) \\ &= \sum_{d|n} d^{\alpha+\beta-2} \sum_{r_1=1}^d \sum_{r_2=1}^d \sum_{r=1}^{dq} P_\alpha\left(\frac{(a+r_1q)r}{dq}\right) P_\beta\left(\frac{(b+r_2q)\bar{r}}{dq}\right) \\ &= \sum_{d|n} d^{\alpha+\beta-2} \sum_{r=1}^{dq} \sum_{r_1=1}^d P_\alpha\left(\frac{ar}{dq} + \frac{r_1r}{d}\right) \sum_{r_2=1}^d P_\beta\left(\frac{b\bar{r}}{dq} + \frac{r_2\bar{r}}{d}\right) \\ &= \sum_{d|n} \sum_{r=1}^{dq} P_\alpha\left(\frac{ar}{q}\right) P_\beta\left(\frac{b\bar{r}}{q}\right). \end{aligned}$$

Since $(n, q) = 1$, we have $(d, q) = 1$. Let $r = xd + yq$ with $x = 1, \dots, q$, $(x, q) = 1$, $y = 1, \dots, d$, $(y, d) = 1$. Then

$$\bar{r} = \bar{x}d^2d + \bar{y}q^2q,$$

where

$$d\bar{d} \equiv 1 \pmod{q}, \quad q\bar{q} \equiv 1 \pmod{d}, \quad x\bar{x} \equiv 1 \pmod{q}, \quad y\bar{y} \equiv 1 \pmod{d}.$$

Therefore

$$\begin{aligned} (5.2) \quad & \sum_{d|n} \sum_{r=1}^{dq}' P_{\alpha}\left(\frac{ar}{q}\right) P_{\beta}\left(\frac{b\bar{r}}{q}\right) \\ &= \sum_{d|n} \sum_{x=1}^q \sum_{y=1}^d P_{\alpha}\left(\frac{a(xd+yq)}{q}\right) P_{\beta}\left(\frac{b(\bar{x}\bar{d}^2d+\bar{y}\bar{q}^2q)}{q}\right) \\ &= \sum_{d|n} \sum_{x=1}^q \sum_{y=1}^d P_{\alpha}\left(\frac{axd}{q}\right) P_{\beta}\left(\frac{b\bar{x}\bar{d}}{q}\right) \\ &= \sum_{d|n} \varphi(d) \sum_{x=1}^q P_{\alpha}\left(\frac{axd}{q}\right) P_{\beta}\left(\frac{b\bar{x}\bar{d}}{q}\right). \end{aligned}$$

As x runs over a reduced residue system modulo q , xd also runs over a reduced residue system modulo q . Thus we have

$$\begin{aligned} (5.3) \quad & \sum_{d|n} \varphi(d) \sum_{x=1}^q P_{\alpha}\left(\frac{axd}{q}\right) P_{\beta}\left(\frac{b\bar{x}\bar{d}}{q}\right) = \sum_{d|n} \varphi(d) \sum_{x=1}^q P_{\alpha}\left(\frac{ax}{q}\right) P_{\beta}\left(\frac{b\bar{x}}{q}\right) \\ &= nc_{\alpha,\beta}(a, b, q). \end{aligned}$$

Combining (5.1)–(5.3), we immediately get Theorem 5.1. \square

Using the same methods and Theorem 4.1 we also get the following generalized Knopp identities for the generalized homogeneous Cochrane-Hardy sums.

Theorem 5.2. *Let n be a positive odd integer with $(abn, q) = 1$. Then we have*

$$\begin{aligned} & \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d c_{\beta}^{(1)}(a+2r_1q, b+r_2q, dq) = nc_{\beta}^{(1)}(a, b, q), \quad \text{if } a \text{ is even;} \\ & \sum_{d|n} d^{\alpha+\beta-2} \sum_{r_1=1}^d \sum_{r_2=1}^d c_{\alpha,\beta}^{(2)}(a+r_1q, b+r_2q, dq) = nc_{\alpha,\beta}^{(2)}(a, b, q), \quad \text{if } q \text{ is even;} \\ & \sum_{d|n} d^{\beta-1} \sum_{r_1=1}^d \sum_{r_2=1}^d c_{\beta}^{(5)}(a+2r_1q, b+r_2q, dq) = nc_{\beta}^{(5)}(a, b, q), \quad \text{if } a+q \text{ is even.} \end{aligned}$$

References

- [1] *T. M. Apostol*: Modular Functions and Dirichlet Series in Number Theory. Springer, New York, Heidelberg, Berlin, 1976.
- [2] *B. C. Berndt*: Analytic Eisenstein series, theta-functions, and series relations in the spirit of Ramanujan. *J. Reine Angew. Math.* *303/304* (1978), 332–365.
- [3] *B. C. Berndt, L. A. Goldberg*: Analytic properties of arithmetic sums arising in the theory of the classical theta-functions. *SIAM J. Math. Anal.* *15* (1984), 143–150.
- [4] *L. A. Goldberg*: An elementary proof of the Petersson-Knopp theorem on Dedekind sums. *J. Number Theory* *12* (1980), 541–542.
- [5] *R. R. Hall, M. N. Huxley*: Dedekind sums and continued fractions. *Acta Arith.* *63* (1993), 79–90.
- [6] *M. I. Knopp*: Hecke operators and an identity for the Dedekind sums. *J. Number Theory* *12* (1980), 2–9.
- [7] *L. A. Parson*: Dedekind sums and Hecke operators. *Math. Proc. Camb. Philos. Soc.* *88* (1980), 11–14.
- [8] *M. R. Pettet, R. Sitaramachandrarao*: Three-term relations for Hardy sums. *J. Number Theory* *25* (1987), 328–339.
- [9] *H. Rademacher, E. Grosswald*: Dedekind Sums. The Carus Mathematical Monographs No. 16, The Mathematical Association of America, Washington, D. C., 1972.
- [10] *R. Sitaramachandrarao*: Dedekind and Hardy sums. *Acta Arith.* *48* (1987), 325–340.
- [11] *W. Zhang*: On a Cochrane sum and its hybrid mean value formula. *J. Math. Anal. Appl.* *267* (2002), 89–96.
- [12] *W. Zhang*: On a Cochrane sum and its hybrid mean value formula. II. *J. Math. Anal. Appl.* *276* (2002), 446–457.
- [13] *W. Zhang, H. Liu*: A note on the Cochrane sum and its hybrid mean value formula. *J. Math. Anal. Appl.* *288* (2003), 646–659.
- [14] *W. Zhang, Y. Yi*: On the upper bound estimate of Cochrane sums. *Soochow J. Math.* *28* (2002), 297–304.
- [15] *Z. Zheng*: On an identity for Dedekind sums. *Acta Math. Sin.* *37* (1994), 690–694.
- [16] *Z. Zheng*: The Petersson-Knopp identity for homogeneous Dedekind sums. *J. Number Theory* *57* (1996), 223–230.

Authors' addresses: Hu aning Liu, Department of Mathematics, Northwest University, Xi'an 710069, Shaanxi, P.R. China, e-mail: hnliumath@hotmail.com; Jing Gao, School of Mathematics and Statistics, Xi'an Jiaotong University, Xi'an 710049, Shaanxi, P.R. China, e-mail: jgao@mail.xjtu.edu.cn.