

Luong Quoc Tuyen

Remarks on sequence-covering maps

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 53 (2012), No. 4, 645--650

Persistent URL: <http://dml.cz/dmlcz/143197>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Remarks on sequence-covering maps

LUONG QUOC TUYEN

*Abstract.* In this paper, we prove that each sequence-covering and boundary-compact map on  $g$ -metrizable spaces is 1-sequence-covering. Then, we give some relationships between sequence-covering maps and 1-sequence-covering maps or weak-open maps, and give an affirmative answer to the problem posed by F.C. Lin and S. Lin in [9].

*Keywords:*  $g$ -metrizable space, weak base,  $sn$ -network, compact map, boundary-compact map, sequence-covering map, 1-sequence-covering map, weak-open map, closed map

*Classification:* 54C10, 54D65, 54E40, 54E99

### 1. Introduction

A study of images of topological spaces under certain sequence-covering maps is an important question in general topology ([1], [2], [8]–[11], [13], [16], [21], for example). In 2000, P. Yan, S. Lin and S.L. Jiang proved that each closed sequence-covering map on metric spaces is 1-sequence-covering ([21]). Furthermore, in 2001, S. Lin and P. Yan proved that each sequence-covering and compact map on metric spaces is 1-sequence-covering ([13]). After that, T.V. An and L.Q. Tuyen proved that each sequence-covering  $\pi$  and  $s$ -map on metric spaces is 1-sequence-covering ([1]). Recently, F.C. Lin and S. Lin proved that each sequence-covering and boundary-compact map on metric spaces is 1-sequence-covering ([8]). Also, the authors posed the following question in [9].

**Question 1.1** ([9, Question 4.6]). *Let  $f : X \rightarrow Y$  be a sequence-covering and boundary-compact map. If  $X$  is  $g$ -metrizable, then is  $f$  an 1-sequence-covering map?*

In this paper, we prove that each sequence-covering and boundary-compact map on  $g$ -metrizable spaces is 1-sequence-covering. Then, we give some relationships between sequence-covering maps and 1-sequence-covering maps or weak-open maps, and give an affirmative answer to the problem posed by F.C. Lin and S. Lin in [9].

Throughout this paper, all spaces are assumed to be Hausdorff, all maps are continuous and onto,  $\mathbb{N}$  denotes the set of all natural numbers,  $\omega$  denotes  $\mathbb{N} \cup \{0\}$ , and convergent sequence includes its limit point. Let  $f : X \rightarrow Y$  be a map and  $\mathcal{P}$  be a collection of subsets of  $X$ , we denote  $\bigcup \mathcal{P} = \bigcup \{P : P \in \mathcal{P}\}$ ,  $\bigcap \mathcal{P} = \bigcap \{P : P \in \mathcal{P}\}$ ,  $f(\mathcal{P}) = \{f(P) : P \in \mathcal{P}\}$ .

**Definition 1.2.** Let  $X$  be a space, and  $P \subset X$ .

- (1) A sequence  $\{x_n\}$  in  $X$  is called *eventually* in  $P$ , if  $\{x_n\}$  converges to  $x$ , and there exists  $m \in \mathbb{N}$  such that  $\{x\} \cup \{x_n : n \geq m\} \subset P$ .
- (2)  $P$  is called a *sequential neighborhood* of  $x$  in  $X$  [5], if whenever  $\{x_n\}$  is a sequence converging to  $x$  in  $X$ , then  $\{x_n\}$  is eventually in  $P$ .

**Definition 1.3.** Let  $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$  be a cover of a space  $X$ . Assume that  $\mathcal{P}$  satisfies the following conditions (a) and (b) for every  $x \in X$ .

- (a)  $\mathcal{P}_x$  is a network at  $x$ .
- (b) If  $P_1, P_2 \in \mathcal{P}_x$ , then there exists  $P \in \mathcal{P}_x$  such that  $P \subset P_1 \cap P_2$ .
- (1)  $\mathcal{P}$  is a *weak base* of  $X$  [3], if for  $G \subset X$ ,  $G$  is open in  $X$  and every  $x \in G$ , there exists  $P \in \mathcal{P}_x$  such that  $P \subset G$ ;  $\mathcal{P}_x$  is said to be a weak neighborhood base at  $x$ .
- (2)  $\mathcal{P}$  is an *sn-network* for  $X$  [10], if each element of  $\mathcal{P}_x$  is a sequential neighborhood of  $x$  for all  $x \in X$ ;  $\mathcal{P}_x$  is said to be an *sn-network* at  $x$ .

**Definition 1.4.** Let  $X$  be a space. Then

- (1)  $X$  is *gf-countable* [3] (resp., *snf-countable* [6]), if  $X$  has a weak base (resp., *sn-network*)  $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$  such that each  $\mathcal{P}_x$  is countable;
- (2)  $X$  is *g-metrizable* [17], if  $X$  is regular and has a  $\sigma$ -locally finite weak base;
- (3)  $X$  is *sequential* [5], if whenever  $A$  is a non-closed subset of  $X$ , then there is a sequence in  $A$  converging to a point not in  $A$ .

*Remark 1.5.* (1) Each *g-metrizable* space or *gf-countable* space is sequential.  
 (2) A space  $X$  is *gf-countable* if and only if it is sequential and *snf-countable*.

**Definition 1.6.** Let  $f : X \rightarrow Y$  be a map. Then

- (1)  $f$  is a *compact* map [4], if each  $f^{-1}(y)$  is compact in  $X$ ;
- (2)  $f$  is a *boundary-compact* map [4], if each  $\partial f^{-1}(y)$  is compact in  $X$ ;
- (3)  $f$  is a *weak-open* map [18], if there exists a weak base  $\mathcal{P} = \bigcup\{\mathcal{P}_y : y \in Y\}$  for  $Y$ , and for  $y \in Y$ , there exists  $x_y \in f^{-1}(y)$  such that for each open neighborhood  $U$  of  $x_y$ ,  $P_y \subset f(U)$  for some  $P_y \in \mathcal{P}_y$ ;
- (4)  $f$  is an *1-sequence-covering* map [10], if for each  $y \in Y$ , there is  $x_y \in f^{-1}(y)$  such that whenever  $\{y_n\}$  is a sequence converging to  $y$  in  $Y$ , there is a sequence  $\{x_n\}$  converging to  $x_y$  in  $X$  with  $x_n \in f^{-1}(y_n)$  for every  $n \in \mathbb{N}$ ;
- (5)  $f$  is a *sequence-covering* map [16], if every convergent sequence of  $Y$  is the image of some convergent sequence of  $X$ ;
- (6)  $f$  is a *quotient* map [4], if whenever  $f^{-1}(U)$  is open in  $X$ , then  $U$  is open in  $Y$ .

*Remark 1.7.* (1) Each compact map is a compact-boundary map.  
 (2) Each 1-sequence-covering map is a sequence-covering map.  
 (3) Each closed map is a quotient map.

**Definition 1.8** ([7]). A function  $g : \mathbb{N} \times X \rightarrow \mathcal{P}(X)$  is a *CWC-map*, if it satisfies the following conditions:

- (1)  $x \in g(n, x)$  for all  $x \in X$  and  $n \in \mathbb{N}$ ;
- (2)  $g(n + 1, x) \subset g(n, x)$  for all  $n \in \mathbb{N}$ ;
- (3)  $\{g(n, x) : n \in \mathbb{N}\}$  is a weak neighborhood base at  $x$  for all  $x \in X$ .

**2. Main results**

**Theorem 2.1.** *Each sequence-covering and boundary-compact map on  $g$ -metrizable spaces is 1-sequence-covering.*

PROOF: Let  $f : X \rightarrow Y$  be a sequence-covering and boundary-compact map and  $X$  be a  $g$ -metrizable space. Firstly, we prove that  $Y$  is *snf*-countable. In fact, since  $X$  is  $g$ -metrizable, it follows from Theorem 2.6 in [20] that there exists a CWC-map  $g$  on  $X$  satisfying that  $y_n \rightarrow x$  whenever  $\{x_n\}, \{y_n\}$  are two sequences in  $X$  such that  $x_n \rightarrow x$  and  $y_n \in g(n, x_n)$  for all  $n \in \mathbb{N}$ . For each  $y \in Y$  and  $n \in \mathbb{N}$ , we put

$$P_{y,n} = f\left(\bigcup\{g(n, x) : x \in \partial f^{-1}(y)\}\right), \text{ and } \mathcal{P}_y = \{P_{y,n} : n \in \mathbb{N}\}.$$

Then each  $\mathcal{P}_y$  is countable and  $P_{y,n+1} \subset P_{y,n}$  for all  $y \in Y$  and  $n \in \mathbb{N}$ . Furthermore, we have

(1)  $\mathcal{P}_y$  is a network at  $y$ . Indeed, let  $y \in U$  with  $U$  open in  $Y$ . Then there exists  $n \in \mathbb{N}$  such that

$$\bigcup\{g(n, x) : x \in \partial f^{-1}(y)\} \subset f^{-1}(U).$$

If not, for each  $n \in \mathbb{N}$ , there exist  $x_n \in \partial f^{-1}(y)$  and  $z_n \in X$  such that  $z_n \in g(n, x_n) - f^{-1}(U)$ . Since  $X$  is  $g$ -metrizable, it follows that each compact subset of  $X$  is metrizable. Since  $\{x_n\} \subset \partial f^{-1}(y)$  and  $f$  is a boundary-compact map, there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  such that  $x_{n_k} \rightarrow x \in \partial f^{-1}(y)$ . Now, for each  $i \in \mathbb{N}$ , we put

$$a_i = \begin{cases} x_{n_1} & \text{if } i \leq n_1 \\ x_{n_{k+1}} & \text{if } n_k < i \leq n_{k+1}; \end{cases}$$

$$b_i = \begin{cases} z_{n_1} & \text{if } i \leq n_1 \\ z_{n_{k+1}} & \text{if } n_k < i \leq n_{k+1}. \end{cases}$$

Then  $a_i \rightarrow x$ . Because  $g(n + 1, x) \subset g(n, x)$  for all  $x \in X$  and  $n \in \mathbb{N}$ , it implies that  $b_i \in g(i, a_i)$  for all  $i \in \mathbb{N}$ . By the property of  $g$ , it implies that  $b_i \rightarrow x$ . Thus,  $z_{n_k} \rightarrow x$ . This contradicts that  $f^{-1}(U)$  is a neighborhood of  $x$  and  $z_{n_k} \notin f^{-1}(U)$  for all  $k \in \mathbb{N}$ . Therefore,  $P_{y,n} \subset U$ , and  $\mathcal{P}_y$  is a network at  $y$ .

(2) Let  $P_{y,m}, P_{y,n} \in \mathcal{P}_y$ . If we take  $k = \max\{m, n\}$ , then  $P_{y,k} \subset P_{y,m} \cap P_{y,n}$ .

(3) Each element of  $\mathcal{P}_y$  is a sequential neighborhood of  $y$ . Let  $P_{y,n} \in \mathcal{P}_y$  and  $\{y_n\}$  be a sequence converging to  $y$  in  $Y$ . Since  $f$  is sequence-covering,  $\{y_n\}$  is an image of some sequence converging to  $x \in \partial f^{-1}(y)$ . On the other hand, since  $g(n, x)$  is a weak neighborhood of  $x$ ,  $\{y_n\}$  is eventually in  $g(n, x)$ . This implies that  $\{y_n\}$  is eventually in  $P_{y,n}$ . Therefore,  $P_{y,n}$  is a sequential neighborhood of  $y$ .

Therefore,  $\bigcup\{\mathcal{P}_y : y \in Y\}$  is an *sn*-network for  $X$ , and  $Y$  is an *snf*-countable space.

Next, let  $\mathcal{B} = \bigcup\{B_x : x \in X\}$  be a  $\sigma$ -locally finite weak base for  $X$ . We prove that for each non-isolated point  $y \in Y$ , there exists  $x_y \in \partial f^{-1}(y)$  such that for each  $B \in \mathcal{B}_{x_y}$ , there exists  $P \in \mathcal{P}_y$  satisfying  $P \subset f(B)$ . Otherwise, there exists a non-isolated point  $y \in Y$  so that for each  $x \in \partial f^{-1}(y)$ , there exists  $B_x \in \mathcal{B}_x$  such that  $P \not\subset f(B_x)$  for all  $P \in \mathcal{P}_y$ . Since  $\mathcal{B}$  is a  $\sigma$ -locally finite weak base and  $\partial f^{-1}(y)$  is compact, it follows that  $\{B_x : x \in \partial f^{-1}(y)\}$  is countable. Assume that  $\{B_x : x \in \partial f^{-1}(y)\} = \{B_m : m \in \mathbb{N}\}$ . Hence, for each  $m, n \in \mathbb{N}$ , there exists  $x_{n,m} \in P_{y,n} - f(B_m)$ . For  $n \geq m$ , we denote  $y_k = x_{n,m}$  with  $k = m + n(n-1)/2$ . Since  $\mathcal{P}_y$  is a network at  $y$  and  $P_{y,n+1} \subset P_{y,n}$  for all  $n \in \mathbb{N}$ ,  $\{y_k\}$  is a sequence converging to  $y$  in  $Y$ . On the other hand, because  $f$  is a sequence-covering map,  $\{y_k\}$  is an image of some sequence  $\{x_n\}$  converging to  $x \in \partial f^{-1}(y)$  in  $X$ . Furthermore, since  $B_x \in \{B_m : m \in \mathbb{N}\}$ , there exists  $m_0 \in \mathbb{N}$  such that  $B_x = B_{m_0}$ . Because  $B_{m_0}$  is a weak neighborhood of  $x$ ,  $\{x\} \cup \{x_k : k \geq k_0\} \subset B_{m_0}$  for some  $k_0 \in \mathbb{N}$ . Thus,  $\{y\} \cup \{y_k : k \geq k_0\} \subset f(B_{m_0})$ . But if we take  $k \geq k_0$ , then there exists  $n \geq m_0$  such that  $y_k = x_{n,m_0}$ , and it implies that  $x_{n,m_0} \in f(B_{m_0})$ . This contradicts to  $x_{n,m_0} \in P_{y,n} - f(B_{m_0})$ .

We now prove that  $f$  is an 1-sequence-covering map. Suppose  $y \in Y$ . By the above proof there is  $x_y \in \partial f^{-1}(y)$  such that whenever  $B \in \mathcal{B}_{x_y}$ , there exists  $P \in \mathcal{P}_y$  satisfying  $P \subset f(B)$ . Let  $\{y_n\}$  be a sequence in  $Y$ , which converges to  $y$ . Since  $\mathcal{B}_{x_y}$  is a weak neighborhood base at  $x_y$ , we can choose a decreasing countable network  $\{B_{y,n} : n \in \mathbb{N}\} \subset \mathcal{B}_{x_y}$  at  $x_y$ . We choose a sequence  $\{z_n\}$  in  $X$  as follows.

Since  $B_{y,n} \in \mathcal{B}_{x_y}$ , by the above argument, there exists  $P_{y,k_n} \in \mathcal{P}_y$  satisfying  $P_{y,k_n} \subset f(B_{y,n})$  for all  $n \in \mathbb{N}$ . On the other hand, since each element of  $\mathcal{P}_y$  is a sequential neighborhood of  $y$ , it follows that for each  $n \in \mathbb{N}$ ,  $f(B_{y,n})$  is a sequential neighborhood of  $y$  in  $Y$ . Hence, for each  $n \in \mathbb{N}$ , there exists  $i_n \in \mathbb{N}$  such that  $y_i \in f(B_{y,n})$  for every  $i \geq i_n$ . Assume that  $1 < i_n < i_{n+1}$  for each  $n \in \mathbb{N}$ . Then for each  $j \in \mathbb{N}$ , we take

$$z_j = \begin{cases} z_j \in f^{-1}(y_j) & \text{if } j < i_1 \\ z_{j,n} \in f^{-1}(y_j) \cap B_{y,n} & \text{if } i_n \leq j < i_{n+1}. \end{cases}$$

If we put  $S = \{z_j : j \geq 1\}$ , then  $S$  converges to  $x_y$  in  $X$ , and  $f(S) = \{y_n\}$ . Therefore,  $f$  is 1-sequence-covering. □

*Remark 2.2.* By Theorem 2.1, we get an affirmative answer to Question 1.1.

**Corollary 2.3.** *Each sequence-covering quotient and boundary-compact map on  $g$ -metrizable spaces is weak-open.*

PROOF: Let  $f : X \rightarrow Y$  be a sequence-covering quotient and boundary-compact map and  $X$  be a  $g$ -metrizable space. By Theorem 2.1,  $f$  is 1-sequence-covering. Since  $f$  is quotient and  $X$  is sequential,  $f$  is weak-open by Corollary 3.5 in [18]. □

By Theorem 2.1 and Remark 1.7(1), the following corollaries hold.

**Corollary 2.4.** *Each sequence-covering and compact map on  $g$ -metrizable spaces is 1-sequence-covering.*

**Corollary 2.5** ([8, Theorem 2.1]). *Each sequence-covering and boundary-compact map on metric spaces is 1-sequence-covering.*

**Corollary 2.6** ([9, Theorem 4.5]). *Each closed sequence-covering map on  $g$ -metrizable spaces is 1-sequence-covering.*

PROOF: Let  $f : X \longrightarrow Y$  be a closed sequence-covering map and  $X$  be a  $g$ -metrizable space. By Lemma 3.1 in [15],  $Y$  is  $gf$ -countable. Furthermore, since  $Y$  is  $gf$ -countable and  $f$  is a closed map, it follows from Corollary 8 in [14] and Corollary 10 in [19] that  $Y$  contains no closed copy of  $S_\omega$ . By Lemma 3.2 in [15],  $f$  is a boundary-compact map. Therefore,  $f$  is 1-sequence-covering by Theorem 2.1.  $\square$

By Corollary 2.6, we have the following corollary.

**Corollary 2.7** ([11, Theorem 3.4.6]). *Each closed sequence-covering map on metric spaces is 1-sequence-covering.*

**Corollary 2.8.** *Each closed sequence-covering map on  $g$ -metrizable spaces is weak-open.*

PROOF: Let  $f : X \longrightarrow Y$  be a closed sequence-covering map and  $X$  be a  $g$ -metrizable space. It follows from Corollary 2.6 that  $f$  is 1-sequence-covering. Since  $f$  is closed and  $X$  is sequential,  $f$  is weak-open by Corollary 3.5 in [18].  $\square$

**Acknowledgment.** The author would like to express their thanks to referee for his/her helpful comments and valuable suggestions.

#### REFERENCES

- [1] An T.V., Tuyen L.Q., *Further properties of 1-sequence-covering maps*, Comment. Math. Univ. Carolin. **49** (2008), no. 3, 477–484.
- [2] An T.V., Tuyen L.Q., *On  $\pi$ -images of separable metric spaces and a problem of Shou Lin*, Mat. Vesnik, (2011), to appear.
- [3] Arhangel'skii A.V., *Mappings and spaces*, Russian Math. Surveys **21** (1966), no. 4, 115–162.
- [4] Engelking R., *General Topology (revised and completed edition)*, Heldermann Verlag, Berlin, 1989.
- [5] Franklin S.P., *Spaces in which sequences suffice*, Fund. Math. **57** (1965), 107–115.
- [6] Ge Y., *Characterizations of  $sn$ -metrizable spaces*, Publ. Inst. Math. (Beograd) (N.S) **74** (88) (2003), 121–128.
- [7] Lee K.B., *On certain  $g$ -first countable spaces*, Pacific J. Math. **65** (1976), no. 1, 113–118.
- [8] Lin F.C., Lin S., *On sequence-covering boundary compact maps of metric spaces*, Adv. Math. (China) **39** (2010), no. 1, 71–78.
- [9] Lin F.C., Lin S., *Sequence-covering maps on generalized metric spaces*, arXiv: 1106.3806.
- [10] Lin S., *On sequence-covering  $s$ -mappings*, Adv. Math. (China) **25** (1996), no. 6, 548–551.
- [11] Lin S., *Point-Countable Covers and Sequence-Covering Mappings*, Chinese Science Press, Beijing, 2002.

- [12] Lin S., Liu C., *On spaces with point-countable cs-networks*, Topology Appl. **74** (1996), 51–60.
- [13] Lin S., Yan P., *Sequence-covering maps of metric spaces*, Topology Appl. **109** (2001), 301–314.
- [14] Lin S., Tanaka Y., *Point-countable  $k$ -networks, closed maps, and related results*, Topology Appl. **59** (1994), 79–86.
- [15] Liu C., *On weak bases*, Topology Appl. **150** (2005), 91–99.
- [16] Siwiec F., *Sequence-covering and countably bi-quotient maps*, General Topology Appl. **1** (1971), 143–154.
- [17] Siwiec F., *On defining a space by a weak base*, Pacific J. Math. **52** (1974), 233–245.
- [18] Xia S., *Characterizations of certain  $g$ -first countable spaces*, Adv. Math. **29** (2000), 61–64.
- [19] Yan P., Lin S., *Point-countable  $k$ -networks,  $cs^*$ -network and  $\alpha_4$ -spaces*, Topology Proc. **24** (1999), 345–354.
- [20] Yan P., Lin S., *CWC-mappings and metrization theorems*, Adv. Math. (China) **36** (2007), no. 2, 153–158.
- [21] Yan P.F., Lin S., Jiang S.L., *Metrizability is preserved by closed sequence-covering maps*, Acta Math. Sinica **47** (2004), no. 1, 87–90.

DEPARTMENT OF MATHEMATICS, DA NANG UNIVERSITY, VIETNAM

*E-mail*: luongtuyench12@yahoo.com

(Received February 11, 2012)