

Xhevat Z. Krasniqi

Some new modified cosine sums and L^1 -convergence of cosine trigonometric series

Archivum Mathematicum, Vol. 49 (2013), No. 1, 43--50

Persistent URL: <http://dml.cz/dmlcz/143299>

Terms of use:

© Masaryk University, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**SOME NEW MODIFIED COSINE SUMS AND
 L^1 -CONVERGENCE OF COSINE TRIGONOMETRIC SERIES**

XHEVAT Z. KRASNIQI

ABSTRACT. In this paper we introduce some new modified cosine sums and then using these sums we study L^1 -convergence of trigonometric cosine series.

1. INTRODUCTION AND PRELIMINARIES

Let

$$(1.1) \quad \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx$$

be cosine trigonometric series and satisfy condition $a_k \rightarrow 0, k \rightarrow \infty$. The partial sum of series (1) we denote by $S_n(x) = \frac{a_0}{2} + \sum_{k=1}^n a_k \cos kx$ and let be $f(x) = \lim_{n \rightarrow \infty} S_n(x)$.

A sequence (a_k) is said to belong to the class S , or briefly $a_k \in S$, if $a_k \rightarrow 0$ as $k \rightarrow \infty$, and there exists a sequence of numbers (A_k) such that

$$A_k \downarrow 0, \\ \sum_{k=1}^{\infty} A_k < \infty,$$

and

$$|\Delta a_k| \leq A_k,$$

for all k , where $\Delta a_k = a_k - a_{k+1}$.

This class of sequences was defined by Sidon in [18] and by Telyakovskii in [21], therefore the class S is sometimes called the Sidon-Telyakovskii class. The class S is generalized later by Tomovski in [22] and by Leindler in [16].

Tomovski defined the class $S_r, r = 1, 2, \dots$ as follows: $\{a_k\}_{k=1}^{\infty} \in S_r$ if $a_k \rightarrow 0$ as $k \rightarrow \infty$ and there exists a monotonically decreasing sequence $\{A_k\}_{k=1}^{\infty}$ such that

2010 *Mathematics Subject Classification*: primary 42A20; secondary 42A32.

Key words and phrases: L^1 -convergence of trigonometric cosine series, new modified cosine sums, Dirichlet kernel.

Dedicated to my Professor Halil Turku on the occasion of his 79th birthday.

Received September 18, 2008, revised March 2013. Editor O. Došlý.

DOI: 10.5817/AM2013-1-43

$\sum_{k=1}^{\infty} k^r A_k < \infty$ and $|\Delta a_k| \leq A_k$ for all k . There was noticed that from $A_k \downarrow 0$ and $\sum_{k=1}^{\infty} k^r A_k < \infty$ it follows $k^{r+1} A_k = o(1), k \rightarrow \infty$. It is clear that $S_{r+1} \subset S_r$ for all $r = 1, 2, \dots$ and for $r = 0$ we get the class $S_0 \equiv S$.

Garret and Stanojević [3] have introduced modified cosine sums

$$f_n(x) = \frac{1}{2} \sum_{k=0}^n \Delta a_k + \sum_{k=1}^n \sum_{j=k}^n \Delta a_j \cos kx.$$

Garret and Stanojević [4], Ram [17], Singh and Sharma [20], and Kaur and Bhatia [11], [6], [10] studied the L^1 -convergence of this cosine sum under different sets of conditions on the coefficients a_n .

Kumari and Ram [15] introduced new modified cosine and sine sums as

$$h_n(x) = \frac{a_0}{2} + \sum_{k=1}^n \sum_{j=k}^n \Delta \left(\frac{a_j}{j} \right) k \cos kx,$$

$$g_n(x) = \sum_{k=1}^n \sum_{j=k}^n \Delta \left(\frac{a_j}{j} \right) k \sin kx$$

and have studied their L^1 -convergence under the condition that the coefficients a_n belong to different classes of sequences. They deduced some results about L^1 -convergence of cosine and sine series as corollaries, as well.

N. Hooda, B. Ram and S. S. Bhatia [5] introduced new modified cosine sums as

$$R_n(x) = \frac{1}{2} \left(a_1 + \sum_{k=0}^n \Delta^2 a_k \right) + \sum_{k=1}^n \left(a_{k+1} + \sum_{j=k}^n \Delta^2 a_j \right) \cos kx$$

and studied the L^1 -convergence of these cosine sums.

K. Kaur [9] introduced new modified sine sums as

$$K_n(x) = \frac{1}{2 \sin x} \sum_{k=1}^n \sum_{j=k}^n (\Delta a_{j-1} - \Delta a_{j+1}) \sin kx,$$

and studied the L^1 -convergence of this modified sine sum with semi-convex coefficients. Also, Kaur et al. [12] introduced a new class of numerical sequences as follows:

Definition 1. If $a_k = o(1)$ as $k \rightarrow \infty$, and

$$\sum_{k=1}^{\infty} k |\Delta^2 a_{k-1} - \Delta^2 a_{k+1}| < +\infty \quad (a_0 = 0)$$

then we say that $\{a_k\}$ belongs to the class **K**.

In their paper they proved the following result regarding to L^1 -convergence of the modified sums $K_n(x)$.

Theorem 1. Let the sequence $\{a_k\}$ belong to the class **K**, then $K_n(x)$ converges to $f(x)$ in the L^1 -norm.

Later on, Singh and Kaur [19] defined new modified generalized sine sums

$$K_{nr}(x) = \frac{1}{2 \sin x} \sum_{k=1}^n (\Delta^r a_{k-1} - \Delta^r a_{k+1}) \tilde{S}_k^{r-1}(x),$$

and a new class of sequences:

Definition 2. Let α be a positive real number. If $a_k = o(1)$ as $k \rightarrow \infty$, and

$$\sum_{k=1}^{\infty} k^\alpha |\Delta^{\alpha+1} a_{k-1} - \Delta^{\alpha+1} a_{k+1}| < +\infty \quad (a_0 = 0)$$

then we say that $\{a_k\}$ belongs to the class \mathbf{K}^α .

They proved the following generalization of Theorem 1.

Theorem 2. Let the sequence $\{a_k\}$ belong to the class \mathbf{K}^α , then $K_{nr}(x)$ converges to $f(x)$ in the L^1 -norm.

Some new modified sums are presented in [13] by present author (see also [14]) as follows

$$H_n(x) = \frac{1}{2 \sin x} \sum_{k=1}^n \sum_{j=k}^n \Delta [(a_{j-1} - a_{j+1}) \sin jx],$$

and also we have proved a new result as below.

Theorem 3. Let (a_n) be a semi-convex null sequence, then $H_n(x)$ converges to $f(x)$ in L^1 -norm.

The interested reader can find some new results in very recently published papers, [7] where the complex form of the sums $K_n(x)$ is introduced, and paper [8] in which it is studied the L^1 -convergence of sine trigonometric series by using a newly introduced modified cosine trigonometric sums under a new class of coefficient sequences (see [8] for details therein).

We recall that with regard to the L^1 -convergence of Ress-Stanojević cosine sums $f_n(x)$ to a cosine trigonometric series, belonging to the class S , Ram [17] proved the following theorem:

Theorem 4. If (1.1) belongs to the class S , then $\|f - f_n\|_{L^1} = o(1)$, $n \rightarrow \infty$.

In order to make an advanced study, on this treating topic, now we shall introduce new modified cosine sums as

$$G_n(x) = \frac{a_0}{2} + \sum_{k_1=1}^n \sum_{k_2=k_1}^n \sum_{k_3=k_2}^n \Delta^2 (a_{k_3} \cos k_3 x),$$

where $\Delta^2 a_k = \Delta(\Delta a_k) = a_k - 2a_{k+1} + a_{k+2}$.

Remark 1. The advantage of introducing of the above modified cosine sums is the following: We have verified that the sums $G_n(x)$ converge in L^1 -norm to $f(x)$, without a new class of null-sequences being defined, in contrary what the other authors previously did in their papers (as examples serve classes \mathbf{K} , \mathbf{K}^α , etc.).

The purpose of this paper is to prove analogous statement with Theorem 4 using new modified cosine sums $G_n(x)$ instead of $g_n(x)$ and the L^1 -convergence of the series (1.1) will be derived as a corollary.

As usual $D_n(x)$ will denote the real Dirichlet kernel, i.e.

$$D_n(x) = \frac{1}{2} + \sum_{k=1}^n \cos kx.$$

For the proof of main result we need the following lemma.

Lemma 1 ([2]). *If $|c_k| \leq 1$, then*

$$\int_0^\pi \left| \sum_{k=0}^n c_k \frac{\sin(k+1/2)x}{2 \sin \frac{x}{2}} \right| dx \leq C(n+1),$$

where C is a positive absolute constant.

2. MAIN RESULTS

We establish the following result.

Theorem 5. *Let (1.1) belong to the class S_2 , then $\|f - G_n\|_{L^1} = o(1)$, as $n \rightarrow \infty$.*

Proof. We have

$$\begin{aligned} G_n(x) &= \frac{a_0}{2} + \sum_{k_1=1}^n \sum_{k_2=k_1}^n \sum_{k_3=k_2}^n \Delta^2(a_{k_3} \cos k_3 x) \\ &= \frac{a_0}{2} + \sum_{k_1=1}^n \sum_{k_2=k_1}^n [\Delta(a_{k_2} \cos k_2 x) - \Delta(a_{k_2+1} \cos(k_2+1)x) \\ &\quad + \cdots + \Delta(a_n \cos nx) - \Delta(a_{n+1} \cos(n+1)x)] \\ &= \frac{a_0}{2} + \sum_{k_1=1}^n \sum_{k_2=k_1}^n [\Delta(a_{k_2} \cos k_2 x) - \Delta(a_{n+1} \cos(n+1)x)] \\ &= \frac{a_0}{2} + \sum_{k_1=1}^n [a_{k_1} \cos k_1 x - a_{k_1+1} \cos(k_1+1)x + \cdots + a_n \cos nx \\ &\quad - a_{n+1} \cos(n+1)x] - \Delta(a_{n+1} \cos(n+1)x) \sum_{k_1=1}^n (n-k_1+1) \\ &= S_n(x) - na_{n+1} \cos(n+1)x - \frac{1}{2}n(n+1)\Delta(a_{n+1} \cos(n+1)x) \\ &= S_n(x) - \frac{1}{2}n(n+3)a_{n+1} \cos(n+1)x \\ (2.1) \quad &+ \frac{1}{2}n(n+1)a_{n+2} \cos(n+2)x. \end{aligned}$$

From $A_k \downarrow 0$ and $\sum_{k=1}^{\infty} k^2 A_k < \infty$ follows $k^3 A_k = o(1)$, $k \rightarrow \infty$, which gives $k^2 A_k = o(1)$, $k \rightarrow \infty$. Therefore from

$$0 \leq n^2 |a_n| = n^2 \left| \sum_{k=n}^{\infty} \Delta a_k \right| \leq \left| \sum_{k=n}^{\infty} k^2 \Delta a_k \right| \leq \sum_{k=n}^{\infty} k^2 A_k = o(1), \quad n \rightarrow \infty$$

follow

$$(2.2) \quad n^2 a_n = o(1), \quad n a_n = o(1), \quad n \rightarrow \infty.$$

Also, $\cos(n+1)x$ and $\cos(n+2)x$ are finite in $[0, \pi]$ therefore from (2.1) and (2.2) we get

$$\lim_{n \rightarrow \infty} G_n(x) = \lim_{n \rightarrow \infty} S_n(x) = f(x).$$

On the other side, using Abel's transformation we have

$$\begin{aligned} f(x) - G_n(x) &= \lim_{m \rightarrow \infty} \left(\sum_{k=n+1}^{m-1} \Delta a_k D_k(x) + a_m D_m(x) - a_{n+1} D_n(x) \right) \\ &\quad + \frac{1}{2} n(n+3) a_{n+1} \cos(n+1)x - \frac{1}{2} n(n+1) a_{n+2} \cos(n+2)x \\ &= \sum_{k=n+1}^{\infty} \Delta a_k D_k(x) - a_{n+1} D_n(x) + \frac{1}{2} n(n+3) a_{n+1} \cos(n+1)x \\ &\quad - \frac{1}{2} n(n+1) a_{n+2} \cos(n+2)x. \end{aligned}$$

Therefore

$$\begin{aligned} \int_0^{\pi} |f(x) - G_n(x)| dx &\leq \int_0^{\pi} \left| \sum_{k=n+1}^{\infty} \Delta a_k D_k(x) \right| dx + |a_{n+1}| \int_0^{\pi} |D_n(x)| dx \\ &\quad + \frac{1}{2} n(n+3) |a_{n+1}| \int_0^{\pi} |\cos(n+1)x| dx \\ &\quad + \frac{1}{2} n(n+1) |a_{n+2}| \int_0^{\pi} |\cos(n+2)x| dx \\ (2.3) \quad &:= \sum_{\nu=1}^4 B_{\nu}(n). \end{aligned}$$

Since $a_k \in S_2 \subset S_0 \equiv S$ then $\sum_{k=n+1}^{\infty} (k+1) \Delta A_k = o(1)$ as $n \rightarrow \infty$, therefore from this fact, Lemma 1, and using Abel's transformation we have

$$\begin{aligned} B_1(n) &= \int_0^{\pi} \left| \sum_{k=n+1}^{\infty} A_k \frac{\Delta a_k}{A_k} D_k(x) \right| dx \leq \sum_{k=n+1}^{\infty} \Delta A_k \int_0^{\pi} \left| \sum_{i=0}^k \frac{\Delta a_i}{A_i} D_i(x) \right| dx \\ (2.4) \quad &= O\left(\sum_{k=n+1}^{\infty} (k+1) \Delta A_k \right) = o(1), \quad n \rightarrow \infty. \end{aligned}$$

By well-known Zygmund's theorem (see [20, p. 458]), for n sufficiently large, the following relation holds

$$\int_0^\pi |D_n(x)| dx \sim \log n,$$

therefore from the last relation and (2.2) we have

$$(2.5) \quad B_2(n) = |a_{n+1}| \log n \leq n|a_{n+1}| = o(1), \quad n \rightarrow \infty.$$

Moreover, from fact that integrals $\int_0^\pi |\cos(n+1)x| dx$, $\int_0^\pi |\cos(n+2)x| dx$ are bounded, and from relation (2.2) we conclude that

$$(2.6) \quad B_3(n) = O(n(n+3)|a_{n+1}|) = o(1), \quad n \rightarrow \infty$$

and similarly

$$(2.7) \quad B_4(n) = O(n(n+1)|a_{n+2}|) = o(1), \quad n \rightarrow \infty.$$

Finally, from (2.3)–(2.7) it follows that

$$\|f - G_n\|_{L^1} = o(1), \quad n \rightarrow \infty.$$

The proof of the Theorem 5 is completed. \square

Corollary 1. *Let (1.1) belong to the class S_2 , then $\|f - S_n\|_{L^1} = o(1)$ as $n \rightarrow \infty$.*

Proof. From Theorem 5, and relations (2.6), (2.7), we have

$$\begin{aligned} \|f - S_n\|_{L^1} &= \|f - G_n + G_n - S_n\|_{L^1} \\ &\leq \|f - G_n\|_{L^1} + \|G_n - S_n\|_{L^1} \\ &\leq \|f - G_n\|_{L^1} + \frac{1}{2}n(n+3)|a_{n+1}| \int_0^\pi |\cos(n+1)x| dx \\ &\quad + \frac{1}{2}n(n+1)|a_{n+2}| \int_0^\pi |\cos(n+2)x| dx = o(1) \end{aligned}$$

as $n \rightarrow \infty$, which completely proves the corollary. \square

Remark 2. A closer examination of the proofs of Theorem 5 and Corollary 1 reveals that condition $a_k \in S_2$ can be replaced by conditions $a_k \in S$ and $n^2|a_n| = o(1)$. This enables us to formulate Theorem 5 and Corollary 1 in the following form:

Theorem 6. *Let (a_k) belong to the class S and $n^2|a_n| = o(1)$, then $\|f - G_n\|_{L^1} = o(1)$ as $n \rightarrow \infty$.*

Corollary 2. *Let (a_k) belong to the class S and $n^2|a_n| = o(1)$, then $\|f - S_n\|_{L^1} = o(1)$ as $n \rightarrow \infty$.*

We would like to finalize this paper with a comment. We have noticed during this study that, if someone tries to introduce some modified sums of the form

$$T_{n,m}(x) = \frac{a_0}{2} + \sum_{k_1=1}^n \sum_{k_2=k_1}^n \sum_{k_3=k_2}^n \dots \sum_{k_m=k_{m-1}}^n \Delta^{m-1}(a_{k_m}/k_m) k_1 \cos k_1 x,$$

where $m \in \mathbb{N}$, $m > 3$, $\Delta a_k = a_k - a_{k+1}$, $\Delta^{m-1} a_k = \Delta (\Delta^{m-2} a_k)$, which is a natural extension of our results, then several difficulties in the proof of the counterpart of Theorem 5 will be appeared. This is why we are focused only on the case $m = 3$.

REFERENCES

- [1] Bary, N. K., *A treatise on trigonometric series*, Pergamon Press, Oxford–London–New York–Paris–Frankfurt, 1964.
- [2] Fomin, G. A., *On linear methods for summing Fourier series*, Math. Sbornik **66** (1964), 144–152.
- [3] Garret, W. J., Stanojević, Č. V., *On integrability and L^1 -convergence of certain cosine sums*, Notices, Amer. Math. Soc. **22** (1975), A–166.
- [4] Garret, W. J., Stanojević, Č. V., *On L^1 -convergence of certain cosine sums*, Proc. Amer. Math. Soc. **54** (1976), 101–105.
- [5] Hooda, N., Ram, B., Bhatia, S. S., *On L^1 -convergence of a modified cosine sum*, Soochow J. of Math. **28** (3) (2002), 305–310.
- [6] Kaur, J., Bhatia, S. S., *Integrability and L^1 -convergence of certain cosine sums*, Kyungpook Math. J. **47** (2007), 323–328.
- [7] Kaur, J., Bhatia, S. S., *Convergence of modified complex trigonometric sums in the metric space L* , Lobachevskii J. Math. **31** (3) (2010), 290–294.
- [8] Kaur, J., Bhatia, S. S., *A class of L^1 -convergence of new modified cosine sums*, Southeast Asian Bull. Math. **36** (2012), 831–836.
- [9] Kaur, K., *On L^1 -convergence of a modified sine sums*, Electronic J. Geogr. Math. **14** (1) (2003).
- [10] Kaur, K., *Integrability and L^{sp1} -convergence of Rees–Stanojević sums with generalized semi-convex coefficients of non-integral orders*, Arch. Math. (Brno) **41** (4) (2005), 423–437.
- [11] Kaur, K., Bhatia, S. S., *Integrability and L^1 -convergence of Rees–Stanojević sums with generalized semi-convex coefficients*, Int. J. Math. Math. Sci. **30** (11) (2002), 645–650.
- [12] Kaur, K., Bhatia, S. S., Ram, B., *Integrability and L^1 -convergence of modified sine sums*, Georgian Math. J. **11** (1) (2004), 99–104.
- [13] Krasniqi, Xh. Z., *A note on L^1 -convergence of the sine and cosine trigonometric series with semi-convex coefficients*, Int. J. Open Probl. Comput. Sci. Math. **2** (2) (2009), 231–239.
- [14] Krasniqi, Xh. Z., *On L^1 -convergence of Rees–Stanojević’s sums with coefficients from the class \mathbf{K}* , Matematiche **65** (2) (2010), 25–32, DOI: 10.4418/2010.65.2.3.
- [15] Kumari, S., Ram, B., *L^1 -convergence of certain trigonometric sums*, Indian J. Pure Appl. Math. **20** (9) (1989), 908–914.
- [16] Leindler, L., *Additions to the Telyakovskii class S* , J. Inequal. Pure Appl. Math. **4** (2) (2003), 5, electronic only.
- [17] Ram, B., *Convergence of certain cosine sums in the metric space L* , Proc. Amer. Math. Soc. **66** (1977), 258–260.
- [18] Sidon, S., *Hinreichende Bedingungen für den Fourier-Charakter einer trigonometrischen Reihe*, J. London Math. Soc. **14** (1939), 1528–160.

- [19] Singh, K., Kaur, K., *On L^1 -convergence of certain generalized modified trigonometric sums*, Mat. Vesnik **61** (3) (2009), 219–226.
- [20] Singh, N., Sharma, K. M., *Convergence of certain cosine sums in the metric space L* , Proc. Amer. Math. Soc. **75** (1978), 117–120.
- [21] Telyakovskii, S. A., *On a sufficient condition of Sidon for integrability of trigonometric series*, Math. Zametki **14** (1973), 317–328.
- [22] Tomovski, Ž., *Some results on L^1 -approximation of the r -th derivative of Fourier series*, J. Inequal. Pure Appl. Math. **1** (2002), 11, electronic only.

UNIVERSITY OF PRISHTINA, FACULTY OF EDUCATION,
DEPARTMENT OF MATHEMATICS AND INFORMATICS,
AGIM RAMADANI ST., N.N., PRISHTINË 10000, REPUBLIC OF KOSOVA
E-mail: xhevat.krasniqi@uni-pr.edu