

Khalida Aissani; Mouffak Benchohra

Semilinear fractional order integro-differential equations with infinite delay in Banach spaces

*Archivum Mathematicum*, Vol. 49 (2013), No. 2, 105--117

Persistent URL: <http://dml.cz/dmlcz/143498>

## Terms of use:

© Masaryk University, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

# SEMILINEAR FRACTIONAL ORDER INTEGRO-DIFFERENTIAL EQUATIONS WITH INFINITE DELAY IN BANACH SPACES

KHALIDA AISSANI AND MOUFFAK BENCHOHRA

**ABSTRACT.** This paper concerns the existence of mild solutions for fractional order integro-differential equations with infinite delay. Our analysis is based on the technique of Kuratowski's measure of noncompactness and Mönch's fixed point theorem. An example to illustrate the applications of main results is given.

## 1. INTRODUCTION

Fractional calculus is a generalization of the ordinary differentiation and integration to arbitrary non-integer order. The subject is as old as the differential calculus since, starting from some speculations of G. W. Leibniz (1697) and L. Euler (1730), it has been progressing up to nowadays. Fractional differential and integral equations have recently been applied in various areas of engineering, science, finance, applied mathematics, bio-engineering, radiative transfer, neutron transport and the kinetic theory of gases and others [5, 10, 11]. There has been a significant development in ordinary and partial fractional differential equations in recent years; see the monographs of Abbas et al. [1], Baleanu et al. [6], Diethelm [13], Hilfer [18], Kilbas et al. [20], Miller and Ross [26], Podlubny [31], Samko et al. [32], and Tarasov [33], and the papers [2, 3, 8, 14, 23, 24, 28, 29, 30].

The theory of functional differential equations has emerged as an important branch of nonlinear analysis. Differential delay equations, or functional differential equations, have been used in modeling scientific phenomena for many years. Often, it has been assumed that the delay is either a fixed constant or is given as an integral in which case it is called a distributed delay; see for instance the books by Hale and Verduyn Lunel [16], Hino et al. [19], Kolmanovskii and Myshkis [21], Lakshmikantham et al. [22], and Wu [34], and the papers [12, 15].

---

2010 *Mathematics Subject Classification*: primary 26A33; secondary 34G20, 34K30, 34K37.

*Key words and phrases*: semilinear differential equations, Caputo fractional derivative, mild solution, measure of noncompactness, fixed point, semigroup, Banach space.

Received February 13, 2013, revised May 2013. Editor R. Šimon Hilscher.

DOI: 10.5817/AM2013-2-105

In this work we discuss the existence of mild solutions for fractional order integro-differential equations with infinite delay of the form

$$(1) \quad \begin{aligned} D_t^q x(t) &= Ax(t) + \int_0^t a(t,s)f(s, x_s, x(s))ds, \quad t \in J = [0, T], \\ x(t) &= \phi(t), \quad t \in (-\infty, 0], \end{aligned}$$

where  $D_t^q$  is the Caputo fractional derivative of order  $0 < q < 1$ ,  $A$  is a generator of an analytic semigroup  $\{S(t)\}_{t \geq 0}$  of uniformly bounded linear operators on  $X$ ,  $f: J \times \mathcal{B} \times X \rightarrow X$ ,  $a: D \rightarrow \mathbb{R}$  ( $D = \{(t, s) \in [0, T] \times [0, T] : t \geq s\}$ ),  $\phi \in \mathcal{B}$  where  $\mathcal{B}$  is called phase space to be defined in Section 2. For any function  $x$  defined on  $(-\infty, T]$  and any  $t \in J$ , we denote by  $x_t$  the element of  $\mathcal{B}$  defined by

$$x_t(\theta) = x(t + \theta), \quad \theta \in (-\infty, 0].$$

Here  $x_t$  represents the history of the state up to the present time  $t$ .

In the present paper we deal with an infinite time delay. Note that in this case, the phase space  $\mathcal{B}$  plays a crucial role in the study of both qualitative and quantitative aspects of theory of functional equations (see [15]).

We present an existence result of mild solutions for the problem (1) by means of the application of Mönch's fixed point theorem combined with the Kuratowski measure of noncompactness. An example illustrating the abstract theory will be presented.

## 2. PRELIMINARIES

Let  $(X, \|\cdot\|)$  be a real Banach space.

$C = C(J, X)$  be the space of all  $X$ -valued continuous functions on  $J$ .

$L(X)$  be the Banach space of all linear and bounded operators on  $X$ .

$L^1(J, X)$  the space of  $X$ -valued Bochner integrable functions on  $J$  with the norm

$$\|y\|_{L^1} = \int_0^T \|y(t)\| dt.$$

$L^\infty(J, \mathbb{R})$  is the Banach space of essentially bounded functions, normed by

$$\|y\|_{L^\infty} = \inf\{d > 0 : |y(t)| \leq d, \text{ a.e. } t \in J\}.$$

**Definition 2.1.** A function  $f: J \times \mathcal{B} \times X \rightarrow X$  is said to be an Carathéodory function if it satisfies:

- (i) for each  $t \in J$  the function  $f(t, \cdot, \cdot): \mathcal{B} \times X \rightarrow X$  is continuous;
- (ii) for each  $(v, w) \in \mathcal{B} \times X$  the function  $f(\cdot, v, w): J \rightarrow X$  is measurable.

Next we give the concept of a measure of noncompactness [7].

**Definition 2.2.** Let  $B$  be a bounded subset of a seminormed linear space  $Y$ . Kuratowski's measure of noncompactness of  $B$  is defined as

$$\alpha(B) = \inf\{d > 0 : B \text{ has a finite cover by sets of diameter } \leq d\}.$$

We note that this measure of noncompactness satisfies interesting regularity properties (for more information, we refer to [7]).

**Lemma 2.3.**

- (1) If  $A \subseteq B$  then  $\alpha(A) \leq \alpha(B)$ ,
- (2)  $\alpha(A) = \alpha(\bar{A})$ , where  $\bar{A}$  denotes the closure of  $A$ ,
- (3)  $\alpha(A) = 0 \Leftrightarrow \bar{A}$  is compact ( $A$  is relatively compact),
- (4)  $\alpha(\lambda A) = |\lambda|\alpha(A)$ , with  $\lambda \in \mathbb{R}$ ,
- (5)  $\alpha(A \cup B) = \max\{\alpha(A), \alpha(B)\}$ ,
- (6)  $\alpha(A + B) \leq \alpha(A) + \alpha(B)$ , where
 
$$A + B = \{x + y : x \in A, y \in B\},$$
- (7)  $\alpha(A + a) = \alpha(A)$  for any  $a \in Y$ ,
- (8)  $\alpha(\overline{\text{conv}}A) = \alpha(A)$ , where  $\overline{\text{conv}}A$  is the closed convex hull of  $A$ .

For  $H \subset C(J, X)$ , we define

$$(2) \quad \int_0^t H(s)ds = \left\{ \int_0^t u(s)ds : u \in H \right\} \quad \text{for } t \in J,$$

where  $H(s) = \{u(s) \in X : u \in H\}$ .

**Lemma 2.4** ([7]). *If  $H \subset C(J, X)$  is a bounded, equicontinuous set, then*

$$(3) \quad \alpha(H) = \sup_{t \in J} \alpha(H(t)).$$

**Lemma 2.5** ([17]). *If  $\{u_n\}_{n=1}^\infty \subset L^1(J, X)$  and there exists  $m \in L^1(J, \mathbb{R}^+)$  such that  $\|u_n(t)\| \leq m(t)$ , a.e.  $t \in J$ , then  $\alpha(\{u_n(t)\}_{n=1}^\infty)$  is integrable and*

$$(4) \quad \alpha\left(\left\{ \int_0^t u_n(s) ds \right\}_{n=1}^\infty\right) \leq 2 \int_0^t \alpha(\{u_n(s)\}_{n=1}^\infty) ds.$$

In this paper, we will employ an axiomatic definition for the phase space  $\mathcal{B}$  which is similar to those introduced by Hale and Kato [15]. Specifically,  $\mathcal{B}$  will be a linear space of functions mapping  $(-\infty, 0]$  into  $X$  endowed with a seminorm  $\|\cdot\|_{\mathcal{B}}$ , and satisfies the following axioms:

**(A1):** If  $x : (-\infty, T] \rightarrow X$  is continuous on  $J$  and  $x_0 \in \mathcal{B}$ , then  $x_t \in \mathcal{B}$  and  $x_t$  is continuous in  $t \in J$  and

$$(5) \quad \|x(t)\| \leq C\|x_t\|_{\mathcal{B}},$$

where  $C \geq 0$  is a constant.

**(A2):** There exist a continuous function  $C_1(t) > 0$  and a locally bounded function  $C_2(t) \geq 0$  in  $t \geq 0$  such that

$$(6) \quad \|x_t\|_{\mathcal{B}} \leq C_1(t) \sup_{s \in [0, t]} \|x(s)\| + C_2(t)\|x_0\|_{\mathcal{B}},$$

for  $t \in [0, T]$  and  $x$  as in (A1).

**(A3):** The space  $\mathcal{B}$  is complete.

**Remark 2.6.** Condition (5) in (A1) is equivalent to  $\|\phi(0)\| \leq C\|\phi\|_{\mathcal{B}}$ , for all  $\phi \in \mathcal{B}$ .

For our purpose we will only need the following fixed point theorem.

**Theorem 2.7** ([4, 27]). *Let  $U$  be a bounded, closed and convex subset of a Banach space such that  $0 \in U$ , and let  $N$  be a continuous mapping of  $U$  into itself. If the implication*

$$V = \overline{\text{conv}}N(V) \text{ or } V = N(V) \cup \{0\} \implies \alpha(V) = 0$$

*holds for every subset  $V$  of  $U$ , then  $N$  has a fixed point.*

Let  $\Omega$  be a set defined by

$$\Omega = \{x : (-\infty, T] \rightarrow X \text{ such that } x|_{(-\infty, 0]} \in \mathcal{B}, x|_J \in C(J, X)\} .$$

### 3. EXISTENCE OF MILD SOLUTIONS

Following [25, 14] we will introduce now the definition of mild solution to (1).

**Definition 3.1.** A function  $x \in \Omega$  is said to be a mild solution of (1) if  $x$  satisfies

$$(7) \quad x(t) = \begin{cases} \phi(t), & t \in (-\infty, 0]; \\ -Q(t)\phi(0) + \int_0^t \int_0^s R(t-s)a(s,\tau)f(\tau, x_\tau, x(\tau)) d\tau ds, & t \in J, \end{cases}$$

where

$$Q(t) = \int_0^\infty \xi_q(\sigma)S(t^q\sigma)d\sigma, \quad R(t) = q \int_0^\infty \sigma t^{q-1}\xi_q(\sigma)S(t^q\sigma)d\sigma$$

and  $\xi_q$  is a probability density function defined on  $(0, \infty)$  such that

$$\xi_q(\sigma) = \frac{1}{q}\sigma^{-1-\frac{1}{q}}\varpi_q(\sigma^{-\frac{1}{q}}) \geq 0,$$

where

$$\varpi_q(\sigma) = \frac{1}{\pi} \sum_{n=1}^\infty (-1)^{n-1} \sigma^{-qn-1} \frac{\Gamma(nq+1)}{n!} \sin(n\pi q), \quad \sigma \in (0, \infty).$$

**Remark 3.2.** Note that  $\{S(t)\}_{t \geq 0}$  is a uniformly bounded semigroup, i.e,

there exists a constant  $M > 0$  such that  $\|S(t)\| \leq M$  for all  $t \in [0, T]$ .

**Remark 3.3.** According to [25], a direct calculation gives that

$$(8) \quad \|R(t)\| \leq C_{q,M}t^{q-1}, \quad t > 0,$$

where  $C_{q,M} = \frac{qM}{\Gamma(1+q)}$ .

We make the following assumptions.

(H1)  $f: J \times \mathcal{B} \times X \rightarrow X$  satisfies the Carathéodory conditions, and there exist two positive functions  $\mu_i(\cdot) \in L^1(J, \mathbb{R}^+)$  ( $i = 1, 2$ ) with

$$(9) \quad \|\mu_2\|_{L^1(J, \mathbb{R}^+)} < \frac{q}{T^q a C_{q,M}},$$

such that

$$(10) \quad \|f(t, v, w)\| \leq \mu_1(t)\|v\|_{\mathcal{B}} + \mu_2(t)\|w\|, \quad (t, v, w) \in J \times \mathcal{B} \times X.$$

(H2) For any bounded sets  $D_1 \subset \mathcal{B}, D_2 \subset X$ , and  $0 \leq s \leq t \leq T$ , there exists an integrable positive function  $\eta$  such that

$$\alpha(R(t-s)f(\tau, D_1, D_2)) \leq \eta_t(s, \tau) \left( \alpha(D_2) + \sup_{-\infty < \theta \leq 0} \alpha(D_1(\theta)) \right),$$

where  $\eta_t(s, \tau) = \eta(t, s, \tau)$  and  $\sup_{t \in J} \int_0^t \int_0^s \eta_t(s, \tau) d\tau ds = \eta^* < \infty$ .

(H3) For each  $t \in J$ ,  $a(t, s)$  is measurable on  $[0, t]$  and  $a(t) = \text{ess sup}\{ |a(t, s)|, 0 \leq s \leq t \}$  is bounded on  $J$ . The map  $t \rightarrow a_t$  is continuous from  $J$  to  $L^\infty(J, \mathbb{R})$ , here,  $a_t(s) = a(t, s)$ .

Set  $a = \sup_{t \in J} a(t)$ .

**Theorem 3.4.** *Suppose that the assumptions (H1)–(H3) hold with*

$$(11) \quad 16a\eta^* < 1,$$

*then the problem (1) has at least one mild solution on  $(-\infty, T]$ .*

**Proof.** We transform the problem (1) into a fixed-point problem. Define a mapping  $\Phi$  from  $\Omega$  into itself by

$$\Phi(x)(t) = \begin{cases} \phi(t), & t \in (-\infty, 0]; \\ -Q(t)\phi(0) + \int_0^t \int_0^s R(t-s)a(s, \tau)f(\tau, x_\tau, x(\tau)) d\tau ds, & t \in J. \end{cases}$$

Clearly, fixed points of the operator  $\Phi$  are mild solutions of the problem (1).

For  $\phi \in \mathcal{B}$ , we will define the function  $y(\cdot): (-\infty, T] \rightarrow X$  by

$$y(t) = \begin{cases} \phi(t), & \text{if } t \in (-\infty, 0]; \\ 0, & \text{if } t \in J. \end{cases}$$

Then  $y_0 = \phi$ . For each  $z \in C(J, X)$  with  $z(0) = 0$ , we denote by  $\bar{z}$  the function defined by

$$\bar{z}(t) = \begin{cases} 0, & \text{if } t \in (-\infty, 0]; \\ z(t), & \text{if } t \in J. \end{cases}$$

If  $x(\cdot)$  verifies (7), we can decompose it as  $x(t) = y(t) + \bar{z}(t)$ , for  $t \in J$ , which implies  $x_t = y_t + \bar{z}_t$ , for every  $t \in J$  and the function  $z(t)$  satisfies

$$z(t) = -Q(t)\phi(0) + \int_0^t \int_0^s R(t-s)a(s,\tau)f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau)) d\tau ds.$$

Let

$$Z_0 = \{z \in \Omega : z_0 = 0\}.$$

For any  $z \in Z_0$ , we have

$$\|z\|_{Z_0} = \sup_{t \in J} \|z(t)\| + \|z_0\|_{\mathcal{B}} = \sup_{t \in J} \|z(t)\|.$$

Thus  $(Z_0, \|\cdot\|_{Z_0})$  is a Banach space. We define the operator  $\tilde{\Phi}: Z_0 \rightarrow Z_0$  by:

$$\tilde{\Phi}(z)(t) = -Q(t)\phi(0) + \int_0^t \int_0^s R(t-s)a(s,\tau)f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau)) d\tau ds.$$

Obviously, the operator  $\Phi$  has a fixed point is equivalent to  $\tilde{\Phi}$  has one, so it turns to prove that  $\tilde{\Phi}$  has a fixed point. Let  $r > 0$  and consider the set

$$B_r = \{z \in Z_0 : \|z\|_{Z_0} \leq r\}.$$

□

We need the following lemma.

**Lemma 3.5.** *Set*

$$(12) \quad C_1^* = \sup_{t \in J} C_1(t); \quad C_2^* = \sup_{\eta \in J} C_2(\eta).$$

Then for any  $z \in B_r$  we have

$$\|y_t + \bar{z}_t\|_{\mathcal{B}} \leq C_2^* \|\phi\|_{\mathcal{B}} + C_1^* r := r^*,$$

and

$$(13) \quad \|f(t, y_t + \bar{z}_t, y(t) + \bar{z}(t))\| \leq \mu_1(t)r^* + \mu_2(t)r.$$

**Proof.** Using (6), (12) and (10), we obtain

$$\begin{aligned} \|y_t + \bar{z}_t\|_{\mathcal{B}} &\leq \|y_t\|_{\mathcal{B}} + \|\bar{z}_t\|_{\mathcal{B}} \\ &\leq C_1(t) \sup_{0 \leq \tau \leq t} \|y(\tau)\| + C_2(t)\|y_0\|_{\mathcal{B}} + C_1(t) \sup_{0 \leq \tau \leq t} \|z(\tau)\| + C_2(t)\|z_0\|_{\mathcal{B}} \\ &\leq C_2(t)\|\phi\|_{\mathcal{B}} + C_1(t) \sup_{0 \leq \tau \leq t} \|z(\tau)\| \\ &\leq C_2^* \|\phi\|_{\mathcal{B}} + C_1^* r := r^*. \end{aligned}$$

Also, we get

$$\begin{aligned} \|f(t, y_t + \bar{z}_t, y(t) + \bar{z}(t))\| &\leq \mu_1(t)\|y_t + \bar{z}_t\|_{\mathcal{B}} + \mu_2(t)\|y(t) + \bar{z}(t)\| \\ &\leq \mu_1(t)r^* + \mu_2(t)r. \end{aligned}$$

The lemma is proved. □

Now we prove that  $\tilde{\Phi}$  has a fixed point. The proof will be given in three steps.

**Step 1:**  $\tilde{\Phi}$  is continuous.

Let  $\{z^k\}_{k \in \mathbb{N}}$  be a sequence such that  $z^k \rightarrow z$  in  $B_r$  as  $k \rightarrow \infty$ . Then for each  $t \in J$ , we have

$$\begin{aligned} \|\tilde{\Phi}(z^k)(t) - \tilde{\Phi}(z)(t)\| &\leq \int_0^t \int_0^s \|R(t-s)a(s, \tau)[f(\tau, y_\tau + \bar{z}_\tau^k, y(\tau) + \bar{z}^k(\tau)) \\ &\quad - f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau))]\| d\tau ds \\ &\leq a C_{q,M} \int_0^t \int_0^s (t-s)^{q-1} \|f(\tau, y_\tau + \bar{z}_\tau^k, y(\tau) + \bar{z}^k(\tau)) \\ &\quad - f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau))\| d\tau ds. \end{aligned}$$

Since  $f$  is of Carathéodory type, we have by the Lebesgue Dominated Convergence Theorem that

$$\|\tilde{\Phi}(z^k)(t) - \tilde{\Phi}(z)(t)\| \rightarrow 0 \quad \text{when } k \rightarrow \infty.$$

Consequently,

$$\lim_{k \rightarrow \infty} \|\tilde{\Phi}(z^k) - \tilde{\Phi}(z)\|_{Z_0} = 0.$$

Thus  $\tilde{\Phi}$  is continuous.

**Step 2:**  $\tilde{\Phi}$  maps  $B_r$  into itself. Let

$$r \geq \frac{M\|\phi\|_{\mathcal{B}} + \frac{T^q a C_{q,M} r^* \|\mu_1\|_{L^1(J, \mathbb{R}^+)}}{q}}{1 - \frac{T^q a C_{q,M} \|\mu_2\|_{L^1(J, \mathbb{R}^+)}}{q}}.$$

Then for each  $z \in B_r$  and  $t \in J$  we have

$$\begin{aligned} \|\tilde{\Phi}(z)(t)\| &\leq \|Q(t)\phi(0)\| \\ &\quad + \int_0^t \int_0^s \|R(t-s)a(s, \tau)f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau))\| d\tau ds \\ &\leq M\|\phi\|_{\mathcal{B}} + a C_{q,M} \int_0^t \int_0^s (t-s)^{q-1} [\mu_1(\tau)r^* + \mu_2(\tau)r] d\tau ds \\ &\leq M\|\phi\|_{\mathcal{B}} + a C_{q,M} r^* \int_0^t \int_0^s (t-s)^{q-1} \mu_1(\tau) d\tau ds \\ &\quad + a C_{q,M} r \int_0^t \int_0^s (t-s)^{q-1} \mu_2(\tau) d\tau ds \\ &\leq M\|\phi\|_{\mathcal{B}} + \frac{T^q a C_{q,M}}{q} [r^* \|\mu_1\|_{L^1(J, \mathbb{R}^+)} + r \|\mu_2\|_{L^1(J, \mathbb{R}^+)}] \\ &\leq r. \end{aligned}$$

**Step 3:**  $\tilde{\Phi}(B_r)$  is bounded and equicontinuous.



By Step 2, it is obvious that  $\tilde{\Phi}(B_r) \subset B_r$  is bounded. For the equicontinuity of  $\tilde{\Phi}(B_r)$ . Let  $\tau_1, \tau_2 \in J$  with  $\tau_1 > \tau_2$ , and let  $z \in B_r$ . Then

$$\begin{aligned} \|\tilde{\Phi}(z)(\tau_1) - \tilde{\Phi}(z)(\tau_2)\| &\leq \|Q(\tau_1) - Q(\tau_2)\| \|\phi(0)\| \\ &+ \left\| \int_0^{\tau_1} \int_0^s R(\tau_1 - s) a(s, \tau) f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau)) d\tau ds \right. \\ &\quad \left. - \int_0^{\tau_2} \int_0^s R(\tau_2 - s) a(s, \tau) f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau)) d\tau ds \right\|. \end{aligned}$$

Set

$$G(\cdot, y + \bar{z}, y(\cdot) + \bar{z}(\cdot)) = \int_0^\cdot a(\cdot, \tau) f(\tau, y_\tau + \bar{z}_\tau, y(\tau) + \bar{z}(\tau)) d\tau,$$

then

$$\begin{aligned} \|\tilde{\Phi}(z)(\tau_1) - \tilde{\Phi}(z)(\tau_2)\| &\leq \|Q(\tau_1) - Q(\tau_2)\| \|\phi(0)\| \\ &+ \left\| \int_0^{\tau_1} R(\tau_1 - s) G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right. \\ &\quad \left. - \int_0^{\tau_2} R(\tau_2 - s) G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right\| \\ &\leq \|Q(\tau_1) - Q(\tau_2)\| \|\phi(0)\| \\ &+ \left\| \int_0^{\tau_2} R(\tau_1 - s) G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right. \\ &\quad \left. + \int_{\tau_2}^{\tau_1} R(\tau_1 - s) G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right. \\ &\quad \left. - \int_0^{\tau_2} R(\tau_2 - s) G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right\| \\ &\leq \|Q(\tau_1) - Q(\tau_2)\| \|\phi(0)\| \\ &+ \left\| \int_0^{\tau_2} [R(\tau_1 - s) - R(\tau_2 - s)] G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right\| \\ &\quad + \int_{\tau_2}^{\tau_1} \|R(\tau_1 - s)\| \|G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s))\| ds \\ &\leq I_1 + I_2 + I_3, \end{aligned}$$

where

$$I_1 = \|Q(\tau_1) - Q(\tau_2)\| \|\phi(0)\|$$

$$I_2 = \left\| \int_0^{\tau_2} [R(\tau_1 - s) - R(\tau_2 - s)] G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right\|$$

$$I_3 = \int_{\tau_2}^{\tau_1} \|R(\tau_1 - s)\| \|G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s))\| ds.$$

$I_1$  tends to zero as  $\tau_2 \rightarrow \tau_1$ , since  $Q(t)$  is a strongly continuous operator. For  $I_2$ , using (8) and (13), we have

$$\begin{aligned}
 I_2 &\leq \left\| \int_0^{\tau_2} [q \int_0^\infty \sigma(\tau_1 - s)^{q-1} \xi_q(\sigma) S((\tau_1 - s)^q \sigma) d\sigma \right. \\
 &\quad \left. - q \int_0^\infty \sigma(\tau_2 - s)^{q-1} \xi_q(\sigma) S((\tau_2 - s)^q \sigma) d\sigma] G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) ds \right\| \\
 &\leq q \int_0^{\tau_2} \int_0^\infty \sigma \left| [(\tau_1 - s)^{q-1} - (\tau_2 - s)^{q-1}] \xi_q(\sigma) S((\tau_1 - s)^q \sigma) \right. \\
 &\quad \times G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s)) \left. \right\| d\sigma ds \\
 &\quad + q \int_0^{\tau_2} \int_0^\infty \sigma(\tau_2 - s)^{q-1} \xi_q(\sigma) \|S((\tau_1 - s)^q \sigma) - S((\tau_2 - s)^q \sigma)\| \\
 &\quad \times \|G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s))\| d\sigma ds \\
 &\leq C_{q,M} \int_0^{\tau_2} |(\tau_1 - s)^{q-1} - (\tau_2 - s)^{q-1}| \|G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s))\| ds \\
 &\quad + q \int_0^{\tau_2} \int_0^\infty \sigma(\tau_2 - s)^{q-1} \xi_q(\sigma) \|S((\tau_1 - s)^q \sigma) - S((\tau_2 - s)^q \sigma)\| \\
 &\quad \times \|G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s))\| d\sigma ds \\
 &\leq a [r^* \|\mu_1\|_{L^1(J, \mathbb{R}^+)} + r \|\mu_2\|_{L^1(J, \mathbb{R}^+)}] \\
 &\quad \times [C_{q,M} \int_0^{\tau_2} |(\tau_1 - s)^{q-1} - (\tau_2 - s)^{q-1}| ds \\
 &\quad + q \int_0^{\tau_2} \int_0^\infty \sigma(\tau_2 - s)^{q-1} \xi_q(\sigma) \|S((\tau_1 - s)^q \sigma) - S((\tau_2 - s)^q \sigma)\| d\sigma ds].
 \end{aligned}$$

Clearly, the first term on the right-hand side of the above inequality tends to zero as  $\tau_2 \rightarrow \tau_1$ . From the continuity of  $S(t)$  in the uniform operator topology for  $t > 0$ , the second term on the right-hand side of the above inequality tends to zero as  $\tau_2 \rightarrow \tau_1$ . In view of (13), we have

$$\begin{aligned}
 I_3 &\leq C_{q,M} \int_{\tau_2}^{\tau_1} (\tau_1 - s)^{q-1} \|G(s, y_s + \bar{z}_s, y(s) + \bar{z}(s))\| ds \\
 &\leq a C_{q,M} [r^* \|\mu_1\|_{L^1(J, \mathbb{R}^+)} + r \|\mu_2\|_{L^1(J, \mathbb{R}^+)}] \int_{\tau_2}^{\tau_1} (\tau_1 - s)^{q-1} ds.
 \end{aligned}$$

As  $\tau_2 \rightarrow \tau_1$ ,  $I_3$  tends to zero. So  $\tilde{\Phi}(B_r)$  is equicontinuous.

Now let  $V$  be a subset of  $B_r$  such that  $V \subset \overline{\text{conv}}(\tilde{\Phi}(V) \cup \{0\})$ . Moreover, for any  $\varepsilon > 0$  and bounded set  $D$ , we can take a sequence  $\{v_n\}_{n=1}^\infty \subset D$  such that  $\alpha(D) \leq 2\alpha(\{v_n\}) + \varepsilon$  ([9, p.125]). Thus, for  $\{v_n\}_{n=1}^\infty \subset V$ , and using Lemmas

2.3–2.5 and (H2), we get

$$\begin{aligned}
\alpha(\tilde{\Phi}V) &\leq 2\alpha(\{\tilde{\Phi}v_n\}) + \varepsilon = 2 \sup_{t \in J} \alpha(\{\tilde{\Phi}v_n(t)\}) + \varepsilon \\
&= 2 \sup_{t \in J} \alpha\left(\left\{\int_0^t R(t-s) \int_0^s a(s,\tau) f(\tau, y_\tau + v_{n\tau}, y(\tau) + v_n(\tau)) d\tau ds\right\}\right) + \varepsilon \\
&\leq 4 \sup_{t \in J} \int_0^t \alpha\left(\left\{R(t-s) \int_0^s a(s,\tau) f(\tau, y_\tau + v_{n\tau}, y(\tau) + v_n(\tau)) d\tau ds\right\}\right) + \varepsilon \\
&\leq 8 \sup_{t \in J} \int_0^t \int_0^s \alpha(\{R(t-s)a(s,\tau)f(\tau, y_\tau + v_{n\tau}, y(\tau) + v_n(\tau)) d\tau ds\}) + \varepsilon \\
&\leq 8 a \sup_{t \in J} \int_0^t \int_0^s \alpha(\{R(t-s)f(\tau, y_\tau + v_{n\tau}, y(\tau) + v_n(\tau)) d\tau ds\}) + \varepsilon \\
&\leq 8 a \sup_{t \in J} \int_0^t \int_0^s \eta_t(s,\tau) [\alpha(v_n(\tau)) + \sup_{-\infty < \theta \leq 0} \alpha(v_n(\theta + \tau))] d\tau ds + \varepsilon \\
&\leq 8 a \sup_{t \in J} \int_0^t \int_0^s \eta_t(s,\tau) [\alpha(v_n) + \sup_{0 < \mu \leq \tau} \alpha(v_n(\mu))] d\tau ds + \varepsilon \\
&\leq 16 a \alpha(v_n) \sup_{t \in J} \int_0^t \int_0^s \eta_t(s,\tau) d\tau ds + \varepsilon \\
&\leq 16 a \eta^* \alpha(V) + \varepsilon.
\end{aligned}$$

Therefore, in view of Lemma 2.3, we have

$$\alpha(V) \leq \alpha(\tilde{\Phi}V) \leq 16 a \eta^* \alpha(V) + \varepsilon,$$

since  $\varepsilon$  is arbitrary we obtain that

$$\alpha(V) \leq 16 a \eta^* \alpha(V).$$

This means that

$$\alpha(V)(1 - 16 a \eta^*) \leq 0.$$

By (11) it follows that  $\alpha(V) = 0$ . In view of the Ascoli-Arzelà theorem,  $V$  is relatively compact in  $B_r$ . Applying now Theorem 2.7, we conclude that  $\tilde{\Phi}$  has a fixed point which is a solution of the problem (1).

#### 4. AN EXAMPLE

In this section we give an example to illustrate the above results. Consider the following fractional integrodifferential equations

$$\begin{aligned}
(14) \quad \frac{\partial^q}{\partial t^q} v(t, \zeta) &= \frac{\partial^2}{\partial \zeta^2} v(t, \zeta) + \int_0^t (t-s) \int_{-\infty}^0 \gamma_1(\theta) \sin(s|v(s+\theta, \zeta)|) d\theta ds \\
&+ \int_0^t (t-s) \frac{s^2}{2} \sin|v(s, \zeta)| \int_0^s \cos v(\iota, \zeta) d\iota ds
\end{aligned}$$

$$v(\theta, \zeta) = v_0(\theta, \zeta), \quad -\infty < \theta \leq 0,$$

where  $t, \zeta \in [0, 1]$ ,  $\gamma_1: (-\infty, 0] \rightarrow \mathbb{R}$ ,  $v_0: (-\infty, 0] \times [0, 1] \rightarrow \mathbb{R}$  are continuous functions, and  $\int_{-\infty}^0 |\gamma_1(\theta)| d\theta < \infty$ .

Set  $X = L^2([0, 1], \mathbb{R})$  and define  $A$  by

$$D(A) = H^2((0, 1)) \cap H_0^1((0, 1)),$$

$$Au = u''.$$

Then,  $A$  generates a compact, analytic semigroup  $S(t)$  of uniformly bounded, linear operators such that  $\|S(t)\| \leq 1$ .

Let the phase space  $\mathcal{B} = C((-\infty, 0], X)$ , the space of bounded uniformly continuous functions endowed with the following norm:

$$\|\varphi\|_{\mathcal{B}} = \sup_{-\infty < \theta \leq 0} |\varphi(\theta)|, \quad \forall \varphi \in \mathcal{B},$$

then we can see that  $C_1(t) = 0$  in (6).

For  $t \in [0, 1]$ ,  $\zeta \in [0, 1]$  and  $\varphi \in C((-\infty, 0], X)$ , we set

$$x(t)(\zeta) = v(t, \zeta),$$

$$\phi(\theta)(\zeta) = v_0(\theta, \zeta), \quad \theta \in (-\infty, 0],$$

$$a(t, s) = t - s,$$

$$f(t, \varphi, x(t))(\zeta) = \int_{-\infty}^0 \gamma_1(\theta) \sin(s|\varphi(\theta)(\zeta)|) d\theta + \frac{s^2}{2} \sin|x(t)(\zeta)| \int_0^s \cos x(\iota)(\zeta) d\iota.$$

Thus, problem (14) can be rewritten as the abstract problem (1).

Moreover, for  $t \in [0, 1]$ , we can see

$$\|f(t, \varphi, x(t))(\zeta)\| \leq t\|\varphi\|_{\mathcal{B}} \int_{-\infty}^0 |\gamma_1(\theta)| d\theta + \frac{t^3}{2} \|x(t)\|$$

$$= \mu_1(t)\|\varphi\|_{\mathcal{B}} + \mu_2(t)\|x(t)\|,$$

where

$$\mu_1(t) = t \int_{-\infty}^0 |\gamma_1(\theta)| d\theta, \quad \mu_2(t) = t^3/2.$$

Then (14) has a mild solution by Theorem 3.4.

For example, if we put

$$\gamma_1(\theta) = e^\theta, \quad q = \frac{1}{2},$$

then

$$C_{q,M} = \frac{1}{\Gamma(\frac{1}{2})} = 1/\sqrt{\pi}, \quad \|\mu_2\|_{L^1(J, \mathbb{R}^+)} = 1/8.$$

Thus, we see

$$\frac{aT^q C_{q,M} \|\mu_2\|_{L^1(J, \mathbb{R}^+)}}{q} = \frac{1}{4\sqrt{\pi}} < 1.$$

**Acknowledgement.** The authors are grateful to the referee for the careful reading of the paper.

#### REFERENCES

- [1] Abbas, S., Benchohra, M., N'Guérékata, G.M., *Topics in Fractional Differential Equations*, Springer, New York, 2012.
- [2] Agarwal, R. P., Belmekki, M., Benchohra, M., *A survey on semilinear differential equations and inclusions involving Riemann-Liouville fractional derivative*, Adv. Differential Equations **2009** (2009), 1–47, Article ID 981728.
- [3] Agarwal, R. P., Benchohra, M., Hamani, S., *A survey on existence results for boundary value problems of nonlinear fractional differential equations and inclusions*, Acta Appl. Math. **109** (2010), 973–1033.
- [4] Agarwal, R. P., Meehan, M., O'Regan, D., *Fixed Point Theory and Applications*, Cambridge University Press, 2001.
- [5] Appell, J. M., Kalitvin, A. S., Zabrejko, P. P., *Partial Integral Operators and Integrodifferential Equations*, vol. 230, Marcel Dekker, Inc., New York, 2000.
- [6] Baleanu, D., Diethelm, K., Scalas, E., Trujillo, J. J., *Fractional Calculus Models and Numerical Methods*, World Scientific Publishing, New York, 2012.
- [7] Banaś, J., Goebel, K., *Measures of Noncompactness in Banach Spaces*, Lecture Notes in Pure and Applied Mathematics, Marcel Dekker, New York, 1980.
- [8] Benchohra, M., Henderson, J., Ntouyas, S., Ouahab, A., *Existence results for fractional order functional differential equations with infinite delay*, J. Math. Anal. Appl. **338** (2008), 1340–1350.
- [9] Bothe, D., *Multivalued perturbations of  $m$ -accretive differential inclusions*, Israel J. Math. **108** (1998), 109–138.
- [10] Case, K. M., Zweifel, P. F., *Linear Transport Theory*, Addison-Wesley, Reading, MA, 1967.
- [11] Chandrasekhe, S., *Radiative Transfer*, Dover Publications, New York, 1960.
- [12] Corduneanu, C., Lakshmikantham, V., *Equations with unbounded delay*, Nonlinear Anal. **4** (1980), 831–877.
- [13] Diethelm, K., *The Analysis of Fractional Differential Equations*, Springer, Berlin, 2010.
- [14] El-Borai, M., *Some probability densities and fundamental solutions of fractional evolution equations*, Chaos, Solitons & Fractals **14** (2002), 433–440.
- [15] Hale, J. K., Kato, J., *Phase space for retarded equations with infinite delay*, Funkcial. Ekvac. **21** (1) (1978), 11–41.
- [16] Hale, J. K., Lunel, S. Verduyn, *Introduction to Functional–Differential Equations*, Applied Mathematical Sciences, vol. 99, Springer-Verlag, New York, 1993.
- [17] Heinz, H.-P., *On the behaviour of measures of noncompactness with respect to differentiation and integration of vector-valued functions*, Nonlinear Anal. **7** (12) (1983), 1351–1371.
- [18] Hilfer, R., *Applications of fractional calculus in physics*, Singapore, World Scientific, 2000.
- [19] Hino, Y., Murakami, S., Naito, T., *Functional Differential Equations with Infinite Delay*, Lecture Notes in Mathematics, vol. 1473, Springer-Verlag, Berlin, 1991.
- [20] Kilbas, A. A., Srivastava, Hari M., Trujillo, Juan J., *Theory and Applications of Fractional Differential Equations*, Elsevier Science B.V., Amsterdam, 2006.
- [21] Kolmanovskii, V., Myshkis, A., *Introduction to the Theory and Applications of Functional–Differential Equations*, Kluwer Academic Publishers, Dordrecht, 1999.
- [22] Lakshmikantham, V., Wen, L., Zhang, B., *Theory of Differential Equations with Unbounded Delay*, Mathematics and its Applications, Kluwer Academic Publishers, Dordrecht, 1994.

- [23] Li, F., Zhang, J., *Existence of mild solutions to fractional integrodifferential equations of neutral type with infinite delay*, Adv. Differential Equations **2011** (2011), 1–15, Article ID 963463.
- [24] Liang, J., Xiao, T.-J., Casteren, J. van, *A note on semilinear abstract functional differential and integrodifferential equations with infinite delay*, Appl. Math. Lett. **17** (4) (2004), 473–477.
- [25] Mainardi, F., Paradisi, P., Gorenflo, R., *Probability distributions generated by fractional diffusion equations*, Econophysics: An Emerging Science (Kertesz, J., Kondor, I., eds.), Kluwer Academic Publishers, Dordrecht, The Netherlands, 2000.
- [26] Miller, K. S., Ross, B., *An Introduction to the Fractional Calculus and Differential Equations*, John Wiley, New York, 1993.
- [27] Mönch, H., *Boundary value problems for nonlinear ordinary differential equations of second order in Banach spaces*, Nonlinear Anal. **4** (1980), 985–999.
- [28] Mophou, G. M., Nakoulima, O., N'Guérékata, G. M., *Existence results for some fractional differential equations with nonlocal conditions*, Nonlinear Stud. **17** (2010), 15–22.
- [29] Mophou, G. M., N'Guérékata, G. M., *Existence of mild solution for some fractional differential equations with nonlocal conditions*, Semigroup Forum **79** (2009), 315–322.
- [30] Obukhovskii, V., Yao, J.-C., *Some existence results for fractional functional differential equations*, Fixed Point Theory **11** (2010), 85–96.
- [31] Podlubny, I., *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [32] Samko, S. G., Kilbas, A. A., Marichev, O. I., *Fractional Integrals and Derivatives. Theory and Applications*, Gordon and Breach, Yverdon, 1993.
- [33] Tarasov, V. E., *Fractional Dynamics: Application of Fractional Calculus to Dynamics of Particles, Fields and Media*, Springer, Heidelberg; Higher Education Press, Beijing, 2010.
- [34] Wu, J., *Theory and Applications of Partial Functional Differential Equations*, Springer Verlag, New York, 1996.

CORRESPONDING AUTHOR:

MOUFFAK BENCHOIRA,

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ DE SIDI BEL-ABBÈS,

B.P. 89, 22000, SIDI BEL-ABBÈS, ALGÉRIE

*E-mail:* benchohra@univ-sba.dz