

Ryuji Kajikiya

Positive solutions of the  $p$ -Laplace Emden-Fowler equation in hollow thin symmetric domains

*Mathematica Bohemica*, Vol. 139 (2014), No. 2, 145–154

Persistent URL: <http://dml.cz/dmlcz/143845>

## Terms of use:

© Institute of Mathematics AS CR, 2014

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

POSITIVE SOLUTIONS OF THE  $p$ -LAPLACE EMDEN-FOWLER  
EQUATION IN HOLLOW THIN SYMMETRIC DOMAINS

RYUJI KAJIKIYA, Honjo-machi

(Received September 3, 2013)

*Abstract.* We study the existence of positive solutions for the  $p$ -Laplace Emden-Fowler equation. Let  $H$  and  $G$  be closed subgroups of the orthogonal group  $O(N)$  such that  $H \subsetneq G \subset O(N)$ . We denote the orbit of  $G$  through  $x \in \mathbb{R}^N$  by  $G(x)$ , i.e.,  $G(x) := \{gx : g \in G\}$ . We prove that if  $H(x) \subsetneq G(x)$  for all  $x \in \bar{\Omega}$  and the first eigenvalue of the  $p$ -Laplacian is large enough, then no  $H$  invariant least energy solution is  $G$  invariant. Here an  $H$  invariant least energy solution means a solution which achieves the minimum of the Rayleigh quotient among all  $H$  invariant functions. Therefore there exists an  $H$  invariant  $G$  non-invariant positive solution.

*Keywords:* Emden-Fowler equation; group invariant solution; least energy solution; positive solution; variational method

*MSC 2010:* 35J20, 35J25

## 1. INTRODUCTION

In this paper, we study the existence of positive solutions with partial symmetry for the  $p$ -Laplace Emden-Fowler equation

$$(1.1) \quad -\Delta_p u = u^{q-1}, \quad u > 0 \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega.$$

Here  $\Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u)$  is the  $p$ -Laplacian and  $\Omega$  is a bounded domain in  $\mathbb{R}^N$  with  $N \geq 2$ . Denote the critical exponent by  $p^* := Np/(N-p)$  if  $p < N$  and  $p^* := \infty$  if  $N \leq p$ . We assume that  $2 \leq p < q < p^*$ . We define the *Rayleigh quotient*  $R(u)$

---

This research was supported by the Grant-in-Aid for Scientific Research (C), Japan Society for the Promotion of Science (No. 24540179).

and the *Nehari manifold*  $\mathcal{N}$  by

$$R(u) := \left( \int_{\Omega} |\nabla u|^p \, dx \right) \left( \int_{\Omega} |u|^q \, dx \right)^{-p/q},$$

$$\mathcal{N} := \left\{ u \in W_0^{1,p}(\Omega) \setminus \{0\} : \int_{\Omega} (|\nabla u|^p - |u|^q) \, dx = 0 \right\},$$

where  $W_0^{1,p}(\Omega)$  denotes the Sobolev space. Let  $G$  be a closed subgroup of the orthogonal group  $O(N)$ . We call  $\Omega$  a *G invariant domain* if  $g(\Omega) = \Omega$  for any  $g \in G$ . We call  $u(x)$  a *G invariant solution* if  $u(gx) = u(x)$  for any  $g \in G$  and  $x \in \Omega$ . Then (1.1) has a  $G$  invariant positive solution. However, we are looking for an  $H$  invariant  $G$  non-invariant solution under a certain assumption on  $H$  and  $G$ , where  $H$  and  $G$  are closed subgroups of  $O(N)$  such that  $H \subsetneq G \subset O(N)$ . When  $\Omega$  is a  $G$  invariant domain, we denote the set of  $G$  invariant functions in  $W_0^{1,p}(\Omega)$  by  $W_0^{1,p}(\Omega, G)$ . Define  $\mathcal{N}(G) := \mathcal{N} \cap W_0^{1,p}(\Omega, G)$  and put

$$(1.2) \quad R_G := \inf\{R(u) : u \in W_0^{1,p}(\Omega, G) \setminus \{0\}\} = \inf\{R(u) : u \in \mathcal{N}(G)\}.$$

We call  $R_G$  a *G invariant least energy* and  $u$  a *G invariant least energy solution* if  $u \in \mathcal{N}(G)$  and  $R(u) = R_G$ . Such a minimizer exists and becomes a  $G$  invariant positive solution of (1.1). For  $x \in \mathbb{R}^N$ , we define the orbit  $G(x)$  through  $x$  by

$$(1.3) \quad G(x) := \{gx : g \in G\}.$$

Let  $\lambda_p(\Omega)$  denote the first eigenvalue of the  $p$ -Laplace eigenvalue problem

$$(1.4) \quad -\Delta_p u = \lambda |u|^{p-2} u \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega.$$

It is well known that the first eigenvalue is simple and the corresponding eigenfunction is positive (see [7]). We state the main result of this paper.

**Theorem 1.1.** *Assume that  $2 \leq p < q < p^*$ . Let  $G$  and  $H$  be closed subgroups of  $O(N)$  and let  $U$  be a  $G$  invariant bounded domain in  $\mathbb{R}^N$  such that  $H \subsetneq G$  and  $H(x) \subsetneq G(x)$  for all  $x \in \overline{U}$ . Then there exists a constant  $C > 0$  depending only on  $G, H, U, p$  and  $q$  such that if  $\Omega$  is a  $G$  invariant subdomain of  $U$  and if  $\lambda_p(\Omega) > C$ , then  $R_H < R_G$ . Therefore no  $H$  invariant least energy solution is  $G$  invariant.*

The existence of multiple positive solutions of (1.1) on the sphere has been obtained by Kristály [6] also, in which the nonlinear term is asymptotically critical. We observe the Faber-Krahn inequality (see [1]),  $\lambda_p(\Omega) \geq C_{N,p} |\Omega|^{-p/N}$ , where  $C_{N,p} > 0$  is a constant independent of  $\Omega$  and  $|\Omega|$  denotes the volume of  $\Omega$ . Then we obtain the next corollary.

**Corollary 1.2.** *Under the assumption of Theorem 1.1, there exists a constant  $\delta > 0$  depending only on  $G, H, U, p$  and  $q$  such that if  $\Omega$  is a  $G$  invariant subdomain of  $U$  and if  $|\Omega| < \delta$ , then  $R_H < R_G$ .*

We give a simple example of  $H, G$  and  $\Omega$ . A subgroup  $H$  of  $O(N)$  is said to be transitive on the sphere  $S^{N-1}$  if  $H(x) = S^{N-1}$  for  $x \in S^{N-1}$ . All transitive Lie groups were classified by Montgomery and Samelson [8] and Borel [2].

**Example 1.3.** Let  $G := O(N)$  and let  $H$  be any non-transitive closed subgroup of  $O(N)$ . Let  $\Omega$  be an annulus  $1 < |x| < 1 + \varepsilon$  with  $\varepsilon > 0$ . If  $\varepsilon > 0$  is small enough, then no  $H$  invariant least energy solution is radially symmetric.

## 2. LEAST ENERGY SOLUTIONS

Let  $L^r(\Omega, G)$  denote the set of  $G$  invariant functions in  $L^r(\Omega)$ . Define the  $L^2(\Omega)$  inner product and the  $H_0^1(\Omega)$  inner product by

$$(u, v)_{L^2} := \int_{\Omega} uv \, dx, \quad (u, v)_{H_0^1} := \int_{\Omega} \nabla u \nabla v \, dx.$$

We define the orthogonal complements of  $L^2(\Omega, G)$  and  $H_0^1(\Omega, G)$  by

$$\begin{aligned} L^2(\Omega, G)^\perp &:= \{u \in L^2(\Omega) : (u, v)_{L^2} = 0 \text{ for all } v \in L^2(\Omega, G)\}, \\ H_0^1(\Omega, G)^\perp &:= \{u \in H_0^1(\Omega) : (u, v)_{H_0^1} = 0 \text{ for all } v \in H_0^1(\Omega, G)\}. \end{aligned}$$

**Lemma 2.1** ([3], Lemma 3.2). *We have the following assertions.*

- (i)  $H_0^1(\Omega, G)^\perp \subset L^2(\Omega, G)^\perp$ .
- (ii) *Let  $1 \leq r, s \leq \infty$  with  $1/r + 1/s = 1$ . If  $u \in L^r(\Omega) \cap L^2(\Omega, G)^\perp$  and  $v \in L^s(\Omega, G)$ , then  $\int_{\Omega} uv \, dx = 0$ .*

Since  $p \geq 2$ , the Rayleigh quotient  $R$  is twice differentiable in the sense of the Fréchet derivative. Then  $R''(u)vw$  is a bilinear form of  $v$  and  $w$ . We need the formula of the special case  $R''(u)w^2$  only.

**Lemma 2.2.** *Let  $u$  be a positive solution of (1.1). For  $w \in W_0^{1,p}(\Omega)$ , we have*

$$(2.1) \quad \begin{aligned} R''(u)w^2 &= p(p-2) \left( \int |\nabla u|^p dx \right)^{-p/q} \int |\nabla u|^{p-4} (\nabla u \cdot \nabla w)^2 dx \\ &\quad + p \left( \int |\nabla u|^p dx \right)^{-p/q} \int |\nabla u|^{p-2} |\nabla w|^2 dx \\ &\quad + p(q-p) \left( \int |\nabla u|^p dx \right)^{-(p+q)/q} \left( \int u^{q-1} w dx \right)^2 \\ &\quad - p(q-1) \left( \int |\nabla u|^p dx \right)^{-p/q} \int u^{q-2} w^2 dx. \end{aligned}$$

Here all integrals are taken over  $\Omega$ .

*Proof.* Multiplying (1.1) by  $u$  or  $w$  and integrating it over  $\Omega$ , we have

$$\int |\nabla u|^p dx = \int u^q dx, \quad \int |\nabla u|^{p-2} \nabla u \nabla w dx = \int u^{q-1} w dx.$$

Using the above identities and differentiating  $R(u + tw)$  twice at  $t = 0$ , we obtain (2.1).  $\square$

The next proposition plays the most important role in the paper.

**Proposition 2.3.** *Let  $u$  be a  $G$  invariant least energy solution of (1.1) and let  $\Omega_1$  be a  $G$  invariant bounded open set such that  $\Omega \subset \Omega_1$ . Let  $\varphi$  be a function in  $H_0^1(\Omega_1, G)^\perp \cap W^{1,\infty}(\Omega_1)$  which satisfies*

$$(2.2) \quad \int_{\Omega} |\nabla u|^{p-2} u^2 |\nabla \varphi|^2 dx < \frac{q-p}{2(2q-p-1)} \int_{\Omega} |\nabla u|^p \varphi^2 dx.$$

Then  $R((1 + \varepsilon\varphi)u) < R(u)$  for  $\varepsilon > 0$  small enough.

*Proof.* Set  $v := (1 + \varepsilon\varphi)u$  and define  $w := \varphi u$ . Then  $v = u + \varepsilon w$ . Since  $u \in C^1(\overline{\Omega}) \cap H_0^1(\Omega)$ ,  $w$  and  $v$  belong to  $H_0^1(\Omega)$ . Since  $u$  is a solution of (1.1),  $R'(u)$  vanishes. The Taylor theorem ensures that

$$R(v) = R(u) + (\varepsilon^2/2)R''(u)w^2 + o(\varepsilon^2),$$

as  $\varepsilon \rightarrow 0$ . Here  $o(\varepsilon^2)/\varepsilon^2 \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . To prove  $R(v) < R(u)$  for  $\varepsilon > 0$  small enough, we have only to show that  $R''(u)w^2 < 0$ . We substitute  $w = \varphi u$  in (2.1) and compute all terms on the right hand side. We extend  $u$  by setting  $u(x) = 0$  outside  $\Omega$ . By Lemma 2.1, we see that

$$u^q \in L^2(\Omega_1, G), \quad \varphi \in H_0^1(\Omega_1, G)^\perp \subset L^2(\Omega_1, G)^\perp.$$

Consequently,

$$\int_{\Omega} u^{q-1} w \, dx = \int_{\Omega_1} u^q \varphi \, dx = 0.$$

It is easy to see that

$$\begin{aligned} \nabla u \cdot \nabla w &= |\nabla u|^2 \varphi + u \nabla u \cdot \nabla \varphi, \\ |\nabla w|^2 &= |\nabla u|^2 \varphi^2 + 2u \varphi \nabla u \cdot \nabla \varphi + u^2 |\nabla \varphi|^2. \end{aligned}$$

Substituting the above identities in (2.1) and putting

$$A := \left( \int |\nabla u|^p \, dx \right)^{-p/q},$$

we have

$$\begin{aligned} (2.3) \quad R''(u)w^2 &= p(p-1)A \int |\nabla u|^p \varphi^2 \, dx \\ &\quad + 2p(p-1)A \int |\nabla u|^{p-2} u \varphi \nabla u \cdot \nabla \varphi \, dx \\ &\quad + p(p-2)A \int |\nabla u|^{p-4} u^2 (\nabla u \cdot \nabla \varphi)^2 \, dx \\ &\quad + pA \int |\nabla u|^{p-2} u^2 |\nabla \varphi|^2 \, dx - p(q-1)A \int u^q \varphi^2 \, dx. \end{aligned}$$

Now, multiplying (1.1) by  $u\varphi^2$  and integrating over  $\Omega$ , we see that

$$\int u^q \varphi^2 \, dx = \int (|\nabla u|^p \varphi^2 + 2|\nabla u|^{p-2} u \varphi \nabla u \cdot \nabla \varphi) \, dx.$$

Substituting the above identity in (2.3), we obtain

$$\begin{aligned} (2.4) \quad R''(u)w^2 &= -p(q-p)A \int |\nabla u|^p \varphi^2 \, dx \\ &\quad - 2p(q-p)A \int |\nabla u|^{p-2} u \varphi \nabla u \cdot \nabla \varphi \, dx \\ &\quad + p(p-2)A \int |\nabla u|^{p-4} u^2 (\nabla u \cdot \nabla \varphi)^2 \, dx \\ &\quad + pA \int |\nabla u|^{p-2} u^2 |\nabla \varphi|^2 \, dx. \end{aligned}$$

We use the Schwarz inequality

$$|u \varphi \nabla u \cdot \nabla \varphi| \leq \frac{1}{4} |\nabla u|^2 \varphi^2 + u^2 |\nabla \varphi|^2$$

in the second integral on the right hand side of (2.4) and employ  $|\nabla u \cdot \nabla \varphi| \leq |\nabla u| |\nabla \varphi|$  in the third integral. Then we obtain

$$\begin{aligned} R''(u)w^2 &\leq -\frac{1}{2}p(q-p)A \int |\nabla u|^p \varphi^2 \, dx \\ &\quad + p(2q-p-1)A \int |\nabla u|^{p-2} u^2 |\nabla \varphi|^2 \, dx. \end{aligned}$$

The right hand side is negative because of (2.2). The proof is complete.  $\square$

To prove the main theorems, we need the Haar measure. Since  $G$  is a compact Lie group, it has a unique Haar measure  $dg$ . It is a positive Lebesgue measure which satisfies

$$\begin{aligned} \int_G f(hg) \, dg &= \int_G f(gh) \, dg = \int_G f(g^{-1}) \, dg = \int_G f(g) \, dg, \\ \int_G f(g) \, dg &> 0 \quad \text{if } f \geq 0, f \not\equiv 0, \quad \int_G 1 \, dg = 1, \end{aligned}$$

for any  $h \in G$  and any real valued integrable function  $f$  on  $G$  (see [9] for more details).

Let  $M(N)$  be a linear space consisting of all  $N \times N$  real matrices, which is equipped with the norm

$$\|g\| := \max_{|x| \leq 1} |gx| \quad \text{for } g \in M(N).$$

For  $g_0 \in G$  and  $r > 0$  we define a ball  $B(g_0, r; G)$  in  $G$  by

$$B(g_0, r; G) := \{g \in G : \|g - g_0\| < r\}.$$

Then the volume of  $B(g_0, r; G)$  is defined by

$$|B(g_0, r; G)| := \int_{B(g_0, r; G)} 1 \, dg.$$

Using the invariance of the Haar measure, we have the next lemma.

**Lemma 2.4** ([4], Lemma 5.6). *Let  $G$  be a closed subgroup of  $O(N)$ . Then the volume  $|B(g_0, r; G)|$  does not depend on  $g_0 \in G$  but does on  $r$  only.*

### 3. PROOF OF THE MAIN RESULTS

In this section, we prove the main theorem. Let  $H$  and  $G$  be as in Theorem 1.1. Since  $G$  and  $H$  are compact groups, we can define

$$Q(x, g) := \min_{h \in H} |gx - hx|, \quad P(x) := \max_{g \in G} Q(x, g).$$

**Lemma 3.1.** *We have*

$$|P(x) - P(y)| \leq 2|x - y| \quad \text{for } x, y \in \mathbb{R}^N.$$

*Proof.* By the same computation as in our paper [3], Lemma 2.1 or [4], Lemma 5.5, we obtain the lemma.  $\square$

Recall the assumption of Theorem 1.1 that  $H(x) \not\subset G(x)$  for all  $x \in \overline{U}$ . This implies that  $P(x) > 0$  for  $x \in \overline{U}$ . Since  $P(x)$  is continuous by Lemma 3.1, the minimum of  $P(x)$  on  $\overline{U}$  is positive. We define

$$(3.1) \quad \delta := \frac{1}{4} \min_{\overline{U}} P(x) > 0.$$

Then for any  $x \in \overline{U}$  there exists a  $g \in G$  such that

$$(3.2) \quad |gx - hx| \geq 4\delta > 0 \quad \text{for any } h \in H.$$

To prove Theorem 1.1, we shall construct a function  $\varphi$  which satisfies (2.2) and belongs to  $H_0^1(\Omega_1, H)$ . Let  $\delta > 0$  be defined by (3.1). Choose  $\Phi \in C^1(\mathbb{R})$  which satisfies  $0 \leq \Phi(r) \leq 1$  in  $\mathbb{R}$ ,  $\Phi(r) = 1$  for  $r \leq \delta$ ,  $\Phi(r) = 0$  for  $r \geq 2\delta$  and  $-2/\delta \leq \Phi'(r) \leq 0$  in  $(\delta, 2\delta)$ . Put  $r = |x|$ . Then  $\Phi(|x|)$  is a radial function whose support is in  $|x| \leq 2\delta$ .

**Definition 3.2.** We denote the Haar measures on  $H$  and  $G$  by  $dh$  and  $dg$ , respectively. Let  $x_0 \in \Omega$  be determined later on. We define

$$\begin{aligned} \varphi(x) &:= \int_G \Phi(|x - gx_0|) dg - \int_H \Phi(|x - hx_0|) dh, \\ \text{dist}(x, \Omega) &:= \inf\{|x - y| : y \in \Omega\}, \\ \Omega_1 &:= \{x \in \mathbb{R}^N : \text{dist}(x, \Omega) < 2\delta\}. \end{aligned}$$

**Lemma 3.3** ([4], [5]). *Function  $\varphi$  belongs to  $H_0^1(\Omega_1, G)^\perp \cap H_0^1(\Omega_1, H)$ .*

Since  $U$  is bounded, we define  $M := \sup_{x \in U} |x|$  and  $\mu := \delta/M$ . Then  $\mu$  depends only on  $G, H$  and  $U$ . We denote the volume of  $B(g_0, \mu; G)$  by  $c_0$ , i.e.,

$$(3.3) \quad c_0 := |B(g_0, \mu; G)| = \int_{B(g_0, \mu; G)} 1 dg.$$

By Lemma 2.4,  $c_0$  depends not on  $g_0$  but on  $\mu$ , hence it depends only on  $G, H$  and  $U$ . Let  $B(x, r)$  denote the ball in  $\mathbb{R}^N$  which is centered at  $x$  with radius  $r > 0$ .



**Lemma 3.4** ([4], [5]). *For any  $x_0 \in \Omega$ , there exists a  $g_0 \in G$  such that*

$$(3.4) \quad \varphi(x) \geq c_0 > 0 \quad \text{for } x \in B(g_0 x_0, \delta/2).$$

*In particular,  $\varphi \not\equiv 0$  in  $\Omega$ .*

Let  $\delta$  be defined by (3.1). We choose a finite covering  $B(y_i, \delta/4)$  with  $y_1, \dots, y_k \in \overline{U}$  such that

$$(3.5) \quad \overline{U} \subset \bigcup_{i=1}^k B(y_i, \delta/4) \quad \text{with some } k \in \mathbb{N}.$$

Hereafter we fix  $k$  and  $y_1, \dots, y_k$  which satisfy the above inclusion.

**Lemma 3.5.** *Let  $\Omega$  be a  $G$  invariant subdomain of  $U$  and let  $u$  be a  $G$  invariant least energy solution. Extend  $u$  by setting  $u(x) = 0$  outside  $\Omega$ . Then there exists an  $x_0 \in \Omega$  such that*

$$\int_{\Omega} |\nabla u|^p \, dx \leq k \int_{B(x_0, \delta/2)} |\nabla u|^p \, dx.$$

*Proof.* Choose  $i \in \{1, 2, \dots, k\}$  such that

$$\int_{B(y_i, \delta/4)} |\nabla u|^p \, dx = \max_j \int_{B(y_j, \delta/4)} |\nabla u|^p \, dx.$$

Then we have

$$\int_{\Omega} |\nabla u|^p \, dx \leq k \int_{B(y_i, \delta/4)} |\nabla u|^p \, dx.$$

Observe that  $\Omega \cap B(y_i, \delta/4) \neq \emptyset$ . Otherwise the right hand side vanishes. We choose an  $x_0 \in \Omega \cap B(y_i, \delta/4)$ . Then we have

$$\int_{B(y_i, \delta/4)} |\nabla u|^p \, dx \leq \int_{B(x_0, \delta/2)} |\nabla u|^p \, dx.$$

Combining the two above inequalities, we obtain the conclusion. □

**Lemma 3.6.** *Let  $\lambda_p$  be the first eigenvalue of (1.4). Then*

$$\int_{\Omega} |\nabla v|^{p-2} v^2 \, dx \leq \lambda_p^{-2/p} \|\nabla v\|_p^p \quad \text{for any } v \in W_0^{1,p}(\Omega).$$

*Proof.* From the variational characterization of the first eigenvalue, it follows that for  $v \in W_0^{1,p}(\Omega)$ ,

$$\lambda_p \int_{\Omega} |v|^p \, dx \leq \int_{\Omega} |\nabla v|^p \, dx,$$

or equivalently

$$\|v\|_p \leq \lambda_p^{-1/p} \|\nabla v\|_p.$$

Using this inequality with the Hölder inequality, we get

$$\int_{\Omega} |\nabla v|^{p-2} v^2 \, dx \leq \|\nabla v\|_p^{p-2} \|v\|_p^2 \leq \lambda_p^{-2/p} \|\nabla v\|_p^p.$$

□

Define  $\delta$ ,  $c_0$  and  $k$  by (3.1), (3.3) and (3.5), respectively, and then determine  $x_0$  by Lemma 3.5. Thus  $\varphi(x)$  is well defined by Definition 3.2. To prove Theorem 1.1, we define

$$C := [32\delta^{-2}kc_0^{-2}(2q-p-1)/(q-p)]^{p/2},$$

which depends only on  $G$ ,  $H$ ,  $U$ ,  $p$  and  $q$ . We conclude this paper by proving Theorem 1.1.

*Proof of Theorem 1.1.* Let  $C$  be as above. Suppose that  $\lambda_p(\Omega) > C$ . We shall show that  $\varphi$  satisfies (2.2). Since  $|\Phi'(r)| \leq 2/\delta$  by the definition of  $\Phi$ , we have  $|\nabla\varphi| \leq 4/\delta$ . This inequality and Lemmas 3.6 and 3.5 show that

$$\begin{aligned} \int_{\Omega} |\nabla u|^{p-2} u^2 |\nabla\varphi|^2 \, dx &\leq 16\delta^{-2} \lambda_p^{-2/p} \|\nabla u\|_p^p \\ &\leq 16\delta^{-2} \lambda_p^{-2/p} k \int_{B(x_0, \delta/2)} |\nabla u|^p \, dx. \end{aligned}$$

By Lemma 3.4, we choose  $g_0 \in G$  satisfying (3.4). Since  $u$  is  $G$  invariant, the last integral is estimated as

$$\int_{B(x_0, \delta/2)} |\nabla u|^p \, dx = \int_{B(g_0 x_0, \delta/2)} |\nabla u|^p \, dx \leq c_0^{-2} \int_{B(g_0 x_0, \delta/2)} |\nabla u|^p \varphi^2 \, dx.$$

Combining the two above inequalities, we have

$$\int_{\Omega} |\nabla u|^{p-2} u^2 |\nabla\varphi|^2 \, dx \leq 16\delta^{-2} \lambda_p^{-2/p} k c_0^{-2} \int_{B(g_0 x_0, \delta/2)} |\nabla u|^p \varphi^2 \, dx.$$

Since  $\lambda_p(\Omega) > C$ , we obtain (2.2). Since  $\varphi \in H_0^1(\Omega_1, H)$  by Lemma 3.3,  $v := (1 + \varepsilon\varphi)u$  belongs to  $H_0^1(\Omega, H)$ . By Proposition 2.3, we conclude that  $R_H \leq R(v) < R(u) = R_G$ . The proof is complete.  $\square$

### References

- [1] *T. Bhattacharya*: A proof of the Faber-Krahn inequality for the first eigenvalue of the  $p$ -Laplacian. *Ann. Mat. Pura Appl.*, IV. Ser. *177* (1999), 225–240.
- [2] *A. Borel*: Le plan projectif des octaves et les sphères comme espaces homogènes. *C. R. Acad. Sci., Paris* *230* (1950), 1378–1380. (In French.)
- [3] *R. Kajikiya*: Least energy solutions of the Emden-Fowler equation in hollow thin symmetric domains. *J. Math. Anal. Appl.* *406* (2013), 277–286.
- [4] *R. Kajikiya*: Multiple positive solutions of the Emden-Fowler equation in hollow thin symmetric domains. *Calc. Var. Partial Differ. Equ.* To appear.
- [5] *R. Kajikiya*: Partially symmetric solutions of the generalized Hénon equation in symmetric domains. *Topol. Methods Nonlinear Anal.* To appear.
- [6] *A. Kristály*: Asymptotically critical problems on higher-dimensional spheres. *Discrete Contin. Dyn. Syst.* *23* (2009), 919–935.
- [7] *P. Lindqvist*: On the equation  $\operatorname{div}(|\nabla u|^{p-2}\nabla u) + \lambda|u|^{p-2}u = 0$ . *Proc. Am. Math. Soc.* *109* (1990), 157–164.
- [8] *D. Montgomery, H. Samelson*: Transformation groups of spheres. *Ann. Math. (2)* *44* (1943), 454–470.
- [9] *L. S. Pontryagin*: *Topological Groups*. Transl. from the Russian, (3rd ed.), *Classics of Soviet Mathematics*, Gordon and Breach, New York, 1986.

*Author's address:* *Ryuji Kajikiya*, Department of Mathematics, Faculty of Science and Engineering, Saga University, 1 Honjo-machi, Saga, 840-8502, Japan, e-mail: [kajikiya@ms.saga-u.ac.jp](mailto:kajikiya@ms.saga-u.ac.jp).