

Dániel T. Soukup; Lajos Soukup
Partitioning bases of topological spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 55 (2014), No. 4, 537--566

Persistent URL: <http://dml.cz/dmlcz/143977>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2014

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Partitioning bases of topological spaces

DÁNIEL T. SOUKUP, LAJOS SOUKUP*

Abstract. We investigate whether an arbitrary base for a dense-in-itself topological space can be partitioned into two bases. We prove that every base for a T_3 Lindelöf topology can be partitioned into two bases while there exists a consistent example of a first-countable, 0-dimensional, Hausdorff space of size 2^ω and weight ω_1 which admits a point countable base without a partition to two bases.

Keywords: base; resolvable; partition

Classification: 54A35, 03E35, 54A25

1. Introduction

At the Trends in Set Theory conference in Warsaw, Barnabás Farkas¹ raised the natural question whether one can partition any given base for a topological space into two bases; we will call this property being *base resolvable*. Note that every space with an isolated point is not base resolvable; hence, from now on by *space* we mean a *dense-in-itself topological space*. The aim of this paper is to present two streams of results: in the first part of the article, we will show that certain natural classes of spaces are base resolvable. In the second part, we present a method to construct non base resolvable spaces.

The paper is structured as follows: In Section 2, we will start with general observations about bases and we prove that metric spaces and weakly separated spaces are base resolvable. This section also serves as an introduction to the methods that will be applied in Section 3 where we prove one of our main results in Theorem 3.7 that every T_3 (locally) Lindelöf space is base resolvable.

In Section 4, we investigate base resolvability from a purely combinatorial viewpoint which leads to further results. We show that every hereditarily Lindelöf space (without any separation axioms) is base resolvable and any base for a T_1 topology which is closed under finite unions can be partitioned into two bases, see Theorem 4.8 and 4.9 respectively.

Next in Theorem 5.5, we prove that every base \mathbb{B} for a space X (resolvable or not) contains a large *negligible* portion, i.e., there is $\mathcal{U} \in [\mathbb{B}]^{|\mathbb{B}|}$ such that $\mathbb{B} \setminus \mathcal{U}$ is still a base for X .

The preparation of this paper was partially supported by OTKA grant K 83726.

*Corresponding author.

DOI 10.14712/1213-7243.014.404

¹personal communication

The second part of the paper starts with Section 6; here, we isolate a partition property, denoted by $\mathbb{P} \rightarrow (I_\omega)_2^1$, of the partial order $\mathbb{P} = (\mathbb{B}, \supseteq)$ associated to a base \mathbb{B} which is closely related to base resolvability. We will construct a partial order \mathbb{P} with this property in Theorem 6.6 and deduce the existence of a T_0 non base resolvable topology (in ZFC) in Corollary 6.15.

Next, in Section 7 we present a ccc forcing (of size ω_1) which introduces a first-countable, 0-dimensional, Hausdorff space X of size 2^ω and weight ω_1 such that X is not base resolvable. The main ideas of the construction already appear in Section 6, however the details here are much more subtle and the proofs are more technical.

The paper finishes with a list of open problems in Section 8. We remark that Section 7 was prepared by the second author and the rest of the paper is the work of the first author.

The first author would like to thank his PhD advisor, William Weiss, for the long hours of useful discussions. Both authors are grateful for the help of all the people they discussed the problems at hand, especially Allan Dow, István Juhász, Arnie Miller, Assaf Rinot, Santi Spadaro, Zoltán Szentmiklóssy and Zoltán Vidnyánszky. Finally, we thank Barnabás Farkas for the excellent question!

2. General results

In this section, we prove some basic results concerning partitions of families of sets and partitions of bases; these proofs will introduce us to the more involved techniques of the upcoming sections.

Definition 2.1. We say that a family of sets \mathcal{A} is **well-founded** if the poset $\langle \mathcal{A}, \supseteq \rangle$ is well-founded, i.e., there is no strictly decreasing infinite chain $A_0 \supsetneq A_1 \supsetneq A_2 \supsetneq \dots$ in \mathcal{A} .

\mathcal{A} is **weakly increasing** if there is a well order \prec of \mathcal{A} such that $A \prec B$ implies that $B \setminus A \neq \emptyset$.

Proposition 2.2. *Every family of sets \mathcal{A} contains a weakly increasing, and so well-founded subfamily \mathcal{B} with*

$$\bigcup \mathcal{A} = \bigcup \mathcal{B}.$$

PROOF: Fix an arbitrary well-ordering \prec of \mathcal{A} and let

$$(2.1) \quad \mathcal{B} = \{B \in \mathcal{A} : B \setminus A \neq \emptyset \text{ for all } A \prec B\}.$$

If $C \prec B$ for $C, B \in \mathcal{B}$, then $B \setminus C \neq \emptyset$, so \prec witnesses that \mathcal{B} is weakly increasing.

To verify $\bigcup \mathcal{A} = \bigcup \mathcal{B}$ pick an arbitrary $p \in \bigcup \mathcal{A}$ and let

$$(2.2) \quad B = \min_{\prec} \{A \in \mathcal{A} : p \in A\}.$$

Then $p \in B \setminus A$ for all $A \prec B$, so $B \in \mathcal{B}$. Thus $\bigcup \mathcal{A} = \bigcup \mathcal{B}$. □

Definition 2.3. A base \mathbb{B} for a space X is **resolvable** if it can be decomposed into two bases. A space X is **base resolvable** if every base of X is resolvable.

Recall that by *space* we will mean a dense-in-itself topological space throughout the paper.

Partitioning sets with additional structure is a highly investigated theme in mathematics; let us cite a classical result of A.H. Stone which is relevant to our case:

Theorem 2.4 (A.H. Stone, [2]). *Every partially ordered set (\mathbb{P}, \leq) without maximal elements can be partitioned into two cofinal subsets.*

Proposition 2.5. *Suppose that (X, τ) is a topological space and $p \in X$.*

- (1) *Every neighborhood base at p can be partitioned into two neighborhood bases.*
- (2) *Every π -base can be partitioned into two π -bases.*
- (3) *If \mathcal{B} is a neighborhood base at p and $\mathcal{B} = \mathcal{B}_0 \cup \mathcal{B}_1$ then either \mathcal{B}_0 or \mathcal{B}_1 is a neighborhood base at p .*
- (4) *If \mathcal{B} is a base and $\mathcal{U} \subset \mathcal{B}$ is well founded then $\mathcal{B} \setminus \mathcal{U}$ is a base.*
- (5) *Every base can be partitioned into a cover and a base.*

PROOF: (1) and (2) follow from Theorem 2.4.

Indeed, write $\tau_x = \{U \in \tau : x \in U\}$ for $x \in X$ and observe that $\mathcal{B} \subset \tau_x$ is a neighborhood base at x if and only if \mathcal{B} is cofinal in $\langle \tau_x, \supset \rangle$. By Theorem 2.4, every neighborhood base at p can be partitioned into two cofinal subsets of $\langle \tau_p, \supset \rangle$, i.e., into two neighborhood bases at p . So (1) holds.

To prove (2), observe that $\mathcal{B} \subset \tau$ is a π -base if and only if \mathcal{U} is cofinal in $\langle \tau, \supset \rangle$. By Theorem 2.4, every π -base can be partitioned into two cofinal subsets, i.e., into two π -bases.

(3) If \mathcal{B}_0 is not a neighborhood base at p then there is an element $V \in \tau_p$ which does not contain any element of \mathcal{B} . Thus $\mathcal{B} \cap \mathcal{P}(V) = \mathcal{B}_1 \cap \mathcal{P}(V)$, so \mathcal{B}_1 is a neighborhood base at p .

(4) Let $x \in X$. Then $\tau_x \cap \mathcal{B}$ is a neighborhood base at x . Since $\tau_x \cap \mathcal{U}$ is well-founded, $\tau_x \cap \mathcal{U}$ is not a neighborhood base at x . Thus, by (3), $\tau_x \cap (\mathcal{B} \setminus \mathcal{U})$ is a neighborhood base at x .

Since x was arbitrary, we proved that $\mathcal{B} \setminus \mathcal{U}$ is a base.

(5) Every base \mathcal{B} contains a well-founded cover \mathcal{U} by Proposition 2.2 while $\mathcal{B} \setminus \mathcal{U}$ is still a base of X by (4). □

A family \mathbb{B} of open subset of a space $\langle X, \tau \rangle$ is a base if and only if every nonempty open set is the union of some subfamily of \mathbb{B} . This fact implies the following:

Observation 2.6. *Suppose that (X, τ) is a topological space, $\mathbb{B}_i \subset \tau$ for $i < 2$ and \mathbb{B}_0 is a base.*

- (1) *If for every $U \in \mathbb{B}_0$ there is $\mathcal{U} \subset \mathbb{B}_1$ with $U = \bigcup \mathcal{U}$ then \mathbb{B}_1 is a base as well.*

- (2) If X is T_3 and for every $U, V \in \mathbb{B}_0$ with $\bar{U} \subset V$ there is $\mathcal{U} \subset \mathbb{B}_1$ with $\bar{U} \subset \bigcup \mathcal{U} \subset V$ then \mathbb{B}_1 is a base as well.

Now we prove our first general result.

Proposition 2.7. *Every space with a σ -disjoint base is base resolvable; in particular, every metrizable space is base resolvable.*

PROOF: Fix a space X with a base $\bigcup \{\mathbb{E}_n : n \in \omega\}$ where \mathbb{E}_n is a disjoint family of open sets for each $n \in \omega$; fix an arbitrary base \mathbb{B} as well which we aim to partition.

By induction on $n \in \omega$, construct $\mathbb{B}_{i,n} \subseteq \mathbb{B}$ for $i < 2$ such that

- (1) $\mathbb{B}_{i,n}$ is well founded for $i < 2, n \in \omega$,
- (2) $\mathbb{B}_{i,n} \cap \mathbb{B}_{j,m} = \emptyset$ if $i, j < 2, n, m \in \omega$ and $(i, n) \neq (j, m)$,
- (3) for every $V \in \mathbb{E}_n$ and $i < 2$ there is $\mathcal{U} \subseteq \mathbb{B}_{i,n}$ such that $\bigcup \mathcal{U} = V$.

Assume that $\{\mathbb{B}_{i,k} : i < 2, k < n\}$ was constructed. By Proposition 2.5(4) property (1) assures that $\mathbb{B} \setminus \bigcup \{\mathbb{B}_{i,k} : i < 2, k < n\}$ is still a base of X . Thus, by Proposition 2.2, for each $E \in \mathbb{E}_n$ we can choose a well-founded family $\mathcal{U}_E \subset \mathbb{B} \setminus \bigcup \{\mathbb{B}_{i,k} : i < 2, k < n\}$ such that $E = \bigcup \mathcal{U}_E$. Let

$$\mathbb{B}_{0,n} = \bigcup \{\mathcal{U}_E : E \in \mathbb{E}_n\}.$$

Since the elements of \mathbb{E}_n are pairwise disjoint, $\mathbb{B}_{0,n}$ is well-founded as well.

To obtain $\mathbb{B}_{1,n}$ repeat the construction of $\mathbb{B}_{0,n}$ using $\mathbb{B} \setminus (\bigcup \{\mathbb{B}_{i,k} : i < 2, k < n\} \cup \mathbb{B}_{0,n})$ instead of $\mathbb{B} \setminus \bigcup \{\mathbb{B}_{i,k} : i < 2, k < n\}$.

Let $\mathbb{B}_i = \bigcup \{\mathbb{B}_{i,n} : n \in \omega\}$ for $i < 2$. Then property (3) and Observation 2.6(1) implies that \mathbb{B}_i is a base for $i < 2$. □

Note that every σ -disjoint base is point countable. On the other hand our example of an irresolvable base constructed in Section 7 is point countable.

A somewhat similar technique, which will be used later as well, gives the following result:

Proposition 2.8. *Suppose that a regular space X satisfies $L(X) < \kappa = w(X) = \min\{\chi(x, X) : x \in X\}$. Then X is base resolvable.*

Recall that $L(X)$, the *Lindelöf number* of X , is the minimal cardinality κ such that every open cover of X contains a subcover of size κ . The *weight* of X is

$$w(X) = \min\{|\mathbb{B}| : \mathbb{B} \text{ is a base of } X\}$$

and the *character* of a point $x \in X$ is

$$\chi(x, X) = \min\{|\mathcal{U}| : \mathcal{U} \text{ is a neighbourhood base of } x\}.$$

PROOF: It is well known that any base contains a base of size $w(X)$; therefore it suffices to show that any base \mathbb{B} of size $w(X)$ can be partitioned into two bases. Let us fix an enumeration $\{(U_\alpha, V_\alpha) : \alpha < \kappa\}$ of all pairs of elements $U, V \in \mathbb{B}$ such that $\bar{U} \subseteq V$.

By induction on $\alpha < \kappa$ construct pairwise disjoint families

$$\{\mathbb{B}_{0,\alpha}, \mathbb{B}_{1,\alpha} : \alpha < \kappa\} \subseteq [\mathbb{B}]^{\leq L(X)}$$

such that

$$(2.3) \quad \overline{U_\alpha} \subseteq \bigcup \mathbb{B}_{i,\alpha} \subseteq V_\alpha \text{ for every } i < 2.$$

Since the cardinality of the family $\mathbb{B}_{<\alpha} = \bigcup \{\mathbb{B}_{i,\beta} : \beta < \alpha, i < 2\}$ is at most $L(X) \cdot |\alpha|$ and $L(X) \cdot |\alpha| < \min\{\chi(x, X) : x \in X\}$, the family $\mathbb{B}_{<\alpha}$ cannot contain a neighborhood base at any point $x \in X$.

Thus, by Proposition 2.5, $\mathbb{B} \setminus \mathbb{B}_{<\alpha}$ is still a base for X for every $\alpha < \kappa$. It follows that the induction can be carried out as we can select disjoint $\mathbb{B}_{\alpha,0}$ and $\mathbb{B}_{\alpha,1}$ from $[\mathbb{B} \setminus \mathbb{B}_{<\alpha}]^{\leq L(X)}$ so that

$$\overline{U_\alpha} \subseteq \bigcup \mathbb{B}_{\alpha,i} \subseteq V_\alpha$$

for $i < 2$.

Thus the disjoint families $\mathbb{B}_i = \bigcup \{\mathbb{B}_{i,\alpha} : \alpha < \kappa\}$ form a base for X by property (2.3) above and Observation 2.6(2); thus X is base resolvable. \square

We end this section by a simple observation. Recall that a space X is *weakly separated* if there is a neighborhood assignment $\{U_x : x \in X\}$ (meaning that U_x is a neighbourhood of x) so that $x \neq y \in X$ implies that $x \notin U_y$ or $y \notin U_x$. Note that left or right separated spaces are weakly separated as well as the Sorgenfrey line.

Observation 2.9. *Every weakly separated space is base resolvable.*

PROOF: Recall that every neighborhood base at some point x can be partitioned into two neighbourhood bases by Proposition 2.5(1). Thus, if \mathbb{B} is a base of X and there is a disjoint family $\{\mathbb{B}_x : x \in X\}$ of subsets of \mathbb{B} such that \mathbb{B}_x is a neighbourhood base at x for any $x \in X$ then by partitioning \mathbb{B}_x for each $x \in X$ into two neighbourhood bases of x we get a partition of \mathbb{B} into two bases of X .

Now, let us fix a base \mathbb{B} we wish to partition and a neighbourhood assignment $\{U_x : x \in X\}$ witnessing that X is weakly separated. Define

$$\mathbb{B}_x = \{U \in \mathbb{B} : x \in U \subset U_x\}$$

for $x \in X$; clearly, \mathbb{B}_x is neighbourhood base at x . Furthermore, if $x \neq y$ and say $x \notin U_y$ then $U \in \mathbb{B}_x$ implies $U \not\subset U_y$; that is, $\mathbb{B}_x \cap \mathbb{B}_y = \emptyset$ if $x \neq y \in X$ which finishes the proof. \square

We thank the referee for pointing out this last observation to us.

3. Lindelöf spaces are base resolvable

Our aim in this section is to prove that T_3 Lindelöf spaces are base resolvable. We start with a definition and some observations while the most important part of the work is done in the proof of Lemma 3.3.

Definition 3.1. Let \mathcal{A}, \mathcal{B} be families of open sets in a space X . We say that \mathcal{A} **weakly fills** \mathcal{B} if for every $U, V \in \mathcal{B}$ such that $\overline{U} \subset V$ there is $\mathcal{W} \subseteq \mathcal{A}$ such that

$$\overline{U} \subseteq \bigcup \mathcal{W} \subset V.$$

$(\mathcal{A}, \mathcal{B})$ is called a **weakly good pair** if \mathcal{A}, \mathcal{B} are disjoint, \mathcal{A} weakly fills \mathcal{B} and \mathcal{B} weakly fills \mathcal{A} .

We remark that in the next section we introduce stronger notions called *filling* and *good pairs*. The first part of the following observation basically restates Observation 2.6(2) with our new terminology:

Observation 3.2. *Suppose that X is a regular space.*

- (1) *If $(\mathcal{A}, \mathcal{B})$ is a weakly good pair in X then \mathcal{A} contains a neighborhood base at x if and only if \mathcal{B} contains a neighborhood base at x , for any $x \in X$.*
- (2) *If $\{\mathcal{A}_\alpha : \alpha < \kappa\}$ and $\{\mathcal{B}_\alpha : \alpha < \kappa\}$ are increasing chains and $(\mathcal{A}_\alpha, \mathcal{B}_\alpha)$ is a weakly good pair in X then $(\bigcup_{\alpha < \kappa} \mathcal{A}_\alpha, \bigcup_{\alpha < \kappa} \mathcal{B}_\alpha)$ is a weakly good pair as well.*

We say that the weakly good pair $(\mathcal{A}', \mathcal{B}')$ **extends** the weakly good pair $(\mathcal{A}, \mathcal{B})$ if $\mathcal{A} \subseteq \mathcal{A}'$ and $\mathcal{B} \subseteq \mathcal{B}'$. A family of pairs $\{(\mathcal{A}_\xi, \mathcal{B}_\xi) : \xi < \Theta\}$ is **pairwise disjoint** if $\mathcal{A}_\xi \cap \mathcal{B}_\zeta = \emptyset$ for each $\xi, \zeta < \Theta$.

Next, we prove that weakly good pairs can be nicely extended in Lindelöf spaces.

Lemma 3.3. *Suppose that X is a T_3 Lindelöf space with a base \mathbb{B} . Given a weakly good pair $(\mathcal{A}, \mathcal{B})$ from elements of \mathbb{B} and a single pair of open sets $\{U, V\}$ such that $\overline{U} \subset V$ there is a weakly good pair $(\mathcal{A}', \mathcal{B}')$ formed by elements of \mathbb{B} extending $(\mathcal{A}, \mathcal{B})$ such that both \mathcal{A}' and \mathcal{B}' weakly fills $\{U, V\}$.*

PROOF: We will show this essentially by induction on the size of \mathcal{A} and \mathcal{B} , however we need to prove something significantly stronger (and more technical) than the statement of the lemma itself.

Let Δ_κ stand for the following statement: For each pairwise disjoint family of weakly good pairs $\{(\mathcal{A}_i, \mathcal{B}_i), (\mathcal{C}_j, \mathcal{D}_j) : i < n, j < k\}$, each a subfamily from \mathbb{B} such that $|\mathcal{A}_i|, |\mathcal{B}_i| \leq \kappa$ and an arbitrary family of open sets \mathcal{E} of size at most κ there is a weakly good pair $(\mathcal{A}, \mathcal{B})$ from \mathbb{B} of size at most κ such that

- (1) $\bigcup_{i < n} \mathcal{A}_i \subset \mathcal{A}$ and $\bigcup_{i < n} \mathcal{B}_i \subset \mathcal{B}$,
- (2) \mathcal{A} and \mathcal{B} weakly fill \mathcal{E} ,
- (3) $\{(\mathcal{A}, \mathcal{B}), (\mathcal{C}_j, \mathcal{D}_j) : j < k\}$ is still pairwise disjoint.

We prove that Δ_κ holds for every infinite κ by induction on κ .

Claim 3.4. Δ_ω holds.

PROOF: Fix $\{(\mathcal{A}_i, \mathcal{B}_i), (\mathcal{C}_j, \mathcal{D}_j) : i < n, j < k\}$ and \mathcal{E} as above. By induction on $m \in \omega$ we build increasing chains $\{\mathcal{A}^m : m \in \omega\}$ and $\{\mathcal{B}^m : m \in \omega\}$ from subsets of \mathbb{B} such that

- (1) $\mathcal{A}^0 = \bigcup_{i < n} \mathcal{A}_i, \mathcal{B}^0 = \bigcup_{i < n} \mathcal{B}_i,$
- (2) $\mathcal{A}^{m+1} \setminus \mathcal{A}^m$ and $\mathcal{B}^{m+1} \setminus \mathcal{B}^m$ are countable well-founded families,
- (3) the family of pairs $\{(\mathcal{A}^m, \mathcal{B}^m), (\mathcal{C}_j, \mathcal{D}_j) : j < k\}$ is pairwise disjoint

for each $m \in \omega$. Furthermore, we want to make sure that $\mathcal{A} = \bigcup_{m \in \omega} \mathcal{A}^m$ and $\mathcal{B} = \bigcup_{m \in \omega} \mathcal{B}^m$ form a weakly good pair and they both weakly fill \mathcal{E} . Therefore, we partition ω into infinite sets $\omega = \bigcup \{D_m : m \in \omega\}$ and at the m^{th} step

- (4) we fix a surjective map

$$f_m : D_m \setminus (m + 1) \rightarrow \{(U, V) \in (\mathcal{A}^m \cup \mathcal{B}^m \cup \mathcal{E})^2 : \overline{U} \subset V\};$$

- (5) if $m \in D_\ell \setminus (\ell + 1)$ and $f_\ell(m) = (U, V)$ then both \mathcal{A}^{m+1} and \mathcal{B}^{m+1} weakly fill $\{U, V\}$.

In particular, it suffices to construct disjoint \mathcal{A}^{m+1} and \mathcal{B}^{m+1} from \mathcal{A}^m and \mathcal{B}^m such that they satisfy (2), (3) and (5) above, especially they both weakly fill a given (U, V) . We construct \mathcal{A}^{m+1} , the proof for \mathcal{B}^{m+1} is analogous.

Subclaim 3.4.1. $\mathbb{B} \setminus (\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j)$ is a base of X .

PROOF OF THE SUBCLAIM: Let $x \in X$ be arbitrary.

If $\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j$ does not contain a neighborhood base at x , then $\mathbb{B} \setminus (\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j)$ should contain a neighborhood base at x by Proposition 2.5(3).

Assume now that $\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j$ contains a neighborhood base at x . Since

$$\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j = (\mathcal{B}^m \setminus \mathcal{B}^0) \cup \bigcup_{i < n} \mathcal{B}_i \cup \bigcup_{j < k} \mathcal{D}_j,$$

applying Proposition 2.5(3) again, one of the sets

$$(3.1) \quad \mathcal{B}^m \setminus \mathcal{B}^0, \mathcal{B}_0, \dots, \mathcal{B}_{n-1}, \mathcal{D}_0, \dots, \mathcal{D}_{k-1}$$

contains a neighborhood base at x . Since $\mathcal{B}^m \setminus \mathcal{B}^0$ is well-founded, it cannot contain a neighborhood base. If \mathcal{B}_i (or \mathcal{D}_j , respectively) contains a neighborhood base at x , then \mathcal{A}_i (or \mathcal{C}_j , respectively) also contains a neighborhood base at x by Observation 3.2(1). In both cases, $\mathbb{B} \setminus (\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j)$ contains a neighborhood base, which proves the Subclaim. \square

Since X is Lindelöf, using the Subclaim above and Proposition 2.2 we can find a countable well-founded cover $\mathcal{Q} \subset \mathbb{B} \setminus (\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j)$ of \overline{U} with $\bigcup \mathcal{Q} \subset V$. Now define $\mathcal{A}^{m+1} = \mathcal{A}^m \cup \mathcal{Q}$. Since \mathcal{Q} and $(\mathcal{B}^m \cup \bigcup_{j < k} \mathcal{D}_j)$ are disjoint, (3) holds. (2) and (5) are clear from the construction. \square

Claim 3.5. Suppose that Δ_λ holds for every $\omega \leq \lambda < \kappa$. Then Δ_κ holds.

PROOF: Fix $\{(\mathcal{A}_i, \mathcal{B}_i), (\mathcal{C}_j, \mathcal{D}_j) : i < n, j < k\}$ and \mathcal{E} , let $\text{cf}(\kappa) = \mu$ and fix a cofinal sequence of ordinals $(\kappa_\xi)_{\xi < \mu}$ in κ . Take a chain of elementary submodels $(M_\xi)_{\xi < \mu}$ of $H(\theta)$ (where θ is large enough) such that everything relevant is in M_0 , $\kappa_\xi \subset M_\xi$ and $|M_\xi| = |\kappa_\xi|$ for $\xi < \mu$. The following is an easy consequence of M_ξ being elementary and X being Lindelöf:

Subclaim 3.5.1. $(\mathcal{A}_i \cap M_\xi, \mathcal{B}_i \cap M_\xi)$ is a weakly good pair and $|\mathcal{A}_i \cap M_\xi|, |\mathcal{B}_i \cap M_\xi| \leq |\kappa_\xi|$ for all $i < n$.

PROOF OF THE SUBCLAIM: If $U, V \in \mathcal{A}_i \cap M_\xi, \bar{U} \subset V$ then $\mathcal{A}_i, \mathcal{B}_i, A \in M_\xi$ implies that

$$M_\xi \models \exists \mathcal{B} \in [\mathcal{B}_i]^\omega \quad \bar{U} \subset \bigcup \mathcal{B} \subset V$$

because X is Lindelöf. So there is $\mathcal{B} \in M_\xi \cap [\mathcal{B}_i]^\omega$ such that $\bar{U} \subset \bigcup \mathcal{B} \subset U$. Since \mathcal{B} is countable, $\mathcal{B} \in M_\xi$ implies $\mathcal{B} \subset M_\xi$. So we have $\mathcal{B} \subset \mathcal{B}_i \cap M_\xi$ with $\bar{U} \subset \bigcup \mathcal{B} \subset V$. This shows that $\mathcal{B}_i \cap M_\xi$ fills $\mathcal{A}_i \cap M_\xi$ and the other direction of the proof is completely analogous. \square

By induction on $\xi < \mu$ construct weakly good pairs $\{(\mathcal{A}^\xi, \mathcal{B}^\xi) : \xi < \mu\}$ so that $\mathcal{A}^\xi \subset \mathcal{A}^\zeta, \mathcal{B}^\xi \subset \mathcal{B}^\zeta$ for $\xi < \zeta < \mu$ and

- (i) $\bigcup_{i < n} (\mathcal{A}_i \cap M_\xi) \subset \mathcal{A}^\xi \subset \mathbb{B}$ and $\bigcup_{i < n} (\mathcal{B}_i \cap M_\xi) \subset \mathcal{B}^\xi \subset \mathbb{B}$,
- (ii) \mathcal{A}^ξ and \mathcal{B}^ξ has size $\leq |\kappa_\xi|$,
- (iii) \mathcal{A}^ξ and \mathcal{B}^ξ weakly fills $\mathcal{E} \cap M_\xi$,
- (iv) $\mathcal{A}^\xi \cap \mathcal{B}_i = \emptyset, \mathcal{A}^\xi \cap \mathcal{D}_j = \emptyset$ and $\mathcal{B}^\xi \cap \mathcal{A}_i = \emptyset, \mathcal{B}^\xi \cap \mathcal{C}_j = \emptyset$.

This can be done using $\Delta_{|\kappa_\xi|}$ at stage ξ . First note that $\mathcal{A}^{<\xi} = \bigcup \{\mathcal{A}^\zeta : \zeta < \xi\}$ and $\mathcal{B}^{<\xi} = \bigcup \{\mathcal{B}^\zeta : \zeta < \xi\}$ are of size at most $|\kappa_\xi|$ and $(\mathcal{A}^{<\xi}, \mathcal{B}^{<\xi})$ is a weakly good pair. Also, the family

$$\{(\mathcal{A}^{<\xi}, \mathcal{B}^{<\xi}), (\mathcal{A}_i \cap M_\xi, \mathcal{B}_i \cap M_\xi); (\mathcal{A}_i, \mathcal{B}_i), (\mathcal{C}_j, \mathcal{D}_j) : i < n, j < k\}$$

is pairwise disjoint. Hence $\Delta_{|\kappa_\xi|}$ implies that there is a weakly good pair $(\mathcal{A}^\xi, \mathcal{B}^\xi)$ from \mathbb{B} of size at most $|\kappa_\xi|$ which fills $\mathcal{E} \cap M_\xi$ and is pairwise disjoint from $\{(\mathcal{A}_i, \mathcal{B}_i), (\mathcal{C}_j, \mathcal{D}_j) : i < n, j < k\}$ while

$$\mathcal{A}^{<\xi} \cup \bigcup_{i < n} (\mathcal{A}_i \cap M_\xi) \subset \mathcal{A}^\xi$$

and

$$\mathcal{B}^{<\xi} \cup \bigcup_{i < n} (\mathcal{B}_i \cap M_\xi) \subset \mathcal{B}^\xi.$$

Note that $\Delta_{|\kappa_\xi|}$ was used to find the common extension of $n + 1$ weakly good pairs such that this extension is disjoint from $n + k$ given weakly good pairs. Now define $\mathcal{A} = \bigcup \{\mathcal{A}^\xi : \xi < \zeta\}$ and $\mathcal{B} = \bigcup \{\mathcal{B}^\xi : \xi < \zeta\}$. Then $(\mathcal{A}, \mathcal{B})$ is the desired extension. \square

This finishes the proof the lemma. \square

Recall that a space is *locally Lindelöf* if every point has a neighbourhood with Lindelöf closure.

Proposition 3.6. *Suppose that X is a T_3 locally Lindelöf space. Then X embeds into a T_3 Lindelöf space X^* with $|X^* \setminus X| = 1$.*

PROOF: Construct X^* on the set $X \cup \{x^*\}$ where neighborhoods of the point x^* are of the form $\{x^*\} \cup X \setminus \overline{U}$ with $U \subset X$ being open and such that there is an open $V \subset X$ with $\overline{U} \subset V$ and \overline{V} is Lindelöf. It is clear that X^* is Hausdorff and Lindelöf.

Note that if U, V are open in X , $\overline{U} \subset V$ and \overline{V} is Lindelöf, then \overline{V} is normal as well, so there is an open $W \subset V$ so that $\overline{U} \subset W \subset \overline{W} \subset V$. So X^* is regular at the point x^* , so X^* is regular. \square

Corollary 3.7. *Every T_3 locally Lindelöf space is base resolvable. In particular, every T_3 locally countable or locally compact space is base resolvable.*

PROOF: Fix a base \mathbb{B} for a T_3 Lindelöf space X and consider the set \mathbb{P} of all weakly good pairs $(\mathcal{A}, \mathcal{B})$ from \mathbb{B} partially ordered by extension. Note that we can apply Zorn’s lemma to \mathbb{P} by Observation 3.2 part (2); pick a maximal weakly good pair $(\mathcal{A}, \mathcal{B}) \in \mathbb{P}$. Lemma 3.3 implies that a maximal weakly good pair must weakly fill every pair of open sets $\{U, V\}$ with $\overline{U} \subset V$, hence both \mathcal{A} and \mathcal{B} are bases of X .

Given a T_3 locally Lindelöf space X with a base \mathbb{B} consider it’s one-point Lindelöfization $X^* = X \cup \{x^*\}$ with the base

$$\mathbb{B}^* = \mathbb{B} \cup \{U \subseteq X^* : U \text{ is an open neighbourhood of } x^* \text{ in } X^*\}.$$

X^* is T_3 Lindelöf hence base resolvable; thus \mathbb{B}^* can be partitioned into two bases, \mathbb{B}_0^* and \mathbb{B}_1^* , which clearly gives a partition of \mathbb{B} namely, $\mathbb{B}_0^* \cap \mathbb{B}$ and $\mathbb{B}_1^* \cap \mathbb{B}$. \square

4. Combinatorics of resolvability

In this section, we will prove a combinatorial lemma which will be our next tool in showing that further classes of space are base resolvable.

Definition 4.1. Let $\mathcal{A}, \mathcal{B} \subseteq \mathcal{P}(X)$. We say that \mathcal{A} **fills** \mathcal{B} if

$$U = \bigcup \{V \in \mathcal{A} : V \subsetneq U\}$$

for every $U \in \mathcal{B}$. \mathcal{A}, \mathcal{B} is called a **good pair** if \mathcal{A}, \mathcal{B} are disjoint, \mathcal{A} fills \mathcal{B} and \mathcal{B} fills \mathcal{A} . \mathcal{A} is **self-filling** if \mathcal{A} fills \mathcal{A} .

Note that if $\mathcal{A} \subseteq \mathcal{P}(X)$ fills $\{\cap \mathcal{B} : \mathcal{B} \in [\mathcal{A}]^{<\omega}\}$ and \mathcal{A} covers X then \mathcal{A} is a base for a topology on X .

Definition 4.2. A self-filling family \mathcal{A} is **resolvable** if there is a partition $\mathcal{A}_0, \mathcal{A}_1$ of \mathcal{A} such that \mathcal{A}_i fills \mathcal{A} for $i < 2$.

The importance of the following lemma is that it shows that resolvability is a local property:

Theorem 4.3. *Suppose that $\mathbb{B} \subseteq \mathcal{P}(X)$ is self-filling. Then the following are equivalent:*

(1) *for every $U \in \mathbb{B}$ there is a good pair $(\mathbb{B}_0^U, \mathbb{B}_1^U)$ from \mathbb{B} such that*

$$U = \bigcup \mathbb{B}_0^U = \bigcup \mathbb{B}_1^U,$$

(2) *\mathbb{B} is resolvable.*

PROOF: (2) implies (1) is trivial.

To see that (1) implies (2), let \mathcal{P} be the set of all good pairs $(\mathbb{B}_0, \mathbb{B}_1)$ formed by elements of \mathbb{B} . Then \mathcal{P} is partially ordered by $(\mathbb{B}_0, \mathbb{B}_1) \leq (\mathbb{B}'_0, \mathbb{B}'_1)$ if and only if $\mathbb{B}_i \subseteq \mathbb{B}'_i$ for $i < 2$. It is clear that every chain in (\mathcal{P}, \leq) has an upper bound hence, by Zorn's lemma, we can pick a \leq -maximal element $(\mathbb{B}_0, \mathbb{B}_1) \in \mathcal{P}$.

We claim that \mathbb{B}_i fills \mathbb{B} for $i < 2$. Pick any $U \in \mathbb{B}$ and consider the good pair $\mathbb{B}_0^U, \mathbb{B}_1^U$ with $U = \bigcup \mathbb{B}_0^U = \bigcup \mathbb{B}_1^U$. Define

$$\mathbb{B}'_i = \mathbb{B}_i \cup (\mathbb{B}_i^U \setminus \mathbb{B}_{1-i})$$

for $i < 2$.

The second statement of the following lemma yields immediately that $(\mathbb{B}'_0, \mathbb{B}'_1)$ forms a good pair which fills $\{U\}$.

Lemma 4.4. (1) *If a family of sets \mathcal{A} fills a family of sets \mathcal{B} and \mathcal{A}' fills \mathcal{B}' then $\mathcal{A} \cup (\mathcal{A}' \setminus \mathcal{B})$ fills \mathcal{B}' .*

(2) *If $(\mathcal{A}, \mathcal{B})$ and $(\mathcal{A}', \mathcal{B}')$ are good pairs then $(\mathcal{A} \cup (\mathcal{A}' \setminus \mathcal{B}), \mathcal{B} \cup (\mathcal{B}' \setminus \mathcal{A}))$ is also a good pair which fills $\bigcup \mathcal{B}'$.*

PROOF OF THE LEMMA: (1) Pick $U \in \mathcal{B}'$. Since \mathcal{A}' fills \mathcal{B}' , there is $\mathcal{A}^+ \subset \mathcal{A}' \setminus \{U\}$ with $U = \bigcup \mathcal{A}^+$. For each $B \in \mathcal{A}^+ \cap \mathcal{B}$ choose $\mathcal{A}_B \subset \mathcal{A}$ with $B = \bigcup \mathcal{A}_B$. Finally let

$$\mathcal{A}^* = (\mathcal{A}^+ \setminus \mathcal{B}) \cup \bigcup \{\mathcal{A}_B : B \in \mathcal{A}^+ \cap \mathcal{B}\}.$$

Then $\mathcal{A}^* \subset \mathcal{A} \cup (\mathcal{A}' \setminus \mathcal{B}) \setminus \{U\}$ and

$$\begin{aligned} \bigcup \mathcal{A}^* &= \bigcup \left((\mathcal{A}^+ \setminus \mathcal{B}) \cup \bigcup \{\mathcal{A}_B : B \in \mathcal{A}^+ \cap \mathcal{B}\} \right) \\ &= \bigcup \left((\mathcal{A}^+ \setminus \mathcal{B}) \cup \{B : B \in \mathcal{A}^+ \cap \mathcal{B}\} \right) = \bigcup \mathcal{A}^+ = U. \end{aligned}$$

(2) The families $\mathcal{A} \cup (\mathcal{A}' \setminus \mathcal{B})$ and $\mathcal{B} \cup (\mathcal{B}' \setminus \mathcal{A})$ are clearly disjoint, $\mathcal{A} \cup (\mathcal{A}' \setminus \mathcal{B})$ fills $\mathcal{B} \cup (\mathcal{B}' \setminus \mathcal{A}) \cup \{\bigcup \mathcal{A}\}$ and $\mathcal{B} \cup (\mathcal{B}' \setminus \mathcal{A})$ fills $\mathcal{A} \cup (\mathcal{A}' \setminus \mathcal{B}) \cup \{\bigcup \mathcal{B}\}$ by (1) which was to be proved. □

Also, $(\mathbb{B}_0, \mathbb{B}_1) \leq (\mathbb{B}'_0, \mathbb{B}'_1)$ and thus by the maximality of $(\mathbb{B}_0, \mathbb{B}_1)$ we have that $\mathbb{B}'_i = \mathbb{B}_i$. This finishes the proof. □

The first corollary is a direct application and shows that resolvability is preserved by unions.

Corollary 4.5. *Suppose that \mathbb{B}_α is a resolvable self-filling family for each $\alpha < \kappa$. Then $\bigcup\{\mathbb{B}_\alpha : \alpha < \kappa\}$ is a resolvable self-filling family as well.*

Corollary 4.6. *Suppose that a self-filling family \mathbb{B} has the property that*

(†) *for every $U \in \mathbb{B}$ there is $\mathcal{U} \in [\mathbb{B} \setminus \{U\}]^{\leq \omega}$ such that $U = \bigcup \mathcal{U}$.*

Then \mathbb{B} is resolvable.

PROOF: We need the following Claim.

Claim 4.7. *If $\mathcal{A} \subset \mathbb{B}$ is well-founded then for every $W \in \mathbb{B}$ there is a countable well-founded family $\mathcal{B}(W, \mathcal{A}) \subset \mathbb{B} \setminus \mathcal{A}$ with $\bigcup \mathcal{B}(W, \mathcal{A}) = W$.*

PROOF: We can assume that $W \in \mathcal{A}$. By (†) there is a countable self-filling family $\mathcal{C} \subset \mathbb{B}$ with $W \in \mathcal{C}$. Let

$$\mathcal{V} = \{V \in \mathcal{C} \setminus \mathcal{A} : V \subsetneq W\}.$$

Since \mathcal{A} is well-founded, for each $x \in W$ the family $\{Z \in \mathcal{A} \cap \mathcal{C} : x \in Z\}$ has a \subset -minimal element Z . Since \mathcal{C} is self-filling, there is $V \in \mathcal{C}$ with $x \in V \subsetneq Z$. Then $V \in \mathcal{V}$.

Thus $\bigcup \mathcal{V} = W$. Now, by Proposition 2.2, there is a well-founded family $\mathcal{B}(W, \mathcal{A}) \subset \mathcal{V}$ with $\bigcup \mathcal{B}(W, \mathcal{A}) = W$. □

By Theorem 4.3, it suffices to prove that for every $U \in \mathbb{B}$ there is a good pair $(\mathbb{B}_0, \mathbb{B}_1)$ from \mathbb{B} such that $U = \bigcup \mathbb{B}_0 = \bigcup \mathbb{B}_1$.

Fix a $U \in \mathbb{B}$. Partition ω into infinite sets $\omega = \bigcup\{D_m : m \in \omega\}$. By induction on $m \in \omega$ we build increasing chains $\{\mathbb{B}_0^m : m \in \omega\}$ and $\{\mathbb{B}_1^m : m \in \omega\}$ from subsets of \mathbb{B} such that

- (1) $\mathbb{B}_0^0 = \mathbb{B}_1^0 = \emptyset$,
- (2) \mathbb{B}_0^m and \mathbb{B}_1^m are disjoint, well founded and countable families,
- (3) fix a surjective map

$$f_m : D_m \setminus (m + 1) \twoheadrightarrow \{U\} \cup B_0^m \cup B_1^m,$$

- (4) if $m \in D_\ell$ and $f_\ell(m) = V$ then

$$(4.1) \quad \mathbb{B}_0^{m+1} = \mathbb{B}_0^m \cup \mathcal{B}(V, \mathbb{B}_1^m)$$

and

$$(4.2) \quad \mathbb{B}_1^{m+1} = \mathbb{B}_1^m \cup \mathcal{B}(V, \mathbb{B}_0^{m+1}).$$

Let $\mathbb{B}_i = \bigcup\{\mathbb{B}_i^m : m \in \omega\}$ for $i < 2$. The $(\mathbb{B}_0, \mathbb{B}_1)$ is a good pair and $U = \bigcup \mathbb{B}_0 = \bigcup \mathbb{B}_1$. Indeed, if $V \in \mathbb{B}_i \cup \{U\}$ then $V \in \mathbb{B}_i^m \cup \{U\}$ for some $m \in \omega$ and so $f_m(\ell) = V$ for some $\ell \in D_m \setminus (m + 1)$. Thus there is a family $\mathcal{B} \subset \mathbb{B}_{1-i}^{\ell+1} \subset \mathbb{B}_{1-i}$ with $\bigcup \mathcal{B} = V$. □

Corollary 4.8. *Locally countable or hereditarily Lindelöf spaces are base resolvable without assuming any separation axioms.*

Our next corollary establishes that every reasonable space admits a resolvable base.

Corollary 4.9. *Suppose that \mathbb{B} is a base closed under finite unions in a T_1 topological space. Then \mathbb{B} is resolvable.*

PROOF: We apply Theorem 4.3 again. Fix $U \in \mathbb{B}$ and we construct a good pair covering U . Fix an arbitrary strictly decreasing sequence $\{U_n : n \in \omega\} \subseteq \mathbb{B}$ such that $U_0 \subseteq U$ and fix $y_n \in U_{n-1} \setminus U_n$ for $n \in \omega \setminus \{0\}$. Let

$$\mathbb{B}_i^U = \{V \in \mathbb{B} \cap \mathcal{P}(U) : \exists k \in \omega \setminus \{0\} : U_{2k+i} \subseteq V \text{ but } U_{2k-1+i} \not\subseteq V\}$$

for $i < 2$. It should be clear that $\mathbb{B}_0^U \cap \mathbb{B}_1^U = \emptyset$.

Next we prove that $U = \bigcup \mathbb{B}_i^U$ for $i < 2$. Fix $i < 2$ and note that $\{U_{2k+i} : k \in \omega \setminus \{0\}\} \subset \mathbb{B}_i^U$. Now fix $x \in U$ and we prove that $x \in \bigcup \mathbb{B}_i^U$; without loss of generality we can suppose that $x \notin U_{2+i}$. Find any $k \in \omega$ so that $y_{2k+i} \neq x$ and take $W \in \mathbb{B}$ so that $x \in W \subset U \setminus \{y_{2k+i}\}$; here we used that \mathbb{B} is a base of a T_1 topology. Note that $V = U_{2k+i} \cup W \in \mathbb{B}$ as \mathbb{B} is closed under finite unions and that $x \in V \in \mathbb{B}_i^U$.

Finally we show that $(\mathbb{B}_0^U, \mathbb{B}_1^U)$ is a good pair; we will show that \mathbb{B}_0^U fills \mathbb{B}_1^U , the other direction is completely analogous. Fix $V \in \mathbb{B}_1^U$ and fix a point $z \in V$. Find an $l \in \omega$ so that $U_{2l-1} \subset V$ and $z \neq y_{2l}$. As \mathbb{B} is a base, there is $W \in \mathbb{B}$ so that $z \in W \subset V \setminus \{y_{2l}\}$. Let $V' = U_{2l} \cup W$. As \mathbb{B} is closed under finite unions we have $V' \in \mathbb{B}$. Moreover $V' \in \mathbb{B}_0^U$ as witnessed by $U_{2l} \subset V'$ but $U_{2l-1} \not\subseteq V'$. Finally, $z \in V' \subset V$ as we wanted. \square

Corollary 4.10. *The set of all open sets in a T_1 topological space is resolvable.*

Let $MA(\text{Cohen})$ denote Martin’s axiom restricted to the partial orderings of the form $Fn(\kappa, 2, \omega)$ for some κ where, $Fn(\kappa, 2, \omega)$ is the poset of functions from some finite subset of κ to 2 ordered by reverse inclusion.

Corollary 4.11. *Under $MA(\text{Cohen})$ every space X of local size $< 2^\omega$ is base resolvable without assuming any separation axioms.*

PROOF: Fix a base \mathbb{B} of X ; we may assume that $|U| < 2^\omega$ for all $U \in \mathbb{B}$. We apply Theorem 4.3 to prove that \mathbb{B} is resolvable as a self-filling family which in turn will imply that \mathbb{B} is a resolvable base. Fix $U \in \mathbb{B}$ and we construct a good pair covering U . Let $\kappa = |U|$ and select $\mathbb{B}_U \in [\mathbb{B}]^\kappa$ which fills itself and $\bigcup \mathbb{B}_U = U$. Now consider the ccc partial order $\mathbb{P} = Fn(\mathbb{B}_U, 2, \omega)$, i.e., the set of all finite partial functions from \mathbb{B}_U to 2. Now consider

$$D_{x,V,i} = \{f \in \mathbb{P} : \text{there is } W \in f^{-1}(i) : x \in W \subset V\}$$

for $i < 2$ and $x \in V \in \mathbb{B}_U$; note that each $D_{x,V,i}$ is dense in \mathbb{P} . Hence there is a filter $G \subseteq \mathbb{P}$ which intersects $D_{x,V,i}$ for $i < 2$ and $x \in V \in \mathbb{B}_U$. Let $\mathbb{B}_i = \{V \in \mathbb{B}_U : (\bigcup G)(V) = i\}$ for $i < 2$ and note that $(\mathbb{B}_0, \mathbb{B}_1)$ is the desired good pair. \square

5. Thinning self-filling families

Let \mathbb{B} be a self-filling family. Note that \mathbb{B} is *redundant* in the sense that $\mathbb{B} \setminus \mathcal{U}$ still fills \mathbb{B} for a finite or more generally, a well founded family \mathcal{U} .

Definition 5.1. We say that $\mathcal{U} \subseteq \mathbb{B}$ is *negligible* if $\mathbb{B} \setminus \mathcal{U}$ still fills \mathbb{B} .

Our aim in this section is to show that every self-filling family \mathbb{B} contains a negligible subfamily of size $|\mathbb{B}|$. Note that a base \mathbb{B} for a space X is *resolvable* if and only if it contains a negligible subfamily $\mathcal{U} \subseteq \mathbb{B}$ such that \mathcal{U} is a base of X as well. We will make use of the following definitions:

Definition 5.2. If \mathbb{B} fills itself then let

$$L(U, \mathbb{B}) = \min\{|\mathcal{V}| : \mathcal{V} \subseteq \mathbb{B} \setminus \{U\}, U = \bigcup \mathcal{V}\}$$

for $U \in \mathbb{B}$.

Observation 5.3. Suppose that \mathbb{B} fills itself and $\mathcal{U} \subseteq \mathbb{B}$.

- (1) If $\mathbb{B} \setminus \mathcal{U}$ fills \mathcal{U} then \mathcal{U} is negligible.
- (2) If \mathcal{U} is well founded then $\mathbb{B} \setminus \mathcal{U}$ fills \mathcal{U} and so \mathcal{U} is negligible; in particular, if \mathcal{U} is weakly increasing, then \mathcal{U} is negligible.

Our first proposition establishes the main result for self-filling families \mathbb{B} with $cf|\mathbb{B}| = |\mathbb{B}|$.

Proposition 5.4. Suppose that \mathbb{B} fills itself and $\kappa = |\mathbb{B}|$ is regular. Then \mathbb{B} contains a negligible family of size κ .

PROOF: We can suppose that $L(U, \mathbb{B}) < \kappa$ for every $U \in \mathbb{B}$; otherwise we can find a weakly increasing subfamily of size κ which is negligible by (1) and (2) of Observation 5.3.

It suffices to define a sequence $\mathcal{U}_\xi, \mathcal{V}_\xi \in [\mathbb{B}]^{<\kappa}$ for $\xi < \kappa$ such that

- (1) $\mathcal{U}_\xi \cap \mathcal{V}_\xi = \emptyset$,
- (2) $\mathcal{U}_\xi \subset \mathcal{U}_\zeta$ and $\mathcal{V}_\xi \subset \mathcal{V}_\zeta$ for $\xi < \zeta < \kappa$,
- (3) \mathcal{V}_ξ fills \mathcal{U}_ξ , and
- (4) $\mathcal{U}_{\xi+1} \setminus \mathcal{U}_\xi \neq \emptyset$.

Clearly, $\mathcal{U} = \bigcup \{\mathcal{U}_\xi : \xi < \kappa\}$ will be a negligible set of size κ in \mathbb{B} by (3) of Observation 5.3. Suppose we have $\mathcal{U}_\xi, \mathcal{V}_\xi \in [\mathbb{B}]^{<\kappa}$ for $\xi < \zeta$ as above for some $\zeta < \kappa$; then $\mathbb{B} \setminus \bigcup \{\mathcal{U}_\xi, \mathcal{V}_\xi : \xi < \zeta\} \neq \emptyset$ by κ being regular. Hence we can select $U_\zeta \in \mathbb{B} \setminus \bigcup \{\mathcal{U}_\xi, \mathcal{V}_\xi : \xi < \zeta\}$ and define

$$\mathcal{U}_\zeta = \bigcup \{\mathcal{U}_\xi : \xi < \zeta\} \cup \{U_\zeta\}.$$

Find $\mathcal{W} \subseteq \mathbb{B} \setminus \{U_\zeta\}$ of size $< \kappa$ such $\bigcup \mathcal{W} = U_\zeta$; define

$$\mathcal{V}_\zeta = \bigcup \{\mathcal{V}_\xi : \xi < \zeta\} \cup (\mathcal{W} \setminus \mathcal{U}_\zeta).$$

Since $\bigcup \{\mathcal{V}_\xi : \xi < \zeta\}$ fills $\bigcup \{\mathcal{U}_\xi : \xi < \zeta\}$ by the inductive hypothesis (3) above, Lemma 4.4(1) implies that \mathcal{V}_ζ fills \mathcal{U}_ζ . □

Theorem 5.5. *Suppose that \mathbb{B} fills itself. Then \mathbb{B} contains a negligible family of size $|\mathbb{B}|$.*

PROOF: We can suppose that $\mu = \text{cf}(\kappa) < \kappa = |\mathbb{B}|$ and that every weakly increasing sequence in \mathbb{B} is of size less than κ by Observation 5.3(2). Fix a cofinal strictly increasing sequence of regular cardinals $(\kappa_\xi)_{\xi < \mu}$ in κ such that $\mu < \kappa_0$ and define

$$\mathbb{B}_\xi = \{U \in \mathbb{B} : L(U, \mathbb{B}) \leq \kappa_\xi\}$$

for every $\xi < \mu$. So we have

$$(5.1) \quad \mathbb{B} = \bigcup_{\xi < \mu} \mathbb{B}_\xi.$$

If there is a $\xi < \mu$ such that every weakly increasing sequence in \mathbb{B} is of size less than κ_ξ then $\mathbb{B} = \mathbb{B}_\xi$. Let us define a set mapping $F : \mathbb{B} \rightarrow [\mathbb{B}]^{<\kappa_\xi^+}$ such that $U = \bigcup F(U)$ where $F(U) \subseteq \mathbb{B} \setminus \{U\}$. As $\kappa_\xi^+ < \kappa$ we can apply Hajnal's Set Mapping theorem (see Theorem 19.2 in [1]) and so there is an F -free set \mathcal{U} of size κ in \mathbb{B} , i.e., $F(U) \cap \mathcal{U} = \emptyset$ for all $U \in \mathcal{U}$. Observe that \mathcal{U} is negligible as $\bigcup \{F(U) : U \in \mathcal{U}\} \subseteq \mathbb{B} \setminus \mathcal{U}$ fills \mathcal{U} .

Now we suppose that $\mathbb{B} \neq \mathbb{B}_\xi$ for $\xi < \mu$, that is, there is a weakly increasing sequence in \mathbb{B} of size κ_ξ for all $\xi < \mu$. It suffices to define sequences $\mathcal{U}_\xi, \mathcal{V}_\xi \in [\mathbb{B}]^{<\kappa}$ for $\xi < \mu$ such that

- (i) $\mathcal{U}_\xi \subset \mathcal{U}_\zeta$ and $\mathcal{V}_\xi \subset \mathcal{V}_\zeta$ for $\xi < \zeta < \mu$,
- (ii) $\mathcal{U}_\xi, \mathcal{V}_\xi$ are disjoint and $\kappa_\xi \leq |\mathcal{U}_\xi|$,
- (iii) \mathcal{V}_ξ fills \mathcal{U}_ξ .

Indeed, the union $\bigcup \{\mathcal{U}_\xi : \xi < \mu\}$ is negligible in \mathbb{B} of size κ by Observation 5.3(1) because $\bigcup \{\mathcal{V}_\xi : \xi < \mu\}$ fills $\bigcup \{\mathcal{U}_\xi : \xi < \mu\}$.

Suppose we defined $\mathcal{U}_\xi, \mathcal{V}_\xi \in [\mathbb{B}]^{<\kappa}$ for $\xi < \mu$ and let

$$\lambda = \left(\left| \bigcup \{\mathcal{U}_\xi \cup \mathcal{V}_\xi : \xi < \mu\} \right| \cdot \kappa_\mu \right)^+.$$

Note that $\lambda < \kappa$ thus we can pick a weakly increasing family $\mathcal{W} \in [\mathbb{B}]^\lambda$. Without loss of generality, we can suppose that \mathcal{W} is disjoint from $\bigcup \{\mathcal{U}_\xi \cup \mathcal{V}_\xi : \xi < \mu\}$. Note that

$$\mathcal{W} = \bigcup \{\mathbb{B}_\delta \cap \mathcal{W} : \delta < \mu\}$$

by (5.1), and that $\mu < \text{cf}(\lambda) = \lambda$, hence there is $\delta < \mu$ such that $\mathcal{W}' = \mathcal{W} \cap \mathbb{B}_\delta$ has size λ . Define

$$\mathcal{U}_\zeta = \bigcup \{\mathcal{U}_\xi : \xi < \zeta\} \cup \mathcal{W}'.$$

Now, for every $U \in \mathcal{W}'$ select $F(U) \in [\mathbb{B} \setminus \{U\}]^{\leq \kappa_\delta}$ such that $U = \bigcup F(U)$. Define

$$\mathcal{V}_\zeta = \bigcup \{\mathcal{V}_\xi : \xi < \zeta\} \cup \bigcup \{F(U) : U \in \mathcal{W}'\} \setminus \mathcal{U}_\zeta.$$

Note that $\kappa_\zeta \leq |\mathcal{U}_\zeta| = \lambda$ and $|\mathcal{V}_\zeta| \leq \lambda \cdot \kappa_\delta < \kappa$. It is only left to prove that \mathcal{V}_ζ fills \mathcal{U}_ζ ; in fact, it suffices to show that \mathcal{V}_ζ fills \mathcal{W}' . Suppose that \prec is the well ordering witnessing that \mathcal{W}' is weakly increasing and suppose that there is a $U \in \mathcal{W}'$ which is not filled by \mathcal{V}_ζ ; we can suppose that U is \prec -minimal. Fix an $x \in U$ witnessing that \mathcal{V}_ζ does not fill U . Pick $V \in F(U)$ such that $x \in V \subset U$; then $V \notin \mathcal{V}_\zeta$, so $V \in \mathcal{W}'$ or $V \in \bigcup\{\mathcal{U}_\xi : \xi < \zeta\}$. If $V \in \mathcal{W}'$ then $V \prec U$, thus V is filled by \mathcal{V}_ζ by the minimality of U . This contradicts the choice of x , hence $V \notin \mathcal{W}'$. Thus $V \in \bigcup\{\mathcal{U}_\xi : \xi < \zeta\}$ which is filled by $\bigcup\{\mathcal{V}_\xi : \xi < \zeta\} \subset \mathcal{V}_\zeta$ by the inductual hypothesis. This again contradicts the choice of x , which finishes the proof. \square

6. Irresolvable self-filling families

The aim of this section is to construct an irresolvable self-filling family and deduce the existence of a non base resolvable T_0 topological space.

Given a partial order (\mathbb{P}, \leq) and $p, q \in \mathbb{P}$ let

$$[p, q] = \{r \in \mathbb{P} : p \leq r \leq q\}.$$

The key to our construction is the following special partition relation:

Definition 6.1. We say that a poset \mathbb{P} without maximal elements satisfies

$$\mathbb{P} \rightarrow (I_\omega)_2^1$$

if for every partition $\mathbb{P} = D_0 \cup D_1$ there is $i < 2$ and strictly increasing $\{p_n : n \in \omega\} \subseteq D_i$ such that $[p_0, p_n] \subseteq D_i$ for every $n \in \omega$. The negation is denoted by $\mathbb{P} \not\rightarrow (I_\omega)_2^1$.

The above definition is motivated by the following:

Observation 6.2. For any irresolvable self-filling family $\mathbb{B} \subseteq \mathcal{P}(X)$ the partial order $\mathbb{P} = (\mathbb{B}, \supseteq)$ satisfies $\mathbb{P} \rightarrow (I_\omega)_2^1$.

PROOF: Consider a partition of $\mathbb{P} = (\mathbb{B}, \supseteq)$ into sets D_0, D_1 . As \mathbb{B} is irresolvable, there is $i < 2$, $x \in X$ and $U \in D_i$ such that $V \in D_i$ for every $V \in \mathbb{B}$ with $x \in V \subseteq U$. Pick a strictly decreasing sequence $\{V_n : n \in \omega\} \subseteq \mathbb{B}$ such that $x \in V_n \subseteq U$ for every $n \in \omega$; clearly, $[V_0, V_n] \subseteq D_i$ for every $n \in \omega$. \square

Our next aim is to find a partial order \mathbb{P} first with $\mathbb{P} \rightarrow (I_\omega)_2^1$; note that trees or $F_n(\kappa, 2)$ cannot satisfy $\mathbb{P} \rightarrow (I_\omega)_2^1$. Moreover:

Proposition 6.3. $\mathbb{P} \not\rightarrow (I_\omega)_2^1$ for every countable poset \mathbb{P} without maximal elements.

PROOF: Fix a countable poset \mathbb{P} without maximal elements. We construct a partition $\mathbb{P} = P_0 \cup P_1$ witnessing $\mathbb{P} \not\rightarrow (I_\omega)_2^1$ as follows: First, fix an enumeration $\{I_n : n \in \omega\}$ of all intervals $I = [p', p]$ in \mathbb{P} which contain an infinite chain and let $\mathbb{P} = \{p_n : n \in \omega\}$ denote a 1-1 enumeration. Construct disjoint $P_{0,n}, P_{1,n} \subseteq \mathbb{P}$ by induction on $n \in \omega$ such that

- (i) $P_{i,n}$ is a finite union of antichains for $i < 2$,
- (ii) $p_n \in P_{0,n} \cup P_{1,n}$,
- (iii) $I_n \cap P_{i,n} \neq \emptyset$ for $i < 2$,
- (iv) whenever $C = \{c_k : k \in \omega\} \subseteq \mathbb{P}$ is a strictly increasing chain, $p_n \in C$ and $[c_i, c_j]$ is well-founded (i.e., $[c_i, c_j] \notin I$) for all $i < j < \omega$ then

$$\bigcup_{k \in \omega} [c_0, c_k] \cap P_{i,n} \neq \emptyset$$

for each $i < 2$.

Provided we can carry out this induction, we have that

Claim 6.4. $\mathbb{P} \dashv\vdash (I_\omega)_2^1$.

PROOF: Let $P_i = \bigcup\{P_{i,n} : n \in \omega\}$ for $i < 2$ and note that this is a partition of \mathbb{P} by (ii). Consider an arbitrary strictly increasing chain $C = \{c_k : k \in \omega\} \subseteq \mathbb{P}$. If there is $k \in \omega$ such that $[c_0, c_k]$ contains an infinite chain in \mathbb{P} then there is an $n \in \omega$ such that $I_n = [c_0, c_k]$; property (iii) from above ensures that $P_i \cap [c_0, c_k] \neq \emptyset$ for $i < 2$. Otherwise, the intervals $[c_i, c_j]$ are all well-founded intervals; in this case, property (iv) ensures that $\bigcup_{k \in \omega} [c_0, c_k] \cap P_i \neq \emptyset$ for $i < 2$. □

Now suppose we constructed $P_{i,n-1}$ satisfying the above conditions for $i < 2$. Note that finitely many elements can be added to both $P_{0,n-1}$ and $P_{1,n-1}$ without violating (i), thus (ii) and (iii) are easy to satisfy (note that $I_n \setminus (P_{0,n-1} \cup P_{1,n-1})$ is infinite since I_n contains an infinite chain).

It suffices to show the following to finish our proof:

Claim 6.5. Fix $p \in \mathbb{P}$ and $A \subseteq \mathbb{P}$ which is covered by finitely many antichains. Then there is an antichain $B \subseteq \mathbb{P} \setminus A$ such that whenever $C = \{c_k : k \in \omega\} \subseteq \mathbb{P}$ is a strictly increasing chain, $p \in C$ and the intervals $[c_i, c_j]$ are all well-founded then

$$\bigcup_{k \in \omega} [c_0, c_k] \cap B \neq \emptyset.$$

PROOF: Let

$$R = \{q \in P : p \leq q \text{ and } [p, q] \text{ does not contain infinite chains}\}.$$

Then $\langle R, \leq \rangle$ is well founded, so we can define, by well-founded recursion, a rank function rk from R into the ordinals such that

$$(6.1) \quad \begin{aligned} rk(p) &= 0, \\ rk(t) &= \sup\{rk(s) + 1 : s \in [p, t)\} \quad \text{if } t \in R, p < t. \end{aligned}$$

Let $Q = R \setminus A$ and define q^- to be the element minimizing rk on $[p, q] \setminus A$ for $q \in Q$. Let

$$B = \{q^- : q \in Q\}.$$

First note that B is an antichain by (6.1). Now fix a strictly increasing chain $C = \{c_k : k \in \omega\} \subseteq P$ such that the intervals $[c_i, c_j]$ are all well-founded and $p \in C$. Since A is covered by finitely many antichains there is $q \in C \setminus A$ such that $p < q$; also, $q \in Q$ by $[p, q]$ being well founded. Thus $q^- \in \bigcup_{k \in \omega} [c_0, c_k] \cap B$. \square

Indeed, to finish the inductive construction, apply the claim twice to find antichain $B_0 \subseteq \mathbb{P} \setminus A$ and $B_1 \subseteq \mathbb{P} \setminus (A \cup B_0)$ such that $\bigcup_{k \in \omega} [c_0, c_k] \cap B_i \neq \emptyset$ whenever $C = \{c_k : k \in \omega\} \subseteq \mathbb{P}$ is a strictly increasing chain, $p \in C$ and the intervals $[c_i, c_j]$ are all well-founded.

Then $P_{0,n} = P_{0,n-1} \cup B_0$ and $P_{1,n} = P_{1,n-1} \cup B_1$ are appropriate extensions satisfying (iv). \square

We will call a countable strictly increasing sequence of elements of a poset \mathbb{P} a *branch*; we say that a branch $x = (x_n)_{n \in \omega}$ goes above an element $p \in \mathbb{P}$ if $p \leq x_n$ for some $n \in \omega$.

Theorem 6.6. *There is a partial order \mathbb{P} of size ω_1 without maximal elements such that $\mathbb{P} \rightarrow (I_\omega)_2^1$. Furthermore,*

- (1) every $p \in \mathbb{P}$ has finitely many predecessors,
- (2) if $p \not\leq q$ in \mathbb{P} then there is a branch x in \mathbb{P} which goes above q but not p .

PROOF: Let us fix a function $c : [\omega_1]^2 \rightarrow \omega$ such that $c(\cdot, \zeta) : \zeta \rightarrow \omega$ is 1-1 for every $\zeta \in \omega_1$. It is easy to see that such functions satisfy the following:

Fact 6.7. *If $c(\cdot, \zeta) : \zeta \rightarrow \omega$ is 1-1 for every $\zeta \in \omega_1$ for some $c : [\omega_1]^2 \rightarrow \omega$ then for every uncountable, disjoint family $\mathcal{A} \subseteq [\omega_1]^{<\omega}$ and $N \in \omega$ there are $a < b$ ¹ in \mathcal{A} such that $c(\xi, \zeta) > N$ for every $\xi \in a, \zeta \in b$.*

Also, fix an enumeration $\{(y_\alpha, w_\alpha) : \omega \leq \alpha < \omega_1\}$ of all pairs of elements of $\omega_1 \times \omega$ such that $y_\alpha, w_\alpha \in \alpha \times \omega$.

We define $\mathbb{P} = (\omega_1 \times \omega, \leq)$ as follows: By induction on $\alpha \in L_1$ (where L_1 stands for the limit ordinals in ω_1) we construct a poset $\mathbb{P}_\alpha = ((\alpha + \omega) \times \omega, \leq_\alpha)$ with properties:

- (i) \mathbb{P}_α has no maximal elements and every $p \in \mathbb{P}_\alpha$ has finitely many predecessors,
- (ii) $\leq_\alpha \upharpoonright \beta = \leq_\beta$ for all $\beta < \alpha$,
- (iii) $(\xi, n) <_\alpha (\zeta, m)$ implies that $\xi < \zeta$ and $\max(n, c(\xi, \zeta)) < m$,
- (iv) there is a $t_\alpha \in \mathbb{P}_\alpha$ such that $t <_\alpha t_\alpha$ if and only if $t \leq_\alpha y_\alpha$ or $t \leq_\alpha w_\alpha$ for any $t \in \mathbb{P}_\alpha$,
- (v) if $p \not\leq q$ in \mathbb{P}_α then there is a branch x in \mathbb{P}_α which goes above q but not p .

We only sketch the inductive step. Suppose that $y_\alpha = (\xi, n)$ and $w_\alpha = (\zeta, m)$. Let $\Gamma = \{\nu < \omega_1 : \text{there is } s \leq y_\alpha \text{ or } s \leq w_\alpha \text{ with } s = (\nu, l) \text{ for some } l \in \omega\}$ and note that $|\Gamma| < \omega$ by (i). Let

$$k = \max\{n, m, c(\nu, \alpha) : \nu \in \Gamma\} + 1.$$

¹ $a < b$ iff $\xi < \zeta$ for all $\xi \in a, \zeta \in b$.

Now define $t_\alpha = (\alpha, k)$ and \leq_α so that $t <_\alpha t_\alpha$ implies that $t \leq_\alpha y_\alpha$ or $t \leq_\alpha w_\alpha$. Extend \leq_α further so that \mathbb{P}_α has no maximal elements and satisfies (v); this can be done by “placing” copies of $2^{<\omega}$ above elements of $\mathbb{P}_\alpha \setminus \bigcup\{\mathbb{P}_\beta : \beta < \alpha\}$.

Let us define $\mathbb{P} = \bigcup\{\mathbb{P}_\alpha : \alpha < \omega_1\}$ and $\leq = \bigcup\{\leq_\alpha : \alpha < \omega_1\}$; observe that (\mathbb{P}, \leq) is well defined and trivially satisfies (1) and (2). In what follows, π_{ω_1} and π_ω denote the projections from $\omega_1 \times \omega$ to the first and second coordinates respectively.

Claim 6.8. $\mathbb{P} \rightarrow (I_\omega)_2^1$.

PROOF: Suppose that $\mathbb{P} = D_0 \cup D_1$; we can assume that D_0 and D_1 are both cofinal in \mathbb{P} . Now suppose that there is no increasing chain with each interval in one of the D_i and reach a contradiction as follows. We will say that an interval $[s, t]$ in \mathbb{P} is i -maximal for some $i < 2$ if $[s, t] \subseteq D_i$ but $[s, t'] \not\subseteq D_i$ for every $t < t' \in \mathbb{P}$. Observe that for every $s \in D_i$ there is $t \in D_i$ such that $[s, t]$ is i -maximal; otherwise, we can construct an increasing chain starting from s with each interval in D_i . Now construct increasing 4-element sequences $R_\alpha = \{\tilde{x}_\alpha, \tilde{y}_\alpha, \tilde{z}_\alpha, \tilde{w}_\alpha\} \subseteq \mathbb{P}$ for $\alpha < \omega_1$ such that $\tilde{x}_\alpha \leq \tilde{y}_\alpha \leq \tilde{z}_\alpha \leq \tilde{w}_\alpha$ and

- (a) $[\tilde{x}_\alpha, \tilde{y}_\alpha] \subseteq \mathbb{P}_0$ is a 0-maximal interval,
- (b) $[\tilde{z}_\alpha, \tilde{w}_\alpha] \subseteq \mathbb{P}_1$ is a 1-maximal interval,
- (c) $\pi''_{\omega_1} R_\alpha < \pi''_{\omega_1} R_\beta$ if $\alpha < \beta$.

By passing to a subsequence of $\{R_\alpha : \alpha < \omega_1\}$ we can suppose that the image of $(\tilde{x}_\alpha, \tilde{y}_\alpha, \tilde{z}_\alpha, \tilde{w}_\alpha)$ under π_ω is independent of $\alpha < \omega_1$ and we let $N = \max \pi''_\omega R_\alpha$. Find $\alpha < \beta$, using Fact 6.7, such that

$$c \upharpoonright [\pi''_{\omega_1} R_\alpha, \pi''_{\omega_1} R_\beta] > N.$$

Observe that $\tilde{x}_\alpha \not\leq \tilde{w}_\beta$ by $\pi''_\omega w_\beta = N < c(\pi''_{\omega_1} \tilde{x}_\alpha, \pi''_{\omega_1} \tilde{w}_\beta)$ and (iii). Now find $\gamma < \omega_1$ such that $(y_\gamma, w_\gamma) = (\tilde{y}_\alpha, \tilde{w}_\beta)$ and consider $t_\gamma \in \mathbb{P}_\gamma$. We claim that t_γ is a minimal extension of \tilde{y}_α and \tilde{w}_β in the following sense:

- (1) $[\tilde{x}_\alpha, t_\gamma] = [\tilde{x}_\alpha, \tilde{y}_\alpha] \cup \{t_\gamma\}$,
- (2) $[\tilde{z}_\beta, t_\gamma] = [\tilde{z}_\beta, \tilde{w}_\beta] \cup \{t_\gamma\}$.

Indeed, if $\tilde{x}_\alpha \leq t' < t_\gamma$ then $t' \leq \tilde{y}_\alpha$ or $t' \leq \tilde{w}_\beta$; $\tilde{x}_\alpha \not\leq \tilde{w}_\beta$ implies that $t' \not\leq w_\beta$ hence $t' \in [\tilde{x}_\alpha, \tilde{y}_\alpha]$. Similarly, if $\tilde{z}_\beta \leq t' < t_\gamma$ then $t' \leq \tilde{y}_\alpha$ or $t' \leq \tilde{w}_\beta$; however, $t' \not\leq \tilde{y}_\alpha$ by $\pi''_\omega t' > \pi''_\omega \tilde{y}_\alpha$ so $t' \in [\tilde{z}_\beta, \tilde{w}_\beta]$.

Note that $t \in \mathbb{P}_0$ contradicts the 0-maximality of $[\tilde{x}_\alpha, \tilde{y}_\alpha]$ and (1) while $t \in \mathbb{P}_1$ contradicts the 1-maximality of $[\tilde{z}_\beta, \tilde{w}_\beta]$ and (2). □

The above claim finishes the proof. □

Using the previous theorem, we construct an irresolvable self-filling family; we can actually realize this family as a system of open sets in a first countable compact space. We remark that this space is base resolvable, as every compact space, by Corollary 3.7.

Theorem 6.9. *There is a first countable Corson compact space (X, τ) and $\mathcal{U} \subseteq \tau$ such that \mathcal{U} fills $\{\bigcap \mathcal{V} : \mathcal{V} \in [\mathcal{U}]^{<\omega}\}$ and \mathcal{U} is irresolvable.*

PROOF: Consider the poset \mathbb{P} in Theorem 6.6. We say that $x \in [\mathbb{P}]^\omega$ is a *maximal chain* if and only if $\{x(n)\}_{n \in \omega}$ is a branch in \mathbb{P} , $x(0)$ is a minimal element of \mathbb{P} and $[x(n), x(n+1)] = \{x(n), x(n+1)\}$. Note that there are no increasing chains of order type $\omega + 1$ in \mathbb{P} . Furthermore, since the intervals are finite we have

Observation 6.10. (1) *Any branch $y \in [\mathbb{P}]^\omega$ can be extended to a maximal chain $\bar{y} \in [\mathbb{P}]^\omega$.*

(2) *There is an $n_0 \in \omega$ such that $\bigcup_{n_0 \leq n} [\bar{y}(n_0), \bar{y}(n)] \subseteq \bigcup_{n \in \omega} [y(0), y(n)]$.*

Note that (2) implies that if $y \in [\mathbb{P}]^\omega$ has homogeneous intervals with respect to some coloring of \mathbb{P} then the end-segment of the maximal extension \bar{y} has the same property.

Now consider $X = \{x \in [\mathbb{P}]^\omega : x \text{ is a maximal chain}\}$ as a subspace of $2^{\mathbb{P}}$; here $2^{\mathbb{P}}$ is equipped with the usual product topology.

Claim 6.11. *X is a compact subspace of $\Sigma(2^{\mathbb{P}}) = \Sigma(2^{\omega_1})$.*

PROOF: $\Sigma(2^{\mathbb{P}}) = \Sigma(2^{\omega_1})$ follows from $|\mathbb{P}| = \omega_1$ and clearly every chain is countable so $X \subseteq \Sigma(2^{\mathbb{P}})$.

We prove that X is a closed subset of $2^{\mathbb{P}}$. Suppose that $y \in 2^{\mathbb{P}} \setminus X$; clearly, if y is not a chain then y can be separated from X . Suppose that y is a chain, then either $y(0)$ is not minimal in \mathbb{P} or there is $n \in \omega$ such that $[y(n), y(n+1)] \neq \{y(n), y(n+1)\}$. In the first case let $\varepsilon \in Fn(\mathbb{P}, 2)$ be defined to be 1 on $y(0)$ and $\varepsilon(p) = 0$ for $p < y(0)$, $p \in \mathbb{P}$ (note that each element in \mathbb{P} has only finitely many predecessors); then $y \in [\varepsilon]$ and $[\varepsilon] \cap X = \emptyset$. In the second case let $\varepsilon \in Fn(\mathbb{P}, 2)$ such that $1 = \varepsilon(y(n)) = \varepsilon(y(n+1))$ and $\varepsilon \upharpoonright [y(n), y(n+1)] \setminus \{y(n), y(n+1)\} = 0$; then $y \in [\varepsilon]$ and $[\varepsilon] \cap X = \emptyset$. □

Claim 6.12. *$\{x\} = \bigcap \{[\chi_{x(n)}] \cap X : n \in \omega\}$ for every $x \in X$. Hence every point in X has countable pseudocharacter; in particular, X is first countable.*

PROOF: Suppose that $y \in \bigcap \{[\chi_{x(n)}] \cap X : n \in \omega\}$, that is, $\{x(n) : n \in \omega\} \subset \{y(n) : n \in \omega\}$. We prove that $x(n) = y(n)$ by induction on $n \in \omega$. First, we have $y(0) = x(0)$ as they are comparable minimal elements in \mathbb{P} . Suppose that $x(i) = y(i)$ for $i < n$; if $x(n) \neq y(n)$ then $x(n) = y(k)$ for some $n < k$, thus $y(n) \in [x(n-1), x(n)] = [y(n-1), y(k)]$ which contradicts the maximality of x . □

Now define

$$V_p = \{x \in X : \exists n \in \omega : x(n) \geq p\} \text{ for } p \in \mathbb{P},$$

and note that V_p is open since $V_p = \bigcup \{[\chi_{\{q\}}] \cap X : p \leq q\}$. We define

$$\mathcal{U} = \{V_p : p \in \mathbb{P}\}.$$

Claim 6.13. *\mathcal{U} fills $\{\bigcap \mathcal{V} : \mathcal{V} \in [\mathcal{U}]^{<\omega}\}$ and \mathcal{U} is irresolvable.*

PROOF: Note that $p < q$ in \mathbb{P} if and only if $V_q \subsetneq V_p$; the nontrivial direction is implied by property (2) of \mathbb{P} in Theorem 6.6. To see that \mathcal{U} fills the finite

intersections from \mathcal{U} let $\mathcal{V} \in [\mathcal{U}]^{<\omega}$ be arbitrary. If $A = \{p \in \mathbb{P} : V_p \in \mathcal{V}\} \in [\mathbb{P}]^{<\omega}$ then

$$\bigcap \mathcal{V} = \bigcup \{V_q : p < q \text{ for all } p \in A\}.$$

We show that \mathcal{U} is irresolvable. Suppose that we partitioned \mathcal{U} , equivalently \mathbb{P} into two parts $\mathbb{P}_0, \mathbb{P}_1$. Applying $\mathbb{P} \rightarrow (I_\omega)_2^1$ we have that there is a chain $y \in \mathbb{P}^\omega$ and $i < 2$ such that $[y(0), y(n)] \subseteq \mathbb{P}_i$ for every $n \in \omega$. By Observation 6.10 there is maximal chain $\bar{y} \in X$ such that $[\bar{y}(n_0), \bar{y}(n)] \subseteq \mathbb{P}_i$ for some $n_0 \in \omega$ and every $n \geq n_0$. We claim that there is no $V \in \{V_p : p \in \mathbb{P}_{1-i}\}$ such that $\bar{y} \in V \subseteq V_{\bar{y}(n_0)}$. Indeed, if $\bar{y} \in V_p \subseteq V_{\bar{y}(n_0)}$ for some $p \in \mathbb{P}$ then $\bar{y}(n_0) \leq p$ and there is $n \in \omega \setminus n_0$ such that $p \leq \bar{y}(n)$; that is, $p \in [\bar{y}(n_0), \bar{y}(n)] \subseteq \mathbb{P}_i$. \square

The last claim finishes the proof of the theorem. \square

Let us finish this section with the following:

Lemma 6.14. *If \mathcal{U} fills $\{\bigcap \mathcal{V} : \mathcal{V} \in [\mathcal{U}]^{<\omega}\}$ and \mathcal{U} is irresolvable then there is a non base resolvable, T_0 topological space.*

PROOF: Suppose that $\mathcal{U} \subset \mathcal{P}(X)$ is as above. Define a relation \sim on X by $x \sim y$ if and only if $\{U \in \mathcal{U} : x \in U\} = \{U \in \mathcal{U} : y \in U\}$; clearly, \sim is an equivalence relation on X . Let $[x] = \{x' \in X : x \sim x'\}$ for $x \in X$ and let $[U] = \{[x] : x \in U\}$ for any $U \subset X$. It is clear that $[U] = \bigcup \{[V] : V \in \mathcal{V}\}$ if $U = \bigcup \mathcal{V}$ and $[U] = \bigcap \{[V] : V \in \mathcal{V}\}$ if $U = \bigcap \mathcal{V}$. Thus $\mathbb{B} = \{[U] : U \in \mathcal{U}\}$ is a base for a T_0 topology on $[X]$; sometimes this is referred to as the Kolmogorov quotient of the original (not necessarily T_0) topology generated by \mathcal{U} .

It remains to show that \mathbb{B} is an irresolvable base. Take a partition $\mathbb{B} = \mathbb{B}_0 \cup \mathbb{B}_1$. Note that

- (1) $[x] \in [U]$ if and only if $x \in U$,
- (2) $[U] = [V]$ if and only if $U = V$,
- (3) $[U] \subset [V]$ if and only if $U \subset V$

for any $U, V \in \mathcal{U}$; thus the partition $\mathbb{B}_0 \cup \mathbb{B}_1$ gives a partition $\mathcal{U}_i = \{U \in \mathcal{U} : [U] \in \mathbb{B}_i\}$ of \mathcal{U} . Now there is an $i < 2$ so that \mathcal{U}_i does not fill \mathcal{U} i.e., there is $x \in X$ and $V \in \mathcal{U}$ so that $x \in U$ implies $U \setminus V \neq \emptyset$ for all $U \in \mathcal{U}_i$. This gives that $[x] \in [U]$ implies $[U] \setminus [V] \neq \emptyset$ for all $[U] \in \mathbb{B}_i$; in particular, \mathbb{B}_i is not a base for the topology generated by \mathbb{B} . \square

In particular, we have the following

Corollary 6.15. *There is a non base resolvable, T_0 topological space.*

7. A 0-dimensional, Hausdorff space with an irresolvable base

In this section, we partially strengthen Corollary 6.15 by showing

Theorem 7.1. *It is consistent that there is a first countable, 0-dimensional, T_2 space which has a point countable, irresolvable base. Furthermore, the space has size \mathfrak{c} and weight ω_1 .*

PROOF: For $\langle \alpha, n \rangle, \langle \beta, m \rangle \in \omega_1 \times \omega$ write $\langle \alpha, n \rangle \triangleleft \langle \beta, m \rangle \in \omega_1 \times \omega$ if $\langle \alpha, n \rangle = \langle \beta, m \rangle$ or $(\alpha < \beta$ and $n < m)$.

Definition 7.2. If $\preceq_1, \preceq_2 \subset \triangleleft$, then let $\preceq_1 \sqcup \preceq_2$ be the partial order generated by $\preceq_1 \cup \preceq_2$.

Definition 7.3. If $\mathcal{A} = \langle \omega_1 \times \omega, \preceq \rangle$ is a poset with $\preceq \subset \triangleleft$, and for each $\alpha \in L_1$ we have a set $T_\alpha \subset \alpha \times \omega$ such that

(C) $\langle T_\alpha, \preceq \rangle$ is an everywhere ω -branching tree,

then we say that the pair $\langle \mathcal{A}, \langle T_\alpha : \alpha \in L_1 \rangle \rangle$ is a *candidate*.

Denote by $T_\alpha(n)$ the n^{th} level of the tree $\langle T_\alpha, \preceq \rangle$.

Definition 7.4. Fix a candidate $\mathbb{A} = \langle \mathcal{A}, \langle T_\alpha : \alpha \in L_1 \rangle \rangle$. We will define a topological space $X(\mathbb{A})$ as follows.

For $\alpha \in L_1$ let $\mathcal{B}(T_\alpha)$ be the collection of the cofinal branches of T_α , and let

$$\mathcal{B}(\mathbb{A}) = \bigcup \{ \mathcal{B}(T_\alpha) : \alpha \in L_1 \}.$$

The underlying set of the space $X(\mathbb{A})$ is $\mathcal{B}(\mathbb{A})$.

For $x \in \omega_1 \times \omega$ let $U(x) = \{ y \in \omega_1 \times \omega : x \preceq y \}$ and

$$V(x) = \{ b \in \mathcal{B}(\mathbb{A}) : \exists y \in b (x \preceq y) \}.$$

Clearly $V(x) = \{ b \in \mathcal{B}(\mathbb{A}) : b \subseteq^* U(x) \}$ where \subseteq^* denotes containment modulo finite.

We declare that the family

$$\mathcal{V} = \{ V(x) : x \in \omega_1 \times \omega \}$$

is the base of $X(\mathbb{A})$.

Lemma 7.5. \mathcal{V} is a base and so $X(\mathbb{A})$ is a topological space. Moreover, \mathcal{V} is point countable.

PROOF: Assume that $b \in V(x) \cap V(y)$. Then there is $z \in b$ such that $x \preceq z$ and $y \preceq z$. Then $b \in V(z) \subset V(x) \cap V(y)$.

To see that \mathcal{V} is point countable, note that $b \notin V(x)$ if $b \in \mathcal{B}(T_\alpha)$ and $x \in (\omega_1 \setminus \alpha) \times \omega$. □

For $x, y \in \omega_1 \times \omega$ with $x \preceq y$ let

$$[x, y] = \{ t \in \omega_1 \times \omega : x \preceq t \preceq y \}.$$

Definition 7.6. We say that a candidate $\mathbb{A} = \langle \mathcal{A}, \langle T_\alpha : \alpha \in L_1 \rangle \rangle$ is *good* if

(G1) $V(u) \supset V(v)$ if and only if $u \preceq v$.

(G2) $\forall \alpha \in L_1 \forall \zeta < \alpha (T_\alpha \setminus (\zeta \times \omega)) \neq \emptyset$.

(G3) (a) $\forall \alpha \in L_1 (\forall x, y \in T_\alpha) U(x) \cap U(y) \neq \emptyset$ if and only if x and y are \preceq -comparable.

(b) for each $\{\alpha, \beta\} \in [L_1]^2$ there is $f(\alpha, \beta) \in \omega$ such that

$$\forall x \in T_\alpha(f(\alpha, \beta)) \forall y \in T_\beta(f(\alpha, \beta)) U(x) \cap U(y) = \emptyset.$$

(G4) For each $x \in \omega_1 \times \omega$ and $\alpha \in L_1$ there is $g(x, \alpha) \in \omega$ such that for each $y \in T_\alpha(g(x, \alpha))$

$$U(y) \subset U(x) \text{ or } U(y) \cap U(x) = \emptyset.$$

(G5) If for all $\alpha \in L_1$ and $\zeta < \alpha$ we choose a four element \preceq -increasing sequence

$$\langle x_\zeta^\alpha, y_\zeta^\alpha, z_\zeta^\alpha, w_\zeta^\alpha \rangle \subset T_\alpha \setminus (\zeta \times \omega)$$

then there are $\{\alpha, \beta\} \in [L_1]^2$, $\zeta < \alpha$, $\xi < \beta$, and $t \in T_\alpha \cap T_\beta$ such that

- (i) $y_\zeta^\alpha \prec t$ and $[x_\zeta^\alpha, t] = [x_\zeta^\alpha, y_\zeta^\alpha] \cup \{t\}$,
- (ii) $w_\xi^\beta \prec t$ and $[z_\xi^\beta, t] = [z_\xi^\beta, w_\xi^\beta] \cup \{t\}$.

Basically (G3) will force the space to be Hausdorff, (G4) ensures that each $V(x)$ is clopen and (G5) will be used in proving irresolvability. Indeed, we have

Lemma 7.7. *If \mathbb{A} is a good candidate, then $X(\mathbb{A})$ is a dense-in-itself, first countable, 0-dimensional T_2 space such that the base $\{V(x) : x \in \omega_1 \times \omega\}$ is point countable and irresolvable.*

PROOF: We prove this lemma in several steps.

Claim 7.8. *$X(\mathbb{A})$ is dense-in-itself.*

Indeed, assume that $b \in B(T_\alpha)$ and $V(x)$ is an open neighbourhood of b . Then there is $y \in b$ with $x \preceq y$ and so $b \in V(y) \subset V(x)$. Thus $V(x) \supset V(y) \supset \{b' \in B(T_\alpha) : y \in b'\}$, and so $V(x)$ has 2^ω many elements. So b is not isolated.

Claim 7.9. *$X(\mathbb{A})$ is T_2 .*

Indeed, let $b \in B(T_\alpha)$ and $c \in B(T_\beta)$ so that $b \neq c$.

If $\alpha = \beta$ then pick $n \in \omega$ such that x , the n^{th} element of b , and y , the n^{th} element of c , are different. Then $b \in V(x)$, $c \in V(y)$ and $V(x) \cap V(y) = \emptyset$ by (G3)(a).

If $\alpha \neq \beta$ then write $n = f(\alpha, \beta)$ (see (G3)(b)), let x be the n^{th} element of b , and let y be the n^{th} element of c . Then $b \in V(x)$, $c \in V(y)$ and $V(x) \cap V(y) = \emptyset$ by (G3)(b).

Claim 7.10. *Each set in $\{V(x) : x \in \omega_1 \times \omega\}$ is clopen, thus $X(\mathbb{A})$ is 0-dimensional.*

Indeed, assume that $x \in \omega_1 \times \omega$, $b \in B(T_\alpha)$ and $b \notin V(x)$. Let $\{y\} = b \cap T_\alpha(g(\alpha, x))$. Then $y \notin U(x)$ because $b \notin V(x)$, so $U(x) \cap U(y) = \emptyset$ by (G4). Thus $V(x) \cap V(y) = \emptyset$ as well.

Claim 7.11. *The base $\{V(x) : x \in \omega_1 \times \omega\}$ is irresolvable.*

Assume on the contrary that there is a partition (K_0, K_1) of $\omega_1 \times \omega$ such that both $\mathcal{V}_0 = \{V(x) : x \in K_0\}$ and $\mathcal{V}_1 = \{V(x) : x \in K_1\}$ are bases.

Assume that $\alpha \in L_1$, $x, y \in T_\alpha$ with $x \preceq y$ and $i \in 2$. We say that interval $[x, y]$ is *i-maximal* in T_α if

- (i) $[x, y] \subset K_i$, but $[x, z] \not\subset K_i$ for any z with $y \prec z \in T_\alpha$.

Subclaim 7.11.1. *If $\alpha \in L_1$ and $x \in T_\alpha \cap K_i$, then there is $x \preceq y \in T_\alpha$ such that the interval $[x, y]$ is i-maximal in T_α .*

PROOF OF THE CLAIM: Assume on the contrary that there is no such y . Then we can construct a strictly increasing sequence $\langle x, y_0, y_1, \dots \rangle$ in T_α such that $[x, y_n] \subset K_i$ for all $n < \omega$.

Then $b = \{y \in T_\alpha : \exists n \in \omega \ y \preceq y_n\} \in \mathcal{B}(T_\alpha)$.

Since $b \in V(x)$, and we assumed that $\{V(z) : z \in K_{1-i}\}$ is a base, there is $z \in K_{1-i}$ with $b \in V(z) \subset V(x)$. Then $x \preceq z$ by (G1). Moreover, there is $y \in b$ with $z \prec y$ because $b \in V(z)$. Thus $z \in [x, y] \cap K_{1-i}$, so $[x, y] \not\subset K_i$. Contradiction, the subclaim is proved. \square

Using the subclaim, for all $\alpha \in L_1$ and for all $\zeta < \alpha$ we will construct a four element \preceq -increasing sequence

$$\langle x_\zeta^\alpha, y_\zeta^\alpha, z_\zeta^\alpha, w_\zeta^\alpha \rangle \subset T_\alpha \setminus (\zeta \times \omega)$$

as follows.

First, using (G2) pick $s_\zeta^\alpha \in T_\alpha \setminus (\zeta \times \omega)$.

If $K_0 \cap U(s_\zeta^\alpha) \cap T_\alpha = \emptyset$, then let $x_\zeta^\alpha = y_\zeta^\alpha = s_\zeta^\alpha$.

Otherwise pick

$$x_\zeta^\alpha \in K_0 \cap U(s_\zeta^\alpha) \cap T_\alpha,$$

and then, using the Subclaim above, pick

$$y_\zeta^\alpha \in U(x_\zeta^\alpha) \cap T_\alpha$$

such that

$$[x_\zeta^\alpha, y_\zeta^\alpha] \text{ is 0-maximal in } T_\alpha.$$

If $K_1 \cap U(y_\zeta^\alpha) \cap T_\alpha = \emptyset$, then let $z_\zeta^\alpha = w_\zeta^\alpha = y_\zeta^\alpha$.

Otherwise pick

$$z_\zeta^\alpha \in K_1 \cap U(y_\zeta^\alpha) \cap T_\alpha,$$

and then, using the Subclaim above, pick

$$w_\zeta^\alpha \in U(z_\zeta^\alpha) \cap T_\alpha$$

such that

$$[z_\zeta^\alpha, w_\zeta^\alpha] \text{ is 1-maximal in } T_\alpha.$$

By (G5), there are $\{\alpha, \beta\} \in [L_1]^2$, $\zeta < \alpha$, $\xi < \beta$, and $t \in T_\alpha \cap T_\beta$ such that

- (i) $y_\zeta^\alpha \prec t$ and $[x_\zeta^\alpha, t] = [x_\zeta^\alpha, y_\zeta^\alpha] \cup \{t\}$,

(ii) $w_\xi^\beta \prec t$ and $[z_\xi^\beta, t] = [z_\xi^\beta, w_\xi^\beta] \cup \{t\}$.

Assume first that $t \in K_0$. Then $t \in K_0 \cap T_\alpha$, and $[x_\zeta^\alpha, t] = [x_\zeta^\alpha, y_\zeta^\alpha] \cup \{t\}$, so $[x_\zeta^\alpha, t] \subset K_0$, i.e., $[x_\zeta^\alpha, y_\zeta^\alpha]$ was not 0-maximal in T_α . Contradiction. If $t \in K_1$, then a similar argument works using the interval $[z_\xi^\beta, w_\xi^\beta]$ and K_1 .

So in both cases we obtained a contradiction, so the base $\{V(x) : x \in \omega_1 \times \omega\}$ is irresolvable, which proves the lemma. \square

Next we show that some c.c.c. forcing introduces a good candidate which finishes the proof the theorem.

Define the poset $\mathcal{P} = \langle P, \preceq \rangle$ as follows. The underlying set consists of 6-tuples

$$\langle A, \preceq, I, \{T_\alpha : \alpha \in I\}, f, g \rangle,$$

where

- (P1) $A \in [\omega_1 \times \omega]^{<\omega}$, $\langle A, \preceq \rangle$ is a poset, $\preceq \subset \triangleleft$, $I \in [\omega_1]^{<\omega}$,
- (P2) $T_\alpha \subset (A \cap \alpha) \times \omega$ and $\langle T_\alpha, \preceq \rangle$ is a tree for $\alpha \in I$,
- (P3) f and g are functions, $\text{dom}(f) \subset [I]^2$, $\text{dom}(g) \subset A \times I$, $\text{ran}(f) \cup \text{ran}(g) \subset \omega$.
- (P4) To simplify our notation write $U(x) = \{y \in A : x \preceq y\}$ for $x \in A$.
 - (a) If $\alpha \in I$ and $x, y \in T_\alpha$ then $U(x) \cap U(y) \neq \emptyset$ if and only if x and y are \preceq -comparable.
 - (b) If $\{\alpha, \beta\} \in [\text{dom}(f)]^2$ and $n = f(\alpha, \beta)$, then

$$U[T_\alpha(n)] \cap U[T_\beta(n)] = \emptyset \text{ and } U[T_\alpha(n)] \cap T_\beta(<n) = \emptyset.$$

- (P5) If $\langle x, \alpha \rangle \in \text{dom}(g)$ then for all $y \in T_\alpha(g(x, \alpha))$ we have $U(y) \subset U(x)$ or $U(y) \cap U(x) = \emptyset$.

For $p \in P$ write $p = \langle A^p, \preceq^p, I^p, \{T_\alpha^p : \alpha \in I^p\}, f^p, g^p \rangle$, and for $x \in A^p$ let $U^p(x) = \{y \in A^p : x \preceq^p y\}$.

For $p, q \in P$ let $p \leq q$ if and only if

- (O1) $A^p \supset A^q$, and $\preceq^q = \preceq^p \upharpoonright A^q$,
- (O2) $I^p \supset I^q$ and $T_\alpha^q = T_\alpha^p \cap A^q$ for $\alpha \in I^q$,
- (O3) if $x \in A^p \setminus A^q$, then $U^p(x) \cap A^q = \emptyset$,
- (O4) $f^p \supset f^q$ and $g^p \supset g^q$,
- (O5) if $U^q(x) \cap U^q(y) = \emptyset$ then $U^p(x) \cap U^p(y) = \emptyset$.

Clearly \leq is a partial order on P .

For $p \in P$ write $\text{supp}(p) = I^p \cup \{\alpha : \langle \alpha, n \rangle \in A^p \text{ for some } n \in \omega\}$.

If \mathcal{G} is a \mathcal{P} -generic filter, then let

$$A = \bigcup \{A^p : p \in \mathcal{G}\},$$

$$\preceq = \bigcup \{\preceq^p : p \in \mathcal{G}\},$$

$$I = \bigcup \{I^p : p \in \mathcal{G}\},$$

$$T_\alpha = \bigcup \{T_\alpha^p : \alpha \in p \in \mathcal{G}\} \text{ for } \alpha \in L_1,$$

$$f = \bigcup \{f^p : p \in \mathcal{G}\},$$

$$g = \bigcup \{g^p : p \in \mathcal{G}\}.$$

We show that \mathcal{P} satisfies c.c.c. and $\mathbb{A} = \langle \langle \omega_1 \times \omega, \preceq \rangle, \{T_\alpha : \alpha \in L_1\} \rangle$ is a good candidate.

Definition 7.12. We say that the conditions p and q are *twins* if conditions (T1)–(T7) below are satisfied:

(T1) $|\text{supp}(p)| = |\text{supp}(q)|$, moreover $\max(\text{supp}(p) \cap \text{supp}(q)) < \min(\text{supp}(p) \triangle \text{supp}(q))$.

Denote by ρ the unique order preserving bijection between $\text{supp}(p)$ and $\text{supp}(q)$, and define the function $\underline{\rho} : \text{supp}(p) \times \omega \rightarrow \text{supp}(q) \times \omega$ by the formula $\underline{\rho}(\langle \alpha, n \rangle) = \langle \rho(\alpha), n \rangle$.

(T2) $\underline{\rho}'' A^p = A^q$.

(T3) $x \preceq^p y$ iff $\underline{\rho}(x) \preceq^q \underline{\rho}(y)$.

(T4) $\rho'' I^p = I^q$.

(T5) $T_{\rho(\alpha)}^q = \underline{\rho}'' T_\alpha$.

(T6) $f^p(x, y) = m$ iff $f^q(\underline{\rho}(x), \underline{\rho}(y)) = m$.

(T7) $g^p(x, \alpha) = m$ iff $g^q(\underline{\rho}(x), \rho(\alpha)) = m$.

Lemma 7.13. *If p and q are twins then*

$$p \oplus q = \langle A^p \cup A^q, \preceq^p \cup \preceq^q, I^p \cup I^q, \{T_\alpha^p \cup T_\alpha^q : \alpha \in I^p \cup I^q\}, f^p \cup f^q, g^p \cup g^q \rangle$$

is a common extension of p and q , where $T_\alpha^p = \emptyset$ for $\alpha \in I^q \setminus I^p$ and $T_\alpha^q = \emptyset$ for $\alpha \in I^p \setminus I^q$.

PROOF: Straightforward. □

Lemma 7.14. *There is a function φ from P into some countable set such that if $\varphi(p) = \varphi(q)$ and $\text{supp}(p) \cap \text{supp}(q) < \text{supp}(p) \triangle \text{supp}(q)$, then p and q are twins.*

PROOF: Let $\varphi(p)$ be the type of the first order structure

$$\langle \text{supp}(p) \times \omega, A^p, \preceq^p, I^p, \{T_\alpha^p : \alpha \in I^p\}, f^p, g^p \rangle. \quad \square$$

Lemmas 7.13 and 7.14 yield that \mathcal{P} satisfies c.c.c

Lemma 7.15. *$A = \omega_1 \times \omega$, $I = L_1$ and $T_\gamma(0) \setminus (\zeta \times \omega)$ is infinite for all $\gamma \in L_1$ and $\zeta < \gamma$, and so (G2) holds.*

PROOF: For $p \in P$, $\gamma \in L_1$ and $y \in (\gamma \times \omega) \setminus A^p$ define $p \uplus \{y\}_\gamma$ as follows:

$$p \uplus \{y\}_\gamma = \langle A^p \cup \{y\}, \preceq^p, I^p \cup \{\gamma\}, \{T_\gamma^p \cup \{y\}, T_\alpha^p : \alpha \in I^p \setminus \{\gamma\}\}, f^p, g^p \rangle.$$

Then $q = p \uplus \{y\}_\gamma \in P$ and $p \uplus \{y\}_\gamma \leq p$. If $y \notin \zeta \times \omega$ then $q \Vdash y \in T_\gamma \setminus (\zeta \times \omega)$ so we are done. □

Lemma 7.16. (a) Assume that $p \in P$, $a \in T_\gamma^p$ and $b \in (\gamma \times \omega) \setminus A^p$ with $a \triangleleft b$. Let

$$p \uplus_a \{b\}_\gamma = \langle A^p \cup \{b\}, \preceq^p \sqcup \{a, b\}, \{T_\gamma^p \cup \{b\}, T_\alpha^p : \alpha \in I^p \setminus \{\gamma\}\}, f^p, g^p \rangle.$$

Then $p \uplus_a \{b\}_\gamma \in P$ and $p \uplus_a \{b\}_\gamma \leq p$.

(b) The structure \mathbb{A} is a candidate.

PROOF: First we check $q = p \uplus_a \{b\}_\gamma \in P$.

(P1)-(P3) are straightforward.

(P4)(a) Since $U^q(b) = \{b\}$, we can assume that $x, y \neq b$. If $U^p(x) \cap U^p(y) \neq \emptyset$ then x and y are \preceq^p -comparable. So we can assume that $b \in U^q(x) \cap U^q(y)$. But then $a \in U^p(x) \cap U^p(y)$, so we are done.

(P4)(b) Assume that $x \in T_\alpha^q(n)$, $y \in T_\beta^q(n)$ with $n = f^p(\alpha, \beta) = f^q(\alpha, \beta)$ and $z \in U^q(x) \cap U^q(y)$. If $z \neq b$ then $z \in U^p(x) \cap U^p(y)$ which is not possible. So $z = b$.

If $x, y \neq b$, then $a \in U^p(x) \cap U^p(y)$ which is not possible. So we can assume that $x = b$ and $\alpha = \gamma$. So $b \in T_\alpha^q(n)$ and so $a \in T_\alpha^p(n-1)$. Thus $T_\alpha^p(n-1) \cap U^p(y) \neq \emptyset$ which is not possible because (P4)(b) holds for p .

Assume that $x \in T_\alpha^q(n)$, $y \in T_\beta^q(<n)$ and $y \in U^q(x)$. If $y \neq b$ then $y \in U^p(x) \cap T_\beta^p(<n)$ which is not possible. So $y = b$ and $\beta = \gamma$. Thus $a \in T_\beta^p(<n) \cap U_\alpha^p(x)$ which is not possible because (P4)(b) holds for p .

(P5) Since $U(b) = \{b\}$, we can assume that $y \in A^p$. Since $b \in U^q(z)$ if and only if $a \in U^q(z)$ for $z \in A^p$, if $U^p(y) \subset U^p(x)$ then $U^q(y) \subset U^q(x)$, and if $U^p(y) \cap U^p(x) = \emptyset$ then $U^q(y) \cap U^q(x) = \emptyset$.

Thus we proved $q \in P$. Since $q \leq p$ is straightforward, we are done.

(b) is clear from (a) by standard density arguments. \square

Now our aim is to prove that \mathbb{A} is a good candidate.

Lemma 7.17. \mathbb{A} has property (G1).

PROOF: Assume that $p \in P$, $u, v \in A^p$, $v \notin U^p(u)$. Pick $\gamma \in L_1 \setminus I^p$ with $\text{supp}(p) \subset \gamma$, and pick $b \in \gamma \times \omega$ with $v \triangleleft b$.

Consider the condition $q = p \uplus_v \{b\}_\gamma \leq p$.

Since $b \in T_\gamma^q$, we have $V(b) \cap \mathcal{B}(T_\gamma) \neq \emptyset$, so $V(b) \neq \emptyset$. Since $U^q(u) \cap U^q(b) = \emptyset$ we have $U(u) \cap U(b) = \emptyset$, and so $V(u) \cap V(b) = \emptyset$, and so $\emptyset \neq V(b) \subset V(v) \setminus V(u)$. \square

Lemma 7.18. $\text{dom}(f) = [L_1]^2$ and $\text{dom}(g) = (\omega_1 \times \omega) \times L_1$. Hence (G3) and (G4) hold.

PROOF: Assume that $\{\gamma, \delta\} \in [I^p]^2 \setminus \text{dom}(f^p)$. Pick m such that $T_\alpha^p(m) = \emptyset$ for all $\alpha \in I^p$. Extend f^p to f^q as follows: $\text{dom}(f^q) = \text{dom}(f^p) \cup \{\{\gamma, \delta\}\}$ and $f^q(\gamma, \delta) = m$. Let

$$q = \langle A^p, \preceq^p, I^p, \{T_\alpha^p : \alpha \in I^p\}, f^q, g^p \rangle.$$

Then $q \in P$ and $q \leq p$.

Similar argument works for g . □

Finally we verify that (G5) also holds. Assume that

$$V^P \models \forall \alpha \in L_1 \forall \zeta < \alpha \langle x_\zeta^\alpha, y_\zeta^\alpha, z_\zeta^\alpha, w_\zeta^\alpha \rangle \subset T_\alpha \setminus (\zeta \times \omega) \text{ is } \preceq\text{-increasing.}$$

For all $\alpha \in L_1$ and $\zeta < \alpha$ pick a condition $p_\zeta^\alpha = \langle A_\zeta^\alpha, \preceq_\zeta^\alpha, \dots \rangle$ which decides the sequence $\langle x_\zeta^\alpha, y_\zeta^\alpha, z_\zeta^\alpha, w_\zeta^\alpha \rangle$ and $\{x_\zeta^\alpha, y_\zeta^\alpha, z_\zeta^\alpha, w_\zeta^\alpha\} \subset T_\zeta^\alpha$.

Let us say that a Δ -system $\mathcal{A} \subset [\omega]^{<\omega}$ is *nice* if $A \cap B < A \Delta B$ for all $A \neq B \in \mathcal{A}$.

Using the Fodor lemma, for each $\zeta \in \omega_1$ find $m_\zeta < \omega$ and $I_\zeta \in [L_1]^{\omega_1}$ such that

- (i) $\varphi(p_\zeta^\alpha) = m_\zeta$ for all $\alpha \in I_\zeta$, where φ is from Lemma 7.14.
- (ii) $\{\text{supp}(p_\zeta^\alpha) : \alpha \in I_\zeta\}$ forms a nice Δ -system with kernel S_ζ , moreover $\alpha \in \text{supp}(p_\zeta^\alpha) \setminus S_\zeta$.
- (iii) $\langle x_\zeta^\alpha, y_\zeta^\alpha, z_\zeta^\alpha, w_\zeta^\alpha \rangle = \langle x_\zeta, y_\zeta, z_\zeta, w_\zeta \rangle$ for $\alpha \in I_\zeta$.

Then $\{x_\zeta^\alpha, y_\zeta^\alpha, z_\zeta^\alpha, w_\zeta^\alpha\} = \{x_\zeta, y_\zeta, z_\zeta, w_\zeta\} \subset S_\zeta \times \omega$.

Find $m \in \omega$ and $I \in [\omega_1]^{\omega_1}$ such that

- (iv) $m_\zeta = m$ for all $\zeta \in I$, and so

$$\forall \zeta \in I \forall \alpha \in I_\zeta \varphi(p_\zeta^\alpha) = m.$$

- (v) $\{S_\zeta : \zeta \in I\}$ forms a nice Δ -system with kernel S .

Pick $\{\xi, \zeta\} \in [I]^2$. Then pick $\alpha \in I_\zeta$ such that $S_\xi \cup S_\zeta < \text{supp}(p_\zeta^\alpha) \setminus S_\zeta$. So

$$S < (S_\xi \cup S_\zeta) \setminus S < \text{supp}(p_\zeta^\alpha) \setminus S_\zeta.$$

Now pick $\beta \in I_\xi$ such that $\text{supp}(p_\zeta^\alpha) < \text{supp}(p_\xi^\beta) \setminus S_\xi$. So

$$S < (S_\xi \cup S_\zeta) \setminus S < \text{supp}(p_\zeta^\alpha) \setminus S_\zeta < \text{supp}(p_\xi^\beta) \setminus S_\xi.$$

Thus $\text{supp}(p_\zeta^\alpha) \cap \text{supp}(p_\xi^\beta) = S$, $\alpha \in \text{supp}(p_\zeta^\alpha) \setminus S_\zeta$ and $\beta \in \text{supp}(p_\xi^\beta) \setminus S_\xi$.

Since $\varphi(p_\zeta^\alpha) = \varphi(p_\xi^\beta)$, the conditions $\varphi(p_\zeta^\alpha)$ and $\varphi(p_\xi^\beta)$ are twins, and


$$q = p_\zeta^\alpha \oplus p_\xi^\beta$$

is a common extension. Pick $t \in (\alpha \times \omega) \setminus (A_\zeta^\alpha \cup A_\xi^\beta)$ with $y_\zeta \triangleleft t$ and $w_\xi \triangleleft t$.

Define r as follows:

$$r = \langle A^q, \preceq_q \underline{\cup} \langle y_\zeta, t \rangle \underline{\cup} \langle w_\xi, t \rangle, I^q,$$

$$\{T_\alpha^q \cup \{t\}, T_\beta^q \cup \{t\}, T^\gamma : \gamma \in I^q \setminus \{\alpha, \beta\}\}, f^q, g^q \rangle.$$



To check $r \in P$ we will use the following observation:

$$(7.1) \quad r \upharpoonright (\text{supp}(p_\zeta^\alpha) \cup \{t\}) = p_\zeta^\alpha \uplus_{y_\zeta^\alpha} \{t\}_\alpha$$

and

$$(7.2) \quad r \upharpoonright (\text{supp}(p_\xi^\beta) \cup \{t\}) = p_\xi^\beta \uplus_{w_\xi^\beta} \{t\}_\beta.$$

Now let us check (P1)–(P5).

(P1) is trivial for r .

(P2) Let $\gamma \in I^q$. If $\gamma \neq \alpha, \beta$, then $T_\gamma^q = T_\gamma^p$, so we are done.

Moreover, $T_\alpha^r = T_\alpha^q \cup \{t\}$, $t \in \alpha \times \omega$, and $\langle T_\alpha^r, \preceq \rangle$ is a tree by (7.1) and (7.2).

The same argument works for T_β^r .

(P3) is trivial.

(P4)(a) Assume that $\gamma \in I^r$, $x, y \in T_\gamma^r$ with $U^r(x) \cap U^r(y) \neq \emptyset$. Since $U^r(t) = \{t\}$ we can assume $x, y \in A^q$.

Assume that $\gamma \in I_\zeta^\alpha$. Then $T_\gamma^q \subset A_\zeta^\alpha$, and so $x, y \in A_\zeta^\alpha$. Thus $t \in U^r(x) \cap U^r(y)$ implies $y_\zeta^\alpha \in U^r(x) \cap U^r(y)$. So $U^q(x) \cap U^q(y) \neq \emptyset$, which yields that x and y are \preceq^q comparable because $q \in P$.

Similar argument works when $\gamma \in I_\xi^\beta$.

(P4)(b) Assume that $\{\alpha', \beta'\} \in \text{dom}(f^r) = \text{dom}(f^q) = \text{dom}(p_\zeta^\alpha) \cup \text{dom}(p_\xi^\beta)$. We can assume that $\{\alpha', \beta'\} \in \text{dom}(p_\xi^\beta)$.

Write $n = f^r(\{\alpha', \beta'\})$.

(i) Assume on the contrary that there are $a \in T_{\alpha'}^r(n)$ and $b \in T_{\beta'}^r(n)$ with $U^r(a) \cap U^r(b) \neq \emptyset$.

First assume that $\{a, b\} \in [A^q]^2$. Since $q \in P$, we have $U^q(a) \cap U^q(b) = \emptyset$. So $t \in U^r(a) \cap U^r(b)$ should hold.

If $c \in A_\xi^\alpha$, then $t \in U(c)$ implies $y_\zeta \in U(c)$ by 7.1. Similarly, if $c \in A_\xi^\beta$, then $t \in U(c)$ implies $w_\xi \in U(c)$ by 7.2.

Since $U^q(a) \cap U^q(b) = \emptyset$, we can assume that $a \in A_\zeta^\alpha \setminus A_\xi^\beta$ and $b \in A_\xi^\beta \setminus A_\zeta^\alpha$.

But then $\alpha' \in \text{supp}(p_\zeta^\alpha) \setminus S$ and $\beta' \in \text{supp}(p_\xi^\beta) \setminus S$, so $f^r(\alpha', \beta')$ is undefined. Contradiction.

So we can assume that, e.g., $t = a$ and $b \in A^q$. Assume first that $b \in A^{p_\zeta^\alpha}$. Then $\alpha' = \alpha$ and $y_\zeta \in A_\zeta^\alpha$ by (7.1). Thus $y_\zeta \in T_\alpha^{p_\zeta^\alpha}(< n) \cap U^{p_\zeta^\alpha}(b)$, and so $T_\alpha^{p_\zeta^\alpha}(< n) \cap U[T_{\beta'}^{p_\zeta^\alpha}(n)] \neq \emptyset$, so (P4)(b) fails for p_ζ^α .

If $b \in A_\xi^\beta$, then we can use similar arguments using (7.2) instead of (7.1).

(ii) Assume on the contrary that there are $a \in T_\alpha^r(n)$ and $b \in T_{\beta'}^r(< n) \cap U^r(a)$.

Clearly $a \neq t$. If $b \neq t$, then $a \in T_{\alpha'}^q(n)$ and $b \in T_{\beta'}^q(< n) \cap U^q(a)$ which contradicts $q \in P$.

Assume that $b = t$. If $b \in A^{p_\zeta^\alpha}$, then (7.1) implies $\beta' = \alpha$ and $y_\zeta \in U^q(a) \cap T^q(< n)$. Thus $y_\zeta \in T_{\beta'}^q(< n) \cap U^q(a)$, which contradicts $q \in P$.

If $b \in A^{p_\xi^\beta}$, then we can use similar arguments using (7.2) instead of (7.1).

(P5) Let $\langle x, \gamma \rangle \in \text{dom}(g^r)$ and $y \in T_\gamma^r(g(x, \gamma))$. Since $U^r(t) = \{t\}$, we can assume that $x, y \neq t$. So $x, y \in A^q$. If $U^q(y) \subset U^q(x)$, then $x \preceq^q y$ and so $U^r(y) \subset U^r(x)$. Assume on the contrary that $U^q(x) \cap U^q(y) = \emptyset$, but $t \in U^r(x) \cap U^r(y)$. We can assume that $\langle x, \gamma \rangle \in g^{p_\zeta^\alpha}$. Thus $x \in A_\zeta^\alpha$ and $\gamma \in I_\zeta^\alpha$. However $T_\gamma^q \subset A_\zeta^\alpha$, so $y \in A_\zeta^\alpha$. Since $x, y \in A_\zeta^\alpha$ and $\gamma \in I_\zeta^\alpha$, $t \in U^r(x) \cap U^r(y)$ implies $y_\zeta \in U^{p_\zeta^\alpha}(x) \cap U^{p_\zeta^\alpha}(y)$ by (7.1), which contradicts $U^q(x) \cap U^q(y) = \emptyset$.

So we proved $r \in P$.

Next we show that $r \leq p_\zeta^\alpha, p_\xi^\beta$. (O1)–(O4) are trivial. To check (O5), assume on the contrary that $U^{p_\zeta^\alpha}(a) \cap U^{p_\zeta^\alpha}(b) = \emptyset$, but $U^r \cap U^r(b) \neq \emptyset$.

Then $t \in U^r(a) \cap U^r(b)$, and so $y_\zeta^\alpha \in U^{p_\zeta^\alpha}(a) \cap U^{p_\zeta^\alpha}(b)$ by (7.1), which is a contradiction.

Finally, it is also straightforward that

$$(7.3) \quad r \Vdash (\text{G5})(i)\text{--}(ii) \text{ holds for } \alpha, \beta, \zeta, \xi, \text{ and } t.$$

So we proved the theorem. □

8. Open problems

In this section, we present a list of open problems which could be of further interest and are closely connected to our results.

Problem 8.1. *Is every linearly ordered space base resolvable?*

Problem 8.2. *Is every T_3 (hereditarily) separable space base resolvable?*

Problem 8.3. *Is every paracompact space base resolvable?*

Note that under PFA, every T_3 hereditarily separable space is Lindelöf hence base resolvable by Corollary 3.7. Also, we conjecture that our forcing construction can be modified to produce a separable non base resolvable space.

Problem 8.4. *Is every power of \mathbb{R} base resolvable? Is it true that base resolvability is preserved by products?*

We know that every π -base is the union of two disjoint π -bases by Proposition 2.5(2). However:

Problem 8.5. *Does every base contain a disjoint base and π -base?*

Bases closed under finite unions are resolvable by Corollary 4.9 which raises to following question:

Problem 8.6. *Is it true that every base which is closed under finite intersections is base resolvable?*

It would be interesting to look into the following:

Problem 8.7. *Is every self-filling family \mathcal{F} of closed (Borel) sets of ω^ω resolvable?*

Concerning negligible subsets we ask the following:

Problem 8.8. *Is there a base \mathbb{B} for some space X such that every $\mathcal{U} \in [\mathbb{B}]^{|\mathbb{B}|}$ contains a neighborhood base at some point?*

REFERENCES

- [1] Hajnal A., Hamburger P., *Set Theory*, London Mathematical Society Student Texts, 48, Cambridge University Press, Cambridge, 1999, ISBN 0 521 59667 X.
- [2] Stone A.H., *On partitioning ordered sets into cofinal subsets*, *Mathematika* **15** (1968), 217–222.

UNIVERSITY OF TORONTO

E-mail: daniel.soukup@mail.utoronto.ca

URL: <http://math.utoronto.ca/~dsoukup>

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES,
BUDAPEST, HUNGARY

E-mail: soukup@renyi.hu

URL: <http://www.renyi.hu/~soukup>

(Received March 28, 2013, revised January 20, 2014)