

Alireza Olfati

On atomic ideals in some factor rings of $C(X, \mathbb{Z})$

Commentationes Mathematicae Universitatis Carolinae, Vol. 62 (2021), No. 2, 259–263

Persistent URL: <http://dml.cz/dmlcz/149015>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2021

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

On atomic ideals in some factor rings of $C(X, \mathbb{Z})$

ALIREZA OLFATI

Abstract. A nonzero R -module M is atomic if for each two nonzero elements a, b in M , both cyclic submodules Ra and Rb have nonzero isomorphic submodules. In this article it is shown that for an infinite P -space X , the factor rings $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ and $C_c(X)/C_F(X)$ have no atomic ideals. This fact generalizes a result published in paper by A. Mozaffarikhah, E. Momtahan, A. R. Olfati and S. Safaeyan (2020), which says that for an infinite set X , the factor ring $\mathbb{Z}^X/\mathbb{Z}^{(X)}$ has no atomic ideal. Another result is that for each infinite P -space X , the socle of the factor ring $C_c(X)/C_F(X)$ is always equal to zero. Also, zero-dimensional spaces X are characterized for which $C^F(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ have atomic ideals.

Keywords: P -space; rings of integer-valued continuous functions; functionally countable subalgebra; atomic ideal; socle

Classification: 54C40

1. Introduction

In this article, by ring, we mean a commutative ring with identity. A submodule N of an R -module M is said to be essential in M and denoted by $N \leq_{\text{ess}} M$ if N intersects each nonzero submodule of M nontrivially. The socle of a module M is the sum of all minimal submodules of M . For every nonzero module M , the socle of M is equal to the intersection of all essential submodules of M .

We assume throughout this paper that any topological space X is Tychonoff. By $C(X)$, we mean the ring of all real valued continuous functions on X . For any $f \in C(X)$, the set $Z(f) := \{x \in X : f(x) = 0\}$ is called the zero-set of f . For any Tychonoff space X , we denote by $C(X, \mathbb{Z})$, the set of all integer valued continuous functions on X . The set of all continuous real valued functions on X with countable image is denoted by $C_c(X)$. Clearly $C(X, \mathbb{Z})$ is a proper subset of $C_c(X)$. Equipped with the pointwise addition and multiplication of \mathbb{R}^X , the sets $C(X, \mathbb{Z})$ and $C_c(X)$ form two subrings of $C(X)$. The ring of integer-valued continuous functions have been studied in many ways and some outstanding results were achieved by some mathematicians, see for example [1], [6], [9] and [10]. Just recently in [2] an extensive study for the subalgebra $C_c(X)$ has been done. In

[5] it is shown that the socle of $C(X)$ is equal to the set of all functions $f \in C(X)$ such that $X \setminus Z(f)$ is finite. We denote the socle of $C(X)$ by $C_F(X)$. It is obvious to see that $C_F(X)$ is a subset of $C_c(X)$. In [3], it was shown that the socle of the factor ring $C(X)/C_F(X)$ is always equal to zero. We also denote by $C_F(X, \mathbb{Z})$ the set $C_F(X) \cap C(X, \mathbb{Z})$. The notion of atomic submodules was introduced and studied in [7] as a generalization for minimal submodules. In [8] it is shown that for an arbitrary infinite set X , the factor ring $\mathbb{Z}^X/\mathbb{Z}^{(X)}$ has no atomic ideal. In this note, we generalize this fact to every P -space and show that for an infinite P -space X the factor rings $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ and $C_c(X)/C_F(X)$ have no atomic ideals. This result leads to the fact that such factor rings have no uniform ideals, i.e., the socle of them equals zero.

2. Main results

Definition 2.1. A nonzero R -module M is called *atomic* if for each a, b in $M \setminus (0)$, the cyclic submodules Ra and Rb have nonzero isomorphic submodules.

Recall that a uniform module is a nonzero module A such that the intersection of any two nonzero submodules of A is nonzero, or, equivalently, such that every nonzero submodule of A is essential in A . It is obvious that every uniform module is atomic. Since every minimal ideal in a semiprime ring is uniform, we observe that every minimal ideal is atomic but the converse is not true. For example, if X is a zero-dimensional space, $C(X, \mathbb{Z})$ has no minimal ideals, but it has an atomic ideal if and only if X has an isolated point. For every module M , we denote by $\Sigma_a(M)$, the sum of all atomic submodules of M . For example, in [8] it is shown that for every zero-dimensional space X , $\Sigma_a(C(X, \mathbb{Z})) = C_F(X, \mathbb{Z})$ and for every Tychonoff space X , $\Sigma_a(C(X)) = C_F(X)$. For every nonzero module M , the factor module $M/\Sigma_a(M)$ may or may not have any atomic submodules, for example see [8]. But it is shown in [8] that for every infinite set X , the factor ring $\mathbb{Z}^X/\mathbb{Z}^{(X)}$ has no atomic ideals, or equivalently $\Sigma_a(\mathbb{Z}^X/\mathbb{Z}^{(X)})$ is equal to zero. The main objective in the sequel is to extend this result to every P -space and show that the two factor rings $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ and $C_c(X)/C_F(X)$ have no atomic ideals. But first we need the following two results.

Proposition 2.2. *If R is a semiprime ring, the ideal I of R is an atomic R -module if and only if I is a uniform ideal.*

PROOF: Let I be an atomic ideal of R . Then for every two $a, b \in I \setminus (0)$, there exist two nonzero submodules M of Ra and N of Rb and an isomorphism $\varphi: M \rightarrow N$. Then for some $r, s \in R$, $\varphi(ra) = sb \neq 0$. Since R is semiprime and φ is a homomorphism, $g\varphi(rf) = \varphi(rgf)$ is nonzero and therefore $fg \neq 0$.

For the converse, suppose that $a, b \in I \setminus (0)$. Since $ab \neq 0$, the cyclic submodule Rab is contained in $Ra \cap Rb$. Thus Ra and Rb have nonzero isomorphic submodules. \square

We recall that a topological space X is a P -space if and only if for each countable family of open sets $\{U_n : n \in \mathbb{N}\}$, the subset $\bigcap_{n \in \mathbb{N}} U_n$ is open in X .

Proposition 2.3. *Every infinite P -space X has a countable infinite family $\{C_i : i \in \mathbb{N}\}$ of nonempty clopen subsets, such that $X = \bigcup_{i \in \mathbb{N}} C_i$ and for each pair of distinct $i, j \in \mathbb{N}$, $C_i \cap C_j = \emptyset$.*

PROOF: Suppose that X is an infinite P -space. By [4, Exercise 4K.2], X is not pseudocompact. Hence there exists a continuous real valued function $f : X \rightarrow \mathbb{R}$ such that $f(X)$ is unbounded and hence infinite. For each $r \in f(X)$, the inverse image $f^{-1}(r)$ is clopen. Choose a countable subset $\{r_n : n \in \mathbb{N}\}$ of $f(X)$. Since two sets $T = \bigcup_{n \in \mathbb{N}} f^{-1}(r_n)$ and $C_0 = f^{-1}(\mathbb{R}) \setminus T$ are open, the subset C_0 is clopen. Now the family $\{C_i : i \in \mathbb{N} \cup \{0\}\}$ is an infinite clopen partition for X . \square

Theorem 2.4. *Assume that X is an infinite P -space. Then both the factor rings $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ and $C_c(X)/C_F(X)$ have no atomic ideals.*

PROOF: Assume that $I/C_F(X, \mathbb{Z})$ is an ideal of $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ and argue by contraposition that it is not an atomic ideal. Let $\bar{f} = f + C_F(X, \mathbb{Z})$ and $\bar{g} = g + C_F(X, \mathbb{Z})$ be two nonzero elements of $I/C_F(X, \mathbb{Z})$ such that $\bar{f}\bar{g} \neq 0$. This means that $fg \notin C_F(X, \mathbb{Z})$ and hence $X \setminus Z(fg)$ is infinite. Since $X \setminus Z(fg)$ is an infinite P -space, by Theorem 2.4, there exists a countably infinite clopen partition $\{C_i : i \in \mathbb{N}\}$ of $X \setminus Z(fg)$. From the fact that $X \setminus Z(fg)$ is clopen in X , each C_i is clopen in X . It is easy to see that the sets $T = \bigcup_{i \in \mathbb{N}} C_{2i}$ and $S = \bigcup_{i \in \mathbb{N}} C_{2i-1}$ are two clopen subsets of X . Let χ_T and χ_S be the characteristic functions of T and S , respectively. Define $h = \chi_T f$ and $k = \chi_S g$. Clearly, $\bar{h} = h + C_F(X, \mathbb{Z})$ and $\bar{k} = k + C_F(X, \mathbb{Z})$ are two nonzero elements of the ideal $I/C_F(X, \mathbb{Z})$ and $\bar{h}\bar{k} = 0$, a contradiction.

The proof for the factor ring $C_c(X)/C_F(X)$ can be repeated verbatim without any extra work. \square

Since each minimal ideal of a commutative ring is atomic, the following corollary is immediate.

Corollary 2.5. *For each infinite P -space X , we have that the socle of the factor ring $C_c(X)/C_F(X)$ is equal to zero.*

In the sequel we observe a necessary and sufficient condition for a compact zero-dimensional space X such that the factor ring $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ has a uniform and hence an atomic ideal. First, we need the following result from [8].

Proposition 2.6. *Let X be a zero-dimensional space. An ideal I of $C(X, \mathbb{Z})$ is atomic if and only if it is a principal ideal generated by the characteristic function of an isolated point of X .*

Corollary 2.7. *Let X be a zero-dimensional space. For the ring $C(X, \mathbb{Z})$,*

$$\Sigma_a(C(X, \mathbb{Z})) = C_F(X, \mathbb{Z}).$$

The set of all isolated points of a topological space X is denoted by $\mathbb{I}(X)$.

Proposition 2.8. *Let X be a compact zero-dimensional space. The factor ring $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ has a uniform (or equivalently atomic) ideal if and only if the subset $X \setminus \mathbb{I}(X)$ has an isolated point.*

PROOF: Let X be a compact zero-dimensional space. Assume that the factor ring $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ has a uniform ideal. Since by [9, Proposition 4.11], the factor ring $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ is isomorphic with the ring $C(X \setminus \mathbb{I}(X), \mathbb{Z})$, it follows that the ring $C(X \setminus \mathbb{I}(X), \mathbb{Z})$ has an atomic ideal and hence by Proposition 2.6, the subspace $X \setminus \mathbb{I}(X)$ has an isolated point. \square

The set of all continuous integer-valued functions with finite images is denoted by $C^F(X, \mathbb{Z})$. We remind the reader that the Banaschewski compactification of a zero-dimensional Hausdorff space X is a compact Hausdorff space $\beta_0 X$ which contains X as a dense subspace and each continuous real valued function $f : X \rightarrow \mathbb{R}$ with a finite image has an extension to $\beta_0 X$; see, e.g. [2].

Proposition 2.9. *Let X be an arbitrary zero-dimensional space. The factor ring $C^F(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ has a uniform (or equivalently atomic) ideal if and only if the subset $\beta_0 X \setminus \mathbb{I}(X)$ has an isolated point.*

PROOF: For each $f \in C^F(X, \mathbb{Z})$, there is a unique $f^{\beta_0} \in C^F(\beta_0 X, \mathbb{Z}) = C(\beta_0 X, \mathbb{Z})$, such that $f^{\beta_0}|_X = f$ and $f^{\beta_0}(X) = f(X)$. Note that since X is dense in $\beta_0 X$, $\mathbb{I}(\beta_0 X) = \mathbb{I}(X)$. Clearly under the isomorphism $f \rightarrow f^{\beta_0}$, $C_F(X, \mathbb{Z})$ is sent to $C_F(\beta_0 X, \mathbb{Z})$. Hence $C^F(X, \mathbb{Z})/C_F(X, \mathbb{Z}) \cong C(\beta_0 X, \mathbb{Z})/C_F(\beta_0 X, \mathbb{Z})$. Now by Proposition 2.8, $C^F(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ has a uniform (or equivalently atomic) ideal if and only if $\beta_0 X \setminus \mathbb{I}(\beta_0 X) = \beta_0 X \setminus \mathbb{I}(X)$ has an isolated point. \square

Example 2.10. It is well known that $\beta\mathbb{N} \setminus \mathbb{N}$ has no isolated points. Now Proposition 2.9 implies that $C^F(\mathbb{N}, \mathbb{Z})/C_F(\mathbb{N}, \mathbb{Z})$ has no uniform (or equivalently atomic) ideals.

With regard to the latter proposition, the interested reader is encouraged to characterize all zero-dimensional topological spaces X for which the factor ring $C(X, \mathbb{Z})/C_F(X, \mathbb{Z})$ has an atomic (equivalently, uniform) ideal, a question which is unsettled yet.

REFERENCES

- [1] Alling N. L., *Rings of continuous integer-valued functions and nonstandard arithmetic*, Trans. Amer. Math. Soc. **118** (1965), 498–525.
- [2] Azarpanah F., Karamzadeh O. A. S., Keshtkar Z., Olfati A. R., *On maximal ideals of $C_c(X)$ and the uniformity of its localizations*, Rocky Mountain J. Math. **48** (2018), no. 2, 345–384.
- [3] Azarpanah F., Karamzadeh O. A. S., Rahmati S., *$C(X)$ vs. $C(X)$ modulo its socle*, Colloq. Math. **111** (2008), no. 2, 315–336.
- [4] Gillman L., Jerison M., *Rings of Continuous Functions*, Graduate Texts in Mathematics, 43, Springer, New York, 1976.
- [5] Karamzadeh O. A. S., Rostami M., *On the intrinsic topology and some related ideals of $C(X)$* , Proc. Amer. Math. Soc. **93** (1985), no. 1, 179–184.
- [6] Martinez J., *$C(X, \mathbb{Z})$ revisited*, Adv. Math. **99** (1993), no. 2, 152–161.
- [7] Momtahan E., Motamedi M., *A study on dimensions of modules*, Bull. Iranian. Math. Soc. **43** (2017), no. 5, 1227–1235.
- [8] Mozaffarikhah A., Momtahan E., Olfati A. R., Safaeeyan S., *p -semisimple modules and type submodules*, J. Algebra Appl. **19** (2020), no. 4, 2050078, 22 pages.
- [9] Olfati A. R., *Homomorphisms from $C(X, \mathbb{Z})$ into a ring of continuous functions*, Algebra Universalis **79** (2018), no. 2, Paper No. 34, 26 pages.
- [10] Pierce R. S., *Rings of integer-valued continuous functions*, Trans. Amer. Math. Soc. **100** (1961), 371–394.

A. Olfati:

DEPARTMENT OF MATHEMATICS, YASOUJ UNIVERSITY, DANESHJOO ST., YASOUJ,
7591874934, IRAN

E-mail: alireza.olfati@yu.ac.ir

E-mail: olfati.alireza@gmail.com

(Received October 1, 2019, revised June 11, 2020)