

Farid Bencherif; Rachid Boumahdi; Tarek Garici

Symmetric identity for polynomial sequences satisfying $A'_{n+1}(x) = (n + 1)A_n(x)$

Communications in Mathematics, Vol. 29 (2021), No. 3, 343–355

Persistent URL: <http://dml.cz/dmlcz/149321>

Terms of use:

© University of Ostrava, 2021

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Symmetric identity for polynomial sequences satisfying $A'_{n+1}(x) = (n + 1)A_n(x)$

Farid Bencherif, Rachid Boumahdi, Tarek Garici

Abstract. Using umbral calculus, we establish a symmetric identity for any sequence of polynomials satisfying $A'_{n+1}(x) = (n + 1)A_n(x)$ with $A_0(x)$ a constant polynomial. This identity allows us to obtain in a simple way some known relations involving Apostol-Bernoulli polynomials, Apostol-Euler polynomials and generalized Bernoulli polynomials attached to a primitive Dirichlet character.

1 Introduction and preliminaries

With each formal power series $S(z) = \sum_{k=0}^{\infty} a_k \frac{z^k}{k!} \in \mathbb{C}[[z]]$ we can associate a polynomial sequence $(A_k(x))_{k \geq 0}$ defined by

$$\sum_{k=0}^{\infty} A_k(x) \frac{z^k}{k!} = S(z)e^{xz}, \tag{1}$$

or equivalently by $A_k(x) = \sum_{i=0}^k \binom{k}{i} a_i x^{k-i}$. Clearly

$$A'_k(x) = kA_{k-1}(x), \quad k \geq 1, \quad A'_0(x) = 0. \tag{2}$$

Reciprocally, it is not difficult to prove that if a polynomial sequence $(A_k(x))_{k \geq 0}$ satisfies (2) then (1) holds for

$$S(z) = \sum_{k=0}^{\infty} A_k(0) \frac{z^k}{k!}.$$

2020 MSC: 05A19, 05A40, 11B68

Key words: Appell sequence, Apostol-Bernoulli polynomial, Apostol-Euler polynomial, generalized Bernoulli polynomial, primitive Dirichlet character.

Affiliation:

Farid Bencherif – Faculté de Mathématiques, USTHB, Alger, Algérie

E-mail: fbencherif@usthb.dz

Rachid Boumahdi – Ecole Nationale Supérieure d’Informatique, Alger, Algérie

E-mail: r_boumehdi@esi.dz

Tarek Garici – Faculté de Mathématiques, USTHB, Alger, Algérie

E-mail: tgarici@usthb.dz

Apostol-Bernoulli polynomials $(\mathcal{B}_k(x; \lambda))_{k \geq 0}$ and Apostol-Euler polynomials $(\mathcal{E}_k(x; \lambda))_{k \geq 0}$ in the variable x defined for $\lambda \in \mathbb{C}$ by

$$\frac{z}{\lambda e^z - 1} e^{xz} = \sum_{k=0}^{\infty} \mathcal{B}_k(x; \lambda) \frac{z^k}{k!}$$

and for $\lambda \in \mathbb{C} \setminus \{-1\}$ by

$$\frac{2}{\lambda e^z + 1} e^{xz} = \sum_{k=0}^{\infty} \mathcal{E}_k(x; \lambda) \frac{z^k}{k!}$$

provide an example of such polynomials. If χ is a primitive Dirichlet character with conductor $f = f_\chi$, then $B_{n,\chi}(x)$, $n = 0, 1, 2, \dots$, the generalized Bernoulli polynomials attached to χ are defined by the generating function [1]

$$F_\chi(t, x) = \frac{t}{e^{ft} - 1} \sum_{a=1}^f \chi(a) e^{at} e^{xt} = \sum_{n=0}^{\infty} B_{n,\chi}(x) \frac{t^n}{n!}, \quad |t| < \frac{2\pi}{f}.$$

The polynomial sequence $(B_{n,\chi}(x))_{n \geq 0}$ is another example of polynomial sequence satisfying (1). Furthermore, if $a_0 \neq 0$, the polynomial sequence $(A_k(x))_{k \geq 0}$ is called an Appell sequence [2]. Which is the case for the classical Bernoulli and Euler polynomials $B_k(x)$ and $E_k(x)$ defined respectively by

$$\frac{z}{e^z - 1} e^{xz} = \sum_{k=0}^{\infty} B_k(x) \frac{z^k}{k!}$$

and

$$\frac{2}{e^z + 1} e^{xz} = \sum_{n=0}^{\infty} E_k(x) \frac{z^k}{k!}.$$

In 2003, motivated by the work of Kaneko [8], Momiyama [10] and Wu et al. [17], Sun [14] derived a general combinatorial identity in terms of polynomials with dual sequences of coefficients. More precisely, he considered the two following polynomial sequences

$$R_k(x) = \sum_{i=0}^k \binom{k}{i} (-1)^i u_i x^{k-i} \quad \text{and} \quad R_k^*(x) = \sum_{i=0}^k \binom{k}{i} (-1)^i u_i^* x^{k-i}, \quad (3)$$

where $(u_n)_{n \geq 0}$ is any complex sequence and $(u_n^*)_{n \geq 0}$ is its dual sequence defined for $k \geq 0$ by $u_k^* = \sum_{i=0}^k \binom{k}{i} (-1)^i u_i$, and proved [14, Theorem 1.1], for any integers $n, m \geq 0$ and $x + y + z = 1$, the following identities

$$\begin{aligned} & (-1)^{n-1} \sum_{k=0}^m \binom{m}{k} x^{m-k} \frac{R_{n+k+1}(y)}{n+k+1} \\ & + (-1)^{m+1} \sum_{k=0}^n \binom{n}{k} x^{n-k} \frac{R_{m+k+1}^*(z)}{m+k+1} = \frac{m!n!x^{m+n+1}u_0}{(m+n+1)!}, \quad (4) \end{aligned}$$

$$(-1)^{n-1} \sum_{k=0}^m \binom{m}{k} x^{m-k} R_{n+k}(y) + (-1)^m \sum_{k=0}^n \binom{n}{k} x^{n-k} R_{m+k}^*(z) = 0, \quad (5)$$

$$\begin{aligned} & (-1)^n \left\{ \sum_{k=0}^m \binom{m+1}{k} x^{m+1-k} (n+1+k) R_{n+k}(y) \right. \\ & \qquad \qquad \qquad \left. + (n+m+2) R_{n+m+1}(y) \right\} \\ & + (-1)^m \left\{ \sum_{k=0}^n \binom{n+1}{k} x^{n+1-k} (m+1+k) R_{m+k}^*(z) \right. \\ & \qquad \qquad \qquad \left. + (n+m+2) R_{m+n+1}^*(z) \right\} = 0. \quad (6) \end{aligned}$$

Which allowed him to derive various known identities involving Bernoulli numbers.

Chen and Sun [3], by applying the extended Zeilberger’s algorithm, established several recurrence relations for Bernoulli numbers and polynomials which generalize the relations of Momiyama [10], Gessel [6] and Gelfand [5]. Prévost [11], by using the Padé approximation of the exponential function, extended Chen and Sun’s results and obtained numerous recurrence relations involving Apostol-Bernoulli polynomials $\mathcal{B}_k(x; \lambda)$ or Apostol-Euler polynomials $\mathcal{E}_k(x; \lambda)$. Recently, Agoh [1] found some shortened recurrence relations for generalized Bernoulli polynomials attached to a primitive Dirichlet character χ which allowed him to reprove and generalize several identities on classical Bernoulli numbers and polynomials such as Saalschütz-Gelfand [5], von Ettiinghausen-Seidel-Stern-Kaneko’s [15], [16], [13], [8] and Chen and Sun [3] formulas. In particular, he showed [1, Theorem 2.1] that for any integers $n, m \geq 0$ and $s \geq 1$, we have

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \frac{(sf)^{m-k} B_{n+1+k, \chi}(x)}{n+1+k} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{(sf)^{n-k} B_{m+1+k, \chi}(x)}{m+1+k} \\ & = \sum_{a=1}^f \chi(a) \sum_{r=0}^{s-1} (a+x+rf)^m (a+x+(r-s)f)^n \\ & \qquad \qquad \qquad + \frac{(-1)^{n+1}}{m+n+1} \binom{n+m}{n}^{-1} (sf)^{n+m+1} B_{0, \chi}(x). \quad (7) \end{aligned}$$

With the use of the well-known relation [1, Eq. (2.4)]

$$B_{n, \chi}(x+sf) = B_{n, \chi}(x) + \sum_{k=0}^{s-1} \sum_{a=1}^f \chi(a) n(a+x+kf)^{n-1}, \quad (8)$$

where n and s are non-negative integers, identity (7) can be deduced from the

following relation, due to He and Zhang [7, Theorem 1.2]

$$\sum_{k=0}^m \binom{m}{k} \frac{\alpha^{m-k} A_{n+k+q}(x)}{(n+q+k)^{\underline{q}}} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{\alpha^{n-k} A_{m+q+k}(x+\alpha)}{(m+q+k)^{\underline{q}}} = \frac{(-1)^{n+1} \alpha^{n+m+1}}{(q-1)!} \int_0^1 (1-t)^m t^n A_{q-1}(x+\alpha t) dt, \tag{9}$$

where $(A_n(x))_{n \geq 0}$ is any sequence of polynomials defined by relation (1), α a complex number, and $n, m \geq 0, q \geq 1$ any integers. Recall that the falling factorial $z^{\underline{q}}$ is a polynomial in z defined by $z^{\underline{0}} = 1$ and

$$z^{\underline{q}} = \prod_{j=0}^{q-1} (z-j),$$

for $q \geq 1$.

It is the main aim of this paper to establish in Theorem 1 a Saalschütz-Gelfand type identity for the polynomial sequence $(A_n(x))_{n \geq 0}$ satisfying (2). This identity allows us to reprove some known identities for generalized Bernoulli polynomials attached to a primitive Dirichlet character due to Agoh [1] and Apostol-Bernoulli polynomials due to Prévost [11, Theorem 2] in a simple way.

2 Main result

Before giving our main result it is convenient to have the following lemma.

Lemma 1. *For any non-negative integers n, m and q , we have the following identity*

$$\sum_{k=0}^m \binom{m}{k} \frac{z^{n+k+q}}{(n+k+q)^{\underline{q}}} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{(1+z)^{m+k+q}}{(m+k+q)^{\underline{q}}} = \frac{(-1)^{n+1}}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \frac{m!(n+q-k-1)!}{(m+n+q-k)!} z^k. \tag{10}$$

Proof. Consider the polynomial $P(z) = z^n(1+z)^m$ and let $Q(z)$ denote the q -fold primitive of $P(z)$ defined by

$$Q(z) = \int_0^z \int_0^{z_1} \cdots \int_0^{z_{q-1}} P(t) dt dz_{q-1} \cdots dz_1. \tag{11}$$

On the one hand, by the Cauchy well-known formula [4, p. 115] for repeated integration, we have

$$Q(z) = \frac{1}{(q-1)!} \int_0^z (z-t)^{q-1} P(t) dt. \tag{12}$$

Changing variable t into $u - 1$ in the right-hand side of (12) gives

$$\begin{aligned} Q(z) &= \frac{1}{(q-1)!} \int_0^{z+1} ((z+1) - u)^{q-1} u^m (u-1)^n du \\ &\quad - \frac{1}{(q-1)!} \int_0^1 (z + (1-u))^{q-1} u^m (u-1)^n du \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{1}{(q-1)!} \int_0^{z+1} ((z+1) - u)^{q-1} u^{m+k} du \\ &\quad - \frac{1}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} (-1)^n z^k \int_0^1 u^m (1-u)^{n+q-k-1} du. \end{aligned}$$

Recall that for any non-negative integers i and j ,

$$\int_0^1 u^i (1-u)^j du = \frac{i!j!}{(i+j+1)!}$$

and for any non-negative integer k , from the Cauchy formula (12) we have

$$\frac{1}{(q-1)!} \int_0^z (z-u)^{q-1} u^k du = \frac{z^{k+q}}{(k+q)^{\underline{q}}}. \tag{13}$$

From this, we can deduce that

$$\begin{aligned} Q(z) &= \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{(z+1)^{m+k+q}}{(m+k+q)^{\underline{q}}} \\ &\quad - \frac{1}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} (-1)^n \frac{m!(n+q-k-1)!}{(m+n+q-k)!} z^k. \end{aligned} \tag{14}$$

On the other hand, expanding $P(t)$ in (11) and using the well-known identity

$$\int_0^z \int_0^{z_1} \dots \int_0^{z_{q-1}} t^{n+k} dt dz_{q-1} \dots dz_1 = \frac{z^{n+k+q}}{(n+k+q)^{\underline{q}}},$$

where k is any non-negative integer, give

$$Q(z) = \sum_{k=0}^m \binom{m}{k} \frac{z^{n+k+q}}{(n+k+q)^{\underline{q}}}. \tag{15}$$

Finally, by equalling both expressions (14) and (15) of $Q(z)$ we obtain (10). \square

Theorem 1. Let $S(z) \in \mathbb{C}[[z]]$ be a formal power series and $(A_k(x))_{k \geq 0}$ be a sequence of polynomials given by

$$\sum_{k=0}^{+\infty} A_k(x) \frac{z^k}{k!} = S(z)e^{xz}. \tag{16}$$

For any complex number α and any integers $n, m \geq 0$ and $q \geq 1$, we have

$$\begin{aligned} \sum_{k=0}^m \binom{m}{k} \frac{\alpha^{m-k} A_{n+q+k}(x)}{(n+q+k)^{\underline{q}}} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{\alpha^{n-k} A_{m+q+k}(x+\alpha)}{(m+q+k)^{\underline{q}}} \\ = \frac{(-1)^{n+1}}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \alpha^{n+m+q-k} \frac{m!(n+q-k-1)!}{(m+n+q-k)!} A_k(x). \end{aligned} \tag{17}$$

Proof. Let

$$S(z) = \sum_{k=0}^{\infty} a_k \frac{z^k}{k!}.$$

Then, from (16) we have, for $r \geq 0$,

$$A_r(x) = \sum_{k=0}^r \binom{r}{k} a_{r-k} x^k.$$

Inspired by the work of Rota [12], let us consider the linear map L from $\mathbb{C}[x, y]$ to $\mathbb{C}[x]$ defined for any non-negative integers r and s by

$$L(x^r y^s) = a_s x^r. \tag{18}$$

For $\alpha = 0$, it is obvious that equation (17) holds. Assume that $\alpha \neq 0$. As for any $r \geq 0$ we have

$$\begin{aligned} L((x+y)^r) &= \sum_{i=0}^r \binom{r}{i} a_{r-i} x^i = A_r(x), \\ L((x+\alpha+y)^r) &= \sum_{i=0}^r \binom{r}{i} a_{r-i} (x+\alpha)^i = A_r(x+\alpha), \end{aligned}$$

replacing z with $(\frac{x+y}{\alpha})$ in (10) and applying L we get the desired relation (17). \square

Remark 1. If I and J denote respectively the right-hand sides of relations (9) and (17), then it is not difficult to show that $I = J$. In fact, for each fixed x and α , the Taylor expansion of the polynomial $A_{q-1}(x + \alpha t)$ in power of t is given by

$$A_{q-1}(x + \alpha t) = \sum_{k=0}^{q-1} \binom{q-1}{k} t^{q-1-k} \alpha^{q-1-k} A_k(x).$$

Thus,

$$J = \frac{\alpha^{n+m+1}}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \alpha^{q-1-k} A_k(x) \int_0^1 t^{n+q-1-k} (1-t)^m dt = I.$$

Which leads to another proof of Theorem 1.2 of He and Zhang [7].

3 Corollaries

Using Theorem 1, we can give an alternative proof of He and Zhang’s theorem [7, Theorem 1.1 and Corollary 1.2] as follows.

Corollary 1. *Let $S(z) \in \mathbb{C}[[z]]$ be a formal power series and $(A_k(x))_{k \geq 0}$ be a sequence of polynomials given by*

$$\sum_{k=0}^{+\infty} A_k(x) \frac{z^k}{k!} = S(z)e^{xz}.$$

For any complex number α and any non-negative integers n, m and q , we have

$$\sum_{k=0}^m \binom{m}{k} (n+k)^q \alpha^{m-k} A_{n-q+k}(x) - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} (m+k)^q \alpha^{n-k} A_{m-q+k}(x+\alpha) = 0. \quad (19)$$

Proof. From (2) we have for each $k \geq 1$, $A'_k(x) = kA_{k-1}(x)$ and $A'_0(x) = 0$. Taking $q = 1$ in relation (17) and then differentiating the obtained relation $q + 1$ times with respect to x gives (19). \square

Remark 2. Relation (19) is clearly satisfied for $\alpha = 0$. Note that in the case $\alpha \neq 0$ relation (19) can also be obtained by using the same method as in the proof of Theorem 1. Indeed, by differentiating the following identity

$$\sum_{k=0}^m \binom{m}{k} z^{n+k} = \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} (1+z)^{m+k}$$

q times with respect to z , we obtain

$$\sum_{k=0}^m \binom{m}{k} (n+k)^q z^{n-q+k} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} (m+k)^q (1+z)^{m-q+k} = 0.$$

Replacing z with $(\frac{x+y}{\alpha})$ and applying the linear map L defined by (18) we get (19).

Next applying Theorem 1 and Corollary 1 for $\alpha = sf$ to the sequence

$$(B_{n,\chi}(x))_{n \geq 0},$$

with the help of relation (8) we get the following two corollaries.

Corollary 2. *For integers $q, s \geq 1$, and $n, m \geq 0$, we have*

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \frac{(sf)^{m-k} B_{n+q+k,\chi}(x)}{(n+q+k)^q} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{(sf)^{n-k} B_{m+q+k,\chi}(x)}{(m+q+k)^q} \\ &= \sum_{i=0}^{s-1} \sum_{a=1}^f \chi(a) \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{(sf)^{n-k} (a+x+if)^{m+q-1+k}}{(m+q-1+k)^{q-1}} \\ &+ \frac{(-1)^{n+1}}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \frac{(sf)^{n+m+q-k} m!(n+q-k-1)!}{(m+n+q-k)!} B_{k,\chi}(x). \quad (20) \end{aligned}$$

With the help of relation (13), the right hand-side of (20) can also be written, for $q \geq 2$, as

$$\begin{aligned} & \frac{1}{(q-2)!} \sum_{i=0}^{s-1} \sum_{a=1}^f \chi(a) \int_0^{a+x+if} (a+x+if-u)^{q-2} u^m (u-sf)^n du \\ & + \frac{(-1)^{n+1}}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \frac{(sf)^{n+m+q-k} m!(n+q-k-1)!}{(m+n+q-k)!} B_{k,\chi}(x). \end{aligned}$$

Taking $q = 1$ in (20) gives immediately Theorem 2.1 of Agoh [1].

Corollary 3. For integers $s \geq 1$, and $n, m, q \geq 0$, we have

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} (n+k)^{\underline{q}} (sf)^{m-k} B_{n-q+k,\chi}(x) \\ & - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} (m+k)^{\underline{q}} (sf)^{n-k} B_{m-q+k,\chi}(x) \\ = & (q+1)! \sum_{a=1}^f \chi(a) \sum_{r=0}^{s-1} \sum_{j=0}^{q+1} \binom{m}{q+1-j} \binom{n}{j} (a+x+rf)^{m-q-1+j} (a+x+(r-s)f)^{n-j}. \end{aligned}$$

One observes that the left hand side of the previous identity can be expressed as

$$\frac{d^{q+1}}{dx^{q+1}} \sum_{a=1}^f \chi(a) \sum_{r=0}^{s-1} (a+x+rf)^m (a+x+(r-s)f)^n.$$

By taking respectively $q = 0$ and $q = 1$ in Corollary 3, we get Theorems 2.3 and 2.4 of Agoh [1].

The following corollary enables us to obtain Sun’s result by giving different values to q .

Corollary 4. Let $(R_k(x))_{n \geq 0}$ and $(R_k^*(x))_{n \geq 0}$ be the polynomial sequences defined by (3), then for any non-negative integers n, m and q and for any complex number α , we have

$$\begin{aligned} & (-1)^{n-1} \sum_{k=0}^m \binom{m}{k} \alpha^{m-k} \frac{R_{n+k+q}(x)}{(n+k+q)^{\underline{q}}} \\ & + (-1)^{m+q} \sum_{k=0}^n \binom{n}{k} \alpha^{n-k} \frac{R_{m+k+q}^*(1-x-\alpha)}{(m+k+q)^{\underline{q}}} \\ = & \frac{1}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \frac{m!(n+q-k-1)!}{(m+n+q-k)!} \alpha^{m+n+q-k} R_k(x), \quad (21) \end{aligned}$$

$$(-1)^{n-1} \sum_{k=0}^m \binom{m}{k} \alpha^{m-k} (n+k)^q R_{n-q+k}(x) + (-1)^{m-q} \sum_{k=0}^n \binom{n}{k} \alpha^{n-k} (m+k)^q R_{m-q+k}^*(1-x-\alpha) = 0. \quad (22)$$

Proof. If

$$F(z) = \sum_{k=0}^{\infty} u_k \frac{z^k}{k!}$$

denotes the exponential generating function for the sequence u , then it is easy to see that the polynomial sequences $(R_k(x))_{k \geq 0}$ and $(R_k^*(x))_{k \geq 0}$ satisfy

$$\sum_{k=0}^{+\infty} R_k(x) \frac{z^k}{k!} = F(-z)e^{zx} \quad \text{and} \quad \sum_{k=0}^{+\infty} R_k^*(x) \frac{z^k}{k!} = F(z)e^{z(x-1)}.$$

Thus, it comes immediately that

$$R_k(x) = (-1)^k R_k^*(1-x).$$

With this, a direct application of Theorem 1 and Corollary 1 for the polynomial sequence $(R_k(x))_{k \geq 0}$ gives (21) and (22). \square

For $q = 1$ in (21) and $q \in \{0, 1\}$ in (22) we obtain respectively Sun's results (4), (5), and (6).

To conclude this section, we deal with Apostol-Bernoulli polynomials $(\mathcal{B}_k(x; \lambda))_{k \geq 0}$ defined [9] for any complex number λ , by means of the following generating function

$$\frac{ze^{xz}}{\lambda e^z - 1} = \sum_{k=0}^{\infty} \mathcal{B}_k(x; \lambda) \frac{z^k}{k!},$$

with $|z + \ln \lambda| < 2\pi$. Let ℓ be a positive integer. Expanding the both sides of the functional relation

$$\frac{\lambda^\ell z e^{(\ell+x)z}}{\lambda e^z - 1} = \frac{z e^{xz}}{\lambda e^z - 1} + \sum_{i=0}^{\ell-1} \lambda^i z e^{iz} e^{xz}$$

into power series of z and then equating the coefficients of $\frac{z^k}{k!}$ ($k \geq 0$), yields

$$\lambda^\ell \mathcal{B}_0(x + \ell; \lambda) = \mathcal{B}_0(x; \lambda)$$

and

$$\lambda^\ell \mathcal{B}_k(x + \ell; \lambda) = \mathcal{B}_k(x; \lambda) + \sum_{i=0}^{\ell-1} \lambda^i k(x+i)^{k-1}$$

for $k \geq 1$. One can see that the previous equality also holds true even when $k = 0$. Therefore, for any positive integer ℓ and any non-negative integer k

$$\lambda^\ell \mathcal{B}_k(x + \ell; \lambda) = \mathcal{B}_k(x; \lambda) + \sum_{i=0}^{\ell-1} \lambda^i k(x+i)^{k-1}. \quad (23)$$

Theorem 1 enables us to obtain simply the following corollary due to Prévost [11, Theorem 2].

Corollary 5. For any integers $n, m, p \geq 0$ and $\ell \geq 1$, if $p \geq n + m + 1$ then

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \lambda^\ell \frac{(n+m-k)! \mathcal{B}_{p-k}(x; \lambda)}{(p-k)! \ell^{p-k}} - \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{(n+m-k)! \mathcal{B}_{p-k}(x; \lambda)}{(p-k)! \ell^{p-k}} \\ &= \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{(n+m-k)! (x+i)^{p-k-1}}{(p-k-1)! \ell^{p-k}} \\ & \quad - \lambda^\ell \frac{(-1)^n n! m!}{p!} \sum_{k=0}^{p-m-n-1} \binom{p}{k} \binom{p-m-k-1}{n} \frac{\mathcal{B}_k(x; \lambda)}{\ell^k}, \end{aligned} \tag{24}$$

and if $p \leq n + m$ then

$$\begin{aligned} & \sum_{k=0}^{\min\{m,p\}} \binom{m}{k} \lambda^\ell \frac{(n+m-k)! \mathcal{B}_{p-k}(x; \lambda)}{(p-k)! \ell^{p-k}} \\ & \quad - \sum_{k=0}^{\min\{n,p\}} \binom{n}{k} (-1)^k \frac{(n+m-k)! \mathcal{B}_{p-k}(x; \lambda)}{(p-k)! \ell^{p-k}} \\ &= \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^{\min(p-1,n)} \binom{n}{k} (-1)^k \frac{(n+m-k)! (x+i)^{p-k-1}}{(p-k-1)! \ell^{p-k}}. \end{aligned} \tag{25}$$

Proof. Applying Theorem 1 and Corollary 1 for $S(z) = \frac{z}{\lambda e^z - 1}$ and $\alpha = \ell$, multiplying the whole by λ^ℓ and using identity (23) give

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \frac{\lambda^\ell \mathcal{B}_{n+q+k}(x; \lambda)}{(n+q+k) \underline{q} \ell^{n+q+k}} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{\mathcal{B}_{m+q+k}(x; \lambda)}{(m+q+k) \underline{q} \ell^{m+q+k}} \\ &= \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \frac{(m+q+k)(x+i)^{m+q+k-1}}{(m+q+k) \underline{q} \ell^{m+q+k}} \\ & \quad + \lambda^\ell \frac{(-1)^{n+1}}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \frac{m!(n+q-k-1)! \mathcal{B}_k(x; \lambda)}{(n+m+q-k)! \ell^k}, \end{aligned}$$

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} (n+k) \underline{q} \frac{\lambda^\ell \mathcal{B}_{n-q+k}(x; \lambda)}{\ell^{n-q+k}} - \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} (m+k) \underline{q} \frac{\mathcal{B}_{m-q+k}(x; \lambda)}{\ell^{m-q+k}} \\ &= \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} (m+k) \underline{q} \frac{(m-q+k)(x+i)^{m-q+k-1}}{\ell^{m-q+k}}. \end{aligned}$$

Reverse the order of summations in the two previous relations to obtain

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \frac{\lambda^\ell \mathcal{B}_{n+m+q-k}(x; \lambda)}{(n+m+q-k)^\underline{q} \ell^{n+m+q-k}} \\ & \quad - \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{\mathcal{B}_{n+m+q-k}(x; \lambda)}{(n+m+q-k)^\underline{q} \ell^{m+n+q-k}} \\ & = \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{(n+m+q-k)(x+i)^{n+m+q-k-1}}{(n+m+q-k)^\underline{q} \ell^{n+m+q-k}} \\ & \quad + \lambda^\ell \frac{(-1)^{n+1}}{(q-1)!} \sum_{k=0}^{q-1} \binom{q-1}{k} \frac{m!(n+q-k-1)! \mathcal{B}_k(x; \lambda)}{(n+m+q-k)! \ell^k}, \quad (26) \end{aligned}$$

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} (n+m-k)^\underline{q} \frac{\lambda^\ell \mathcal{B}_{n+m-q-k}(x; \lambda)}{\ell^{n+m-q-k}} \\ & \quad - \sum_{k=0}^n \binom{n}{k} (-1)^k (n+m-k)^\underline{q} \frac{\mathcal{B}_{n+m-q-k}(x; \lambda)}{\ell^{n+m-q-k}} \\ & = \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^n \binom{n}{k} (-1)^k (n+m-k)^\underline{q} \frac{(n+m-q-k)(x+i)^{n+m-q-k-1}}{\ell^{n+m-q+k}}. \quad (27) \end{aligned}$$

Suppose first that $p \geq n+m+1$. Substituting q by $p-n-m$ in (26) gives

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \lambda^\ell \frac{(n+m-k)! \mathcal{B}_{p-k}(x; \lambda)}{(p-k)! \ell^{p-k}} - \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{(n+m-k)! \mathcal{B}_{p-k}(x; \lambda)}{(p-k)! \ell^{p-k}} \\ & = \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{(n+m-k)(x+i)^{p-k-1}}{(p-k-1)! \ell^{p-k}} \\ & \quad - \lambda^\ell \frac{(-1)^n n! m!}{p!} \sum_{k=0}^{p-n-m-1} \binom{p}{k} \binom{p-m-k-1}{n} \frac{\mathcal{B}_k(x; \lambda)}{\ell^k}. \end{aligned}$$

Thus, we obtain relation (24). Suppose now that $p \leq n+m$. Taking $q = n+m-p$ in (27) gives

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} (n+m-k)^{n+m-p} \frac{\lambda^\ell \mathcal{B}_{p-k}(x; \lambda)}{\ell^{p-k}} \\ & \quad - \sum_{k=0}^n \binom{n}{k} (-1)^k (n+m-k)^{n+m-p} \frac{\mathcal{B}_{p-k}(x; \lambda)}{\ell^{p-k}} \\ & = \sum_{i=0}^{\ell-1} \lambda^i \sum_{k=0}^n \binom{n}{k} (-1)^k (n+m-k)^{n+m-p} \frac{(p-k)(x+i)^{p-k-1}}{\ell^{p+k}}. \quad (28) \end{aligned}$$

Noting that the factor $(n+m-k)^{\frac{n+m-p}{p-k}}$ in the left hand side vanishes when $k > p$, and the factor $(n+m-k)^{\frac{n+m-p}{p-k}}$ in the right hand side vanishes when $k > p-1$, relation (28) leads to relation (25). \square

Remark 3. In a similar way, by making use of

$$\lambda^\ell \mathcal{E}_k(x+\ell; \lambda) = (-1)^\ell \mathcal{E}_k(x; \lambda) + 2(-1)^{\ell-1} \sum_{i=0}^{\ell-1} (-1)^i \lambda^i (x+i)^k,$$

we obtain an identity due to Prévost [11, Theorem 2], for Apostol-Euler polynomials $(\mathcal{E}_k(x; \lambda))_{k \geq 0}$ defined [9] for $\lambda \in \mathbb{C} \setminus \{-1\}$ by means of the following generating function

$$\frac{2}{\lambda e^z + 1} e^{xz} = \sum_{n=0}^{\infty} \mathcal{E}_n(x; \lambda) \frac{z^n}{n!} \quad (|z + \ln \lambda| < \pi).$$

References

- [1] T. Agoh: Shortened recurrence relations for generalized Bernoulli numbers and polynomials. *Journal of Number Theory* 176 (2017) 149–173.
- [2] P. Appell: Sur une classe de polynômes. *Annales Scientifiques de l'École Normale Supérieure* 9 (1880) 119–144.
- [3] W.Y.C. Chen, L.H. Sun: Extended Zeilberger's algorithm for identities on Bernoulli and Euler polynomials. *Journal of Number Theory* 129 (9) (2009) 2111–2132.
- [4] I.M. Gel'fand, G.E. Shilov: *Generalized Functions, Vol. 1: Properties and Operations*. Acad. Press, New York (1964).
- [5] M.B. Gelfand: A note on a certain relation among Bernoulli numbers. *Bašk. Gos. Univ. Ucen. Zap. Vyp* 31 (1968) 215–216.
- [6] I.M. Gessel: Applications of the classical umbral calculus. *Algebra Univers.* 49 (4) (2003) 397–434.
- [7] Y. He, W. Zhang: Some symmetric identities involving a sequence of polynomials. *The Electronic Journal of Combinatorics* 17 (N7) (2010) 7pp.
- [8] M. Kaneko: A recurrence formula for the Bernoulli numbers. *Proceedings of the Japan Academy, Series A, Mathematical Sciences* 71 (8) (1995) 192–193.
- [9] Q.-M. Luo, H.M. Srivastava: Some generalizations of the Apostol-Bernoulli and Apostol-Euler polynomials. *Journal of Mathematical Analysis and Applications* 308 (1) (2005) 290–302.
- [10] H. Momiyama: A new recurrence formula for Bernoulli numbers. *Fibonacci Quarterly* 39 (3) (2001) 285–288.
- [11] M. Prévost: Padé approximation and Apostol-Bernoulli and Apostol-Euler polynomials. *Journal of computational and applied mathematics* 233 (11) (2010) 3005–3017.
- [12] G.-C. Rota: The number of partitions of a set. *The American Mathematical Monthly* 71 (5) (1964) 498–504.
- [13] M. Stern: Beiträge zur Theorie der Bernoullischen und Eulerschen Zahlen. *Abhandlungen der Königlichen Gesellschaft der Wissenschaften in Göttingen* 26 (1880) 3–46.

- [14] Z.-W. Sun: Combinatorial identities in dual sequences. *European Journal of Combinatorics* 24 (6) (2003) 709–718.
- [15] A. von Ettingshausen: *Vorlesungen über die höhere Mathematik*. C. Gerold, Vienna (1827).
- [16] P.L. von Seidel: Über eine einfache Entstehungsweise der Bernoullischen Zahlen und einiger verwandten Reihen, *Sitzungsberichte der Münch. Akad. Math. Phys. Classe* 7 (1877) 157–187.
- [17] K.-J. Wu, Z.-W. Sun, H. Pan: Some identities for Bernoulli and Euler polynomials. *Fibonacci Quarterly* 42 (4) (2004) 295–299.

Received: 18 January 2019

Accepted for publication: 19 June 2020

Communicated by: Karl Dilcher