

Abdelmalek Azizi; Mohamed Mahmoud Chems-Eddin; Abdelkader Zekhnini  
Note on the Hilbert 2-class field tower

*Mathematica Bohemica*, Vol. 147 (2022), No. 4, 513–524

Persistent URL: <http://dml.cz/dmlcz/151096>

## Terms of use:

© Institute of Mathematics AS CR, 2022

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## NOTE ON THE HILBERT 2-CLASS FIELD TOWER

ABDELMALEK AZIZI, Oujda, MOHAMED MAHMOUD CHEMS-EDDIN, Fez,  
 ABDELKADER ZEKHINI, Oujda

Received April 27, 2021. Published online January 18, 2022.

Communicated by Clemens Fuchs

*Abstract.* Let  $k$  be a number field with a 2-class group isomorphic to the Klein four-group. The aim of this paper is to give a characterization of capitulation types using group properties. Furthermore, as applications, we determine the structure of the second 2-class groups of some special Dirichlet fields  $\mathbb{k} = \mathbb{Q}(\sqrt{d}, \sqrt{-1})$ , which leads to a correction of some parts in the main results of A. Azizi and A. Zekhini (2020).

*Keywords:* multiquadratic field; fundamental systems of units; 2-class group; 2-class field tower; capitulation

*MSC 2020:* 11R11, 11R16, 11R20, 11R27, 11R29, 11R37

## 1. INTRODUCTION

Let  $k$  be an algebraic number field and let  $\mathbb{C}\mathbb{L}_2(k)$  denote its 2-class group, that is the 2-Sylow subgroup of the ideal class group  $\mathbb{C}\mathbb{L}(k)$  of  $k$ . Denote by  $k^{(1)}$  the first Hilbert 2-class field of  $k$ , that is the maximal abelian unramified extension of  $k$  such that the degree  $[k^{(1)} : k]$  is a power of 2, and by  $k^{(2)}$  the Hilbert 2-class field of  $k^{(1)}$ . Let  $G_k = \text{Gal}(k^{(2)}/k)$  be the Galois group of  $k^{(2)}/k$  and  $G'_k$  be its derived subgroup. Then it is well known, by class field theory, that  $\text{Gal}(k^{(1)}/k) \simeq \mathbb{C}\mathbb{L}_2(k) \simeq G_k/G'_k$ .

The determination of the structure of  $G_k$  is a classical and difficult open problem of class field theory that is related to many other problems such as the capitulation and the length of the Hilbert 2-class field tower. Actually, our goal in the present paper is to investigate these problems for fields with 2-class groups of type  $(2, 2)$ . Note that if  $\mathbb{C}\mathbb{L}_2(k)$  is of type  $(2, 2)$ , the Hilbert 2-class field tower of  $k$  terminates in at most two steps and the structure of  $G_k$  is based on the capitulation problem in unramified quadratic extensions of  $k$ . In fact,  $G_k$  is isomorphic to one of the groups  $A$ ,  $Q_m$ ,  $D_m$  or  $S_m$ , where  $A$  is the Klein four-group, and  $Q_m$ ,  $D_m$ , or  $S_m$  denote the

quaternion, dihedral or semidihedral groups, respectively, of order  $2^m$ , with  $m \geq 3$  and  $m \geq 4$  for  $S_m$  (cf. [13]).

In this paper, we give a characterization of the capitulation types (see Table 2) using some group properties and as an application, we determine the structure of the second 2-class groups of some special Dirichlet fields  $\mathbb{k} = \mathbb{Q}(\sqrt{d}, \sqrt{-1})$ .

If  $k$  is a number field, we use the following notations:

$h_2(k)$ : the 2-class number of  $k$ ,

$h_2(d)$ : the 2-class number of the quadratic field  $\mathbb{Q}(\sqrt{d})$ ,

$\varepsilon_d$ : the fundamental unit of the quadratic field  $\mathbb{Q}(\sqrt{d})$ ,

$E_k$ : the unit group of  $k$ ,

FSU: abbreviation of “fundamental system of units”,

$k^{(1)}$ : the Hilbert 2-class field of  $k$ ,

$k^{(2)}$ : the Hilbert 2-class field of  $k^{(1)}$ ,

$G_k$ : the Galois group of  $k^{(2)}/k$ ,

$k^+$ : the maximal real subfield of  $k$ ,

$q(k) = [E_k : \prod_i E_{k_i}]$ : the unit index of  $k$ , if  $k$  is multiquadratic and  $k_i$  are the quadratic subfields of  $k$ ,

$N_{k'/k}$ : the norm map of an extension  $k'/k$ .

## 2. PRELIMINARIES

Let  $Q_m$ ,  $D_m$ , and  $S_m$  denote the quaternion, dihedral, and semidihedral groups, respectively, of order  $2^m$ , where  $m \geq 3$  and  $m \geq 4$  for  $S_m$ ; in addition let  $A$  be the Klein four-group. Each of these groups is generated by two elements  $x$  and  $y$ , and admits the following presentations:

$$\begin{aligned} x^2 = y^2 = 1, \quad y^{-1}xy = x & \quad \text{for } A, \\ x^{2^{m-2}} = y^2 = a, \quad a^2 = 1, \quad y^{-1}xy = x^{-1} & \quad \text{for } Q_m, \\ x^{2^{m-1}} = y^2 = 1, \quad y^{-1}xy = x^{-1} & \quad \text{for } D_m, \\ x^{2^{m-1}} = y^2 = 1, \quad y^{-1}xy = x^{2^{m-2}-1} & \quad \text{for } S_m. \end{aligned}$$

We recall some well known properties of 2-groups  $G_k$  such that  $G_k/G'_k$  is of type  $(2, 2)$ , where  $G'_k$  denotes the commutator subgroup of  $G_k$ . For more details about these properties, we refer the reader to [13], pages 272–273, [7], pages 1467–1469, and [9], Chapter 5.

Let  $x$  and  $y$  be as above. Note that the commutator subgroup  $G'_k$  of  $G$  is always cyclic and  $G'_k = \langle x^2 \rangle$ . The group  $G_k$  possesses exactly three subgroups of index 2, which are,

$$H_1 = \langle x \rangle, \quad H_2 = \langle x^2, y \rangle, \quad H_3 = \langle x^2, xy \rangle.$$

Note also that for the two cases  $Q_3$  and  $A$ , each  $H_i$  is cyclic. For the case  $D_m$  with  $m > 3$ ,  $H_2$  and  $H_3$  are also dihedral. For  $Q_m$  with  $m > 3$ ,  $H_2$  and  $H_3$  are quaternion. Finally for  $S_m$ ,  $H_2$  is dihedral whereas  $H_3$  is quaternion. Furthermore, if  $G_k$  is isomorphic to  $A$  (or  $Q_3$ ), then the subgroups  $H_i$  are cyclic of order 2 (or 4, respectively). If  $G_k$  is isomorphic to  $Q_m$  with  $m > 3$ ,  $D_m$  with  $m > 3$  or  $S_m$ , then  $H_1$  is cyclic and  $H_i/H'_i$  is of type  $(2, 2)$  for  $i \in \{2, 3\}$ , where  $H'_i$  is the commutator subgroup of  $H_i$ .

Let  $F_i$  be the subfield of  $k^{(2)}$  fixed by  $H_i$ , where  $i \in \{1, 2, 3\}$ . If  $k^{(2)} \neq k^{(1)}$ ,  $\langle x^4 \rangle$  is the unique subgroup of  $G'_k$  of index 2. Let  $L$  ( $L$  is defined only if  $k^{(2)} \neq k^{(1)}$ ) be the subfield of  $k^{(2)}$  fixed by  $\langle x^4 \rangle$ . Then  $F_1$ ,  $F_2$  and  $F_3$  are the three quadratic subextensions of  $k^{(1)}/k$  and  $L$  is the unique subfield of  $k^{(2)}$  such that  $L/k$  is a nonabelian Galois extension of degree 8.

Let us recall the definition of Taussky's conditions A and B. Let  $k'$  be a cyclic unramified extension of a number field  $k$  and  $j$  denotes the basic homomorphism  $j_{k'/k} : \mathbb{C}\mathbb{L}(k) \rightarrow \mathbb{C}\mathbb{L}(k')$ , induced by the extension of ideals from  $k$  to  $k'$ . Thus, we say:

- ▷  $k'/k$  satisfies condition A if and only if  $|\ker(j_{k'/k}) \cap N_{k'/k}(\mathbb{C}\mathbb{L}(k'))| > 1$ .
- ▷  $k'/k$  satisfies condition B if and only if  $|\ker(j_{k'/k}) \cap N_{k'/k}(\mathbb{C}\mathbb{L}(k'))| = 1$ .

Set  $j_{F_i/k} = j_i$ ,  $i = 1, 2, 3$ . Then we have:

**Theorem 2.1** ([13], Theorem 2).

- (1) If  $k^{(1)} = k^{(2)}$ , then all  $F_i$  satisfy condition A,  $|\ker(j_i)| = 4$  for  $i = 1, 2, 3$  and  $G_k$  is abelian of type  $(2, 2)$ .
- (2) If  $\text{Gal}(L/k) \simeq Q_3$ , then all  $F_i$  satisfy condition A and  $|\ker(j_i)| = 2$  for  $i = 1, 2, 3$  and  $G_k \simeq Q_3$ .
- (3) If  $\text{Gal}(L/k) \simeq D_3$ , then  $F_2, F_3$  satisfy condition B and  $|\ker j_2| = |\ker j_3| = 2$ . Furthermore, if  $F_1$  satisfies condition B, then  $|\ker j_1| = 2$  and  $G_k \simeq S_m$ , if  $F_1$  satisfies condition A and  $|\ker j_1| = 2$ , then  $G_k \simeq Q_m$ . If  $F_1$  satisfies condition A and  $|\ker j_1| = 4$ , then  $G_k \simeq D_m$ .

We summarize these results in Table 1.

$ \ker(j_1) $	(A/B)	$ \ker(j_2) $	(A/B)	$ \ker(j_3) $	(A/B)	$G_k$
4	A	4	A	4	A	$(2, 2)$
2	A	2	A	2	A	$Q_3$
4	A	2	B	2	B	$D_m, m \geq 3$
2	A	2	B	2	B	$Q_m, m > 3$
2	B	2	B	2	B	$S_m, m > 3$

Table 1. Capitulation types.

Therefore, one can easily deduce the following remark.



**Theorem 3.1.** *Keep the above notations.*

- (1) *Assume  $h_2(F_1) = 4$ . If the 2-class group of  $F_2$  or  $F_3$  is cyclic, then the 2-class group of  $F_i$  is cyclic for all  $i = 1, 2, 3$ . Furthermore,  $G_k$  is quaternion. Otherwise,  $G_k$  is dihedral.*
- (2) *Assume now that  $h_2(F_1) > 4$ . Then*
- $\triangleright$   *$G_k$  is a quaternion group if and only if  $G_{F_2}$  and  $G_{F_3}$  are quaternion groups.*
  - $\triangleright$   *$G_k$  is a dihedral group if and only if  $G_{F_2}$  and  $G_{F_3}$  are dihedral groups.*
  - $\triangleright$   *$G_k$  is a semi-dihedral group if and only if one of the two groups  $G_{F_2}$  and  $G_{F_3}$  is quaternion and the other is dihedral.*

**Proof.** (1) If  $h_2(F_1) = 4$ , then  $|G_k| = 8$ . Thus Remark 2.2 gives the first item. (2) Let  $i = 2, 3$ . Since  $F_i^{(2)} = k^{(2)}$ , this implies that each  $G_{F_i}$  is a subgroup of index 2 in  $G_k$ . Thus the group theoretic properties given in Section 2 complete the proof.  $\square$

The results of the second item can be summarized in Table 2.

$ \ker(j_1) $	$ \ker(j_2) $	$G_{F_2}$	$ \ker(j_3) $	$G_{F_3}$	$G_k$
4	2	(2, 2)	2	(2, 2)	$D_3$
4	2	$D_{m-1}$	2	$D_{m-1}$	$D_m, m > 3$
2	2	$Q_{m-1}$	2	$Q_{m-1}$	$Q_m, m > 3$
2	2	$D_{m-1}$	2	$Q_{m-1}$	$S_m, m > 3$

Table 2. Capitulation types for the case  $h_2(F_1) > 4$ .

#### 4. APPLICATIONS

Let  $d = 2q_1q_2$ , where  $q_1 \equiv q_2 \equiv -1 \pmod{4}$  are two distinct positive prime integers such that

$$\left(\frac{2}{q_j}\right) = -\left(\frac{2}{q_k}\right) = \left(\frac{q_j}{q_k}\right) = -\left(\frac{q_k}{q_j}\right) = 1, \quad 1 \leq j \neq k \leq 2.$$

Let  $\mathbb{k} = \mathbb{Q}(\sqrt{d}, \sqrt{-1})$  be an imaginary bicyclic biquadratic number field, which is called, by Hilbert (see [11]), a special Dirichlet field, and denote by  $\mathbb{k}^{(1)}$  the Hilbert 2-class field of  $\mathbb{k}$  and  $\mathbb{k}^{(2)}$  the Hilbert 2-class field of  $\mathbb{k}^{(1)}$ . Put  $G_{\mathbb{k}} = \text{Gal}(\mathbb{k}^{(2)}/\mathbb{k})$ . By [1], the 2-class group of  $\mathbb{k}$  is of type (2, 2). In this section we will apply the results of the above sections to determine the structure of  $G_{\mathbb{k}}$ .

**4.1. Preliminary results.** Let us first collect some results that will be useful in what follows. Let  $k_j$ ,  $1 \leq j \leq 3$ , be the three real quadratic subfields of a biquadratic real number field  $K_0$  and  $\varepsilon_j > 1$  be the fundamental unit of  $k_j$ . Since the square of any unit of  $K_0$  is in the group generated by the  $\varepsilon_j$ 's,  $1 \leq j \leq 3$ , then to

determine a fundamental system of units of  $K_0$  it suffices to determine which of the units in  $B := \{\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_1\varepsilon_2, \varepsilon_1\varepsilon_3, \varepsilon_2\varepsilon_3, \varepsilon_1\varepsilon_2\varepsilon_3\}$  are squares in  $K_0$  (see [14]). Hence, by Dirichlet's unit theorem, a fundamental system of units of  $K_0$  consists of three positive units chosen among  $B' := B \cup \{\sqrt{\eta} : \eta \in B \text{ and } \sqrt{\eta} \in K_0\}$ . We need the two following lemmas.

**Lemma 4.1** ([5]). *Let  $d \equiv 1 \pmod{4}$  be a positive square free integer and  $\varepsilon_d = x + y\sqrt{d}$  be the fundamental unit of  $\mathbb{Q}(\sqrt{d})$ . Assume  $N(\varepsilon_d) = 1$ , then*

- (1)  $x + 1$  and  $x - 1$  are not squares in  $\mathbb{N}$ , i.e.,  $2\varepsilon_d$  is not a square in  $\mathbb{Q}(\sqrt{d})$ .
- (2) For every prime  $p$  dividing  $d$ ,  $p(x + 1)$  and  $p(x - 1)$  are not squares in  $\mathbb{N}$ .

In the following lemma, we state a refinement to Lemma 4.1 above.

**Lemma 4.2.** *Let  $d \equiv 1 \pmod{4}$  be a positive square free integer and  $\varepsilon_d = \frac{1}{2}(x + y\sqrt{d})$  the fundamental unit of  $\mathbb{Q}(\sqrt{d})$ . Assume  $N(\varepsilon_d) = 1$ .*

- (1) If  $d \equiv 1 \pmod{8}$ , then both  $x$  and  $y$  are even.
- (2) If  $d \equiv 5 \pmod{8}$ , then  $x$  and  $y$  can be either even or odd. Moreover, if  $x$  and  $y$  are odd, then  $x + 2$  and  $x - 2$  are not squares in  $\mathbb{N}$ .

*Proof.* (1) Assume  $d \equiv 1 \pmod{8}$ . As  $N(\varepsilon_d) = 1$ , then  $x^2 - 4 = y^2d$ , hence  $x^2 - 4 \equiv y^2 \pmod{8}$ . On the other hand, if we suppose that  $x$  and  $y$  are odd, then  $x^2 \equiv y^2 \equiv 1 \pmod{8}$ , but this implies the contradiction  $-3 \equiv 1 \pmod{8}$ . Thus  $x$  and  $y$  are even.

(2) Assume  $d \equiv 5 \pmod{8}$ . To prove the first assertion of (2), it suffices to give examples justifying the existence of the two cases. By the PARI/GP system we have:

$d$	$d \pmod{8}$	$N(\varepsilon_d)$	$x$	$y$
21	5	1	5	1
69	5	1	25	3
77	5	1	9	1
93	5	1	29	3
133	5	1	173	15
141	5	1	190	16
381	5	1	2030	104
781	5	1	135212398	4838280

For the second assertion, suppose that  $x \pm 2 = y_1^2$ ,  $x \mp 2 = dy_2^2$ , then

$$\varepsilon_d = \frac{x + y\sqrt{d}}{2} = \frac{1}{4}(y_2\sqrt{d} + y_1)^2.$$

This in turn implies that  $\sqrt{\varepsilon_d} \in \mathbb{Q}(\sqrt{d})$ , which is absurd. □

Now we state a lemma which is very useful for getting a FSU of a real biquadratic subfield or imaginary triquadratic subfield of  $\mathbb{Q}(\sqrt{2}, \sqrt{q_1}, \sqrt{q_2}, \sqrt{-1})$ .

**Lemma 4.3.** *Let  $q_1 \equiv 7 \pmod{8}$  and  $q_2 \equiv 3 \pmod{8}$  be two primes such that  $(q_2/q_1) = -1$ .*

- (1) *Let  $x$  and  $y$  be two integers or semi-integers such that  $\varepsilon_{q_1 q_2} = x + y\sqrt{q_1 q_2}$ , then*
- (1a)  $2q_1(x + 1)$  *is a square in  $\mathbb{N}$ ,*
  - (1b)  $\sqrt{\varepsilon_{q_1 q_2}} = y_1\sqrt{q_1} + y_2\sqrt{q_2}$  *and  $1 = q_1 y_1^2 - q_2 y_2^2$  for some integers or semi-integers  $y_1$  and  $y_2$  such that  $y = 2y_1 y_2$ .*
- (2) *Let  $a$  and  $b$  be two integers such that  $\varepsilon_{2q_1 q_2} = a + b\sqrt{2q_1 q_2}$ . Then we have*
- (2a)  $2q_1(a + 1)$  *is a square in  $\mathbb{N}$ ,*
  - (2b)  $\sqrt{2\varepsilon_{2q_1 q_2}} = b_1\sqrt{2q_1} + b_2\sqrt{q_2}$  *and  $2 = 2q_1 b_1^2 - q_2 b_2^2$  for some integers  $b_1$  and  $b_2$  such that  $b = b_1 b_2$ .*
- (3) *Let  $c$  and  $d$  be two integers such that  $\varepsilon_{2q_1} = c + d\sqrt{2q_1}$  and let  $\alpha$  and  $\beta$  be two integers such that  $\varepsilon_{q_1} = \alpha + \beta\sqrt{q_1}$ . Then we have*
- (3a)  $\sqrt{2\varepsilon_{q_1}} = \beta_1 + \beta_2\sqrt{q_1}$  *and  $2 = \beta_1^2 - q_1 \beta_2^2$  for some integers  $\beta_1$  and  $\beta_2$  such that  $\beta = \beta_1 \beta_2$ ,*
  - (3b)  $\sqrt{2\varepsilon_{2q_1}} = d_1 + d_2\sqrt{2q_1}$  *and  $2 = d_1^2 - 2q_1 d_2^2$  for some integers  $d_1$  and  $d_2$  such that  $d = d_1 d_2$ .*
- (4) *Let  $c$  and  $d$  be two integers such that  $\varepsilon_{2q_2} = c + d\sqrt{2q_2}$  and let  $\alpha$  and  $\beta$  be two integers such that  $\varepsilon_{q_2} = \alpha + \beta\sqrt{q_2}$ . Then we have*
- (4a)  $\sqrt{2\varepsilon_{q_2}} = \beta_1 + \beta_2\sqrt{q_2}$  *and  $2 = -\beta_1^2 + q_2 \beta_2^2$  for some integers  $\beta_1$  and  $\beta_2$  such that  $\beta = \beta_1 \beta_2$ ,*
  - (4b)  $\sqrt{2\varepsilon_{2q_2}} = d_1 + d_2\sqrt{2q_2}$  *and  $2 = -d_1^2 + 2q_2 d_2^2$  for some integers  $d_1$  and  $d_2$  such that  $d = d_1 d_2$ .*

*Proof.* Using Lemmas 4.1 and 4.2, we get the statements of this lemma, for more details see [6]. □

**4.2. Capitulation.** Let  $q_1 \equiv q_2 \equiv -1 \pmod{4}$  be primes. Without loss of generality, we can assume that  $q_1$  and  $q_2$  satisfy the conditions

$$\left(\frac{2}{q_1}\right) = -\left(\frac{2}{q_2}\right) = \left(\frac{q_1}{q_2}\right) = -\left(\frac{q_2}{q_1}\right) = 1.$$

Then, by [1], the 2-class group of  $\mathbb{k}$  is of type  $(2, 2)$ , so denote by  $\mathbb{K}_1 = \mathbb{k}(\sqrt{q_1}) = \mathbb{Q}(\sqrt{q_1}, \sqrt{2q_2}, i)$ ,  $\mathbb{K}_2 = \mathbb{k}(\sqrt{q_2}) = \mathbb{Q}(\sqrt{q_2}, \sqrt{2q_1}, i)$  and  $\mathbb{K}_3 = \mathbb{k}(\sqrt{2}) = \mathbb{Q}(\sqrt{2}, \sqrt{q_1 q_2}, i)$  the three unramified quadratic extensions, within  $\mathbb{k}_1^{(1)}$ , of  $\mathbb{k}$ .

Now, we correct the error made in the article [4]. The fundamental systems of units given in [4], Proposition 3.1, for  $\mathbb{K}_1^+$  and  $\mathbb{K}_1$  are not correct. In fact, the error was committed in the FSU of  $\mathbb{K}_1^+$ , this affected that of  $\mathbb{K}_1$ , and thus the main theorem.



**Proposition 4.4.** *Let  $q_1$  and  $q_2$  be two primes defined as above. Then*

- (1) *A FSU of  $\mathbb{K}_1^+$  is  $\{\varepsilon_{q_1}, \sqrt{\varepsilon_{2q_1q_2}}, \sqrt{\varepsilon_{q_1\varepsilon_{2q_2}}}\}$  and that of  $\mathbb{K}_1$  is  $\{\sqrt{\varepsilon_{2q_1q_2}}, \sqrt{\varepsilon_{q_1\varepsilon_{2q_2}}, \sqrt{\varepsilon_{2q_2}}}\}$ .*
- (2) *A FSU of  $\mathbb{K}_2^+$  is  $\{\varepsilon_{2q_1q_2}, \sqrt{\varepsilon_{q_2\varepsilon_{2q_1q_2}}, \sqrt{\varepsilon_{2q_1\varepsilon_{2q_1q_2}}}\}$  and that of  $\mathbb{K}_2$  is  $\{\sqrt{\varepsilon_{q_2\varepsilon_{2q_1q_2}}, \sqrt{\varepsilon_{2q_1\varepsilon_{2q_1q_2}}, \sqrt{\varepsilon_{2q_1\varepsilon_{2q_1q_2}}}\}$ .*
- (3) *A FSU of both  $\mathbb{K}_3^+$  and  $\mathbb{K}_3$  is  $\{\varepsilon_2, \varepsilon_{2q_1q_2}, \sqrt{\varepsilon_{q_1q_2\varepsilon_{2q_1q_2}}}\}$ .*

*Proof.* Using Lemma 4.3 and the method described in the beginning of this subsection (page 6), we easily deduce the result for  $\mathbb{K}_i^+$ ,  $i = 1, 2, 3$  (we proceed as in the proof of [4], Proposition 3.1). Again Lemma 4.3 and [2], Proposition 2, give the result for  $\mathbb{K}_i$ ,  $i = 1, 2, 3$ .  $\square$

Denote by  $\kappa_{\mathbb{K}_j}$  the set of classes of  $\mathbb{k}$  capitulating in  $\mathbb{K}_j$ . Then proceeding as in the proof of [4], Theorem 3.3, we get the following result.

**Theorem 4.5.** *Let  $\mathbb{K}_j$ ,  $1 \leq j \leq 3$ , be the three unramified quadratic extensions of  $\mathbb{k}$  defined above. Then  $|\kappa_{\mathbb{K}_1}| = |\kappa_{\mathbb{K}_2}| = |\kappa_{\mathbb{K}_3}| = 2$ .*

**Lemma 4.6.** *Keep the above notations and conditions satisfied by  $q_1$  and  $q_2$ . Then, the 2-class group of  $\mathbb{K}_2$  is cyclic and those of  $\mathbb{K}_1$  and  $\mathbb{K}_3$  are of type  $(2, 2)$ .*

*Proof.* Let us compute the class number of  $\mathbb{K}_2$ . For the values of class numbers of quadratic fields, see [8], [12]. Proposition 4.4 implies that  $q(\mathbb{K}_2) = 8$ , so by the class number formula (cf. [14]) we obtain

$$\begin{aligned} h_2(\mathbb{K}_2) &= \frac{1}{2^5} q(\mathbb{K}_2) h_2(-1) h_2(q_2) h_2(-q_2) h_2(2q_1) h_2(-2q_1) h_2(2q_1q_2) h_2(-2q_1q_2) \\ &= \frac{1}{2^5} \cdot 8 \cdot h_2(-2q_1) \cdot 2 \cdot 4 \\ &= 2h_2(-2q_1). \end{aligned}$$

Since, by [8], Corollaries (19.6) and (18.4),  $h_2(-2q_1)$  is divisible by 4, so  $h_2(\mathbb{K}_2)$  is divisible by 8. Therefore, the 2-class group of  $\mathbb{K}_2$  cannot be of type  $(2, 2)$ . It follows that the 2-class group of  $\mathbb{K}_2$  is cyclic and those of  $\mathbb{K}_1$  and  $\mathbb{K}_3$  are of type  $(2, 2)$ .  $\square$

**Lemma 4.7** ([6]). *Keep the above hypothesis. The Hilbert 2-class field of  $\mathbb{k}$  is  $\mathbb{k}^{(1)} = \mathbb{Q}(\sqrt{2}, \sqrt{q_1}, \sqrt{q_2}, \sqrt{-1})$  and we have*

$$\begin{aligned} E_{\mathbb{k}^{(1)}} &= \left\langle \zeta_8, \varepsilon_2, \sqrt{\varepsilon_{2q_2}}, \sqrt{\varepsilon_{q_1q_2}}, \sqrt{\varepsilon_{2q_1q_2}}, \sqrt[4]{\varepsilon_{q_1\varepsilon_{q_2\varepsilon_{2q_2}\varepsilon_{q_1q_2\varepsilon_{2q_1q_2}}}}, \right. \\ &\quad \left. \sqrt[4]{\varepsilon_{2^2\varepsilon_{2q_1\varepsilon_{q_1q_2\varepsilon_{2q_1q_2}}}}, \sqrt[4]{\zeta_8^2\varepsilon_{q_1\varepsilon_{q_1q_2\varepsilon_{2q_1q_2}}}} \right\rangle. \end{aligned}$$

**Lemma 4.8.** *Keep the above hypothesis. We have:*

$$\begin{aligned}
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\varepsilon_2) &= -1, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{2q_2}}) &= -\varepsilon_{2q_2}, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{q_1q_2}}) &= 1, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{2q_1q_2}}) &= \varepsilon_{2q_1q_2}, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\zeta_8) &= -i, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt[4]{\varepsilon_{q_1}\varepsilon_{q_2}\varepsilon_{2q_2}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) &= \pm\sqrt{\varepsilon_{q_1}\varepsilon_{2q_2}}\sqrt{\varepsilon_{2q_1q_2}}, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt[4]{\varepsilon_2^2\varepsilon_{2q_1}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) &= \pm\sqrt{\varepsilon_{2q_1q_2}}, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt[4]{\zeta_8^2\varepsilon_2^2\varepsilon_{q_1}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) &= \pm i\sqrt{i\varepsilon_{q_1}}\sqrt{\varepsilon_{2q_1q_2}}.
\end{aligned}$$

If  $q_2 = 3$ , we have  $N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\zeta_6) = 1$ .

**Proof.** Assume that  $q_1 \equiv 7 \pmod{8}$ ,  $q_2 \equiv 3 \pmod{8}$  and  $(q_2/q_1) = -1$ . By the relations given in Lemma 4.3, we have

$$\begin{aligned}
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\varepsilon_2) &= (1 + \sqrt{2})(1 - \sqrt{2}) = -1, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{2q_2}}) &= \frac{1}{\sqrt{2}}(d_1 + d_2\sqrt{2q_2})\frac{1}{-\sqrt{2}}(d_1 + d_2\sqrt{2q_2}) = -\varepsilon_{2q_2}, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{q_1q_2}}) &= (y_1\sqrt{q_1} + y_2\sqrt{q_2})(y_1\sqrt{q_1} - y_2\sqrt{q_2}) = y_1^2q_1 - y_2^2q_2 = 1, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{2q_1q_2}}) &= \frac{1}{\sqrt{2}}(b_1\sqrt{2q_1} + b_2\sqrt{q_2})\frac{1}{-\sqrt{2}}(-b_1\sqrt{2q_1} - b_2\sqrt{q_2}) = \varepsilon_{2q_1q_2}, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{q_1}}) &= \frac{1}{\sqrt{2}}(\beta_1 + \beta_2\sqrt{q_1})\frac{1}{-\sqrt{2}}(\beta_1 + \beta_2\sqrt{q_1}) = -\varepsilon_{q_1}, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{2q_1}}) &= \frac{1}{\sqrt{2}}(d_1 + d_2\sqrt{2q_1})\frac{1}{-\sqrt{2}}(d_1 - d_2\sqrt{2q_2}) = -1, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{q_2}}) &= \frac{1}{\sqrt{2}}(\beta_1 + \beta_2\sqrt{q_2})\frac{1}{-\sqrt{2}}(\beta_1 - \beta_2\sqrt{q_2}) = 1, \\
N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\zeta_8) &= N_{\mathbb{k}^{(1)}/\mathbb{K}_1}\left(\frac{1+i}{-\sqrt{2}}\right) = -\zeta_8^2 = -i.
\end{aligned}$$

So we have

$$N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt{\varepsilon_{q_1}\varepsilon_{q_2}\varepsilon_{2q_2}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) = (-\varepsilon_{q_1}) \cdot 1 \cdot (-\varepsilon_{2q_2}) \cdot 1 \cdot \varepsilon_{2q_1q_2} = \varepsilon_{q_1}\varepsilon_{2q_2}\varepsilon_{2q_1q_2}.$$

Then  $N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(\sqrt[4]{\varepsilon_{q_1}\varepsilon_{q_2}\varepsilon_{2q_2}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) = \pm\sqrt{\varepsilon_{q_1}\varepsilon_{2q_2}}\sqrt{\varepsilon_{2q_1q_2}}$ . We similarly get the rest.  $\square$

**Lemma 4.9.** *Keep the above hypothesis. We have:*

$$\begin{aligned}
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\varepsilon_2) &= \varepsilon_2^2, \\
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\sqrt{\varepsilon_{2q_2}}) &= -1, \\
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\sqrt{\varepsilon_{q_1q_2}}) &= -\varepsilon_{q_1q_2}, \\
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\sqrt{\varepsilon_{2q_1q_2}}) &= -\varepsilon_{2q_1q_2}, \\
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\zeta_8) &= i, \\
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\sqrt[4]{\varepsilon_{q_1}\varepsilon_{q_2}\varepsilon_{2q_2}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) &= \pm\sqrt{\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}, \\
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\sqrt[4]{\varepsilon_2^2\varepsilon_{2q_1}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) &= \pm\varepsilon_2\sqrt{\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}, \\
 N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\sqrt[4]{\zeta_8^2\varepsilon_2^2\varepsilon_{q_1}\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}) &= \pm\zeta_8\varepsilon_2\sqrt{\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}}.
 \end{aligned}$$

If  $q_2 = 3$ , we have  $N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(\zeta_6) = 1$ .

*Proof.* The proof is similar to that of Lemma 4.8. □

From the two above lemmas, Proposition 4.4 and [10] we have:

**Corollary 4.10.** *Keep the above hypothesis. We have:*

(1) *The number of classes of  $\mathbb{K}_1$  which capitulate in  $\mathbb{k}^{(1)}$  is*

$$[\mathbb{k}^{(1)} : \mathbb{K}_1][E_{\mathbb{K}_1} : N_{\mathbb{k}^{(1)}/\mathbb{K}_1}(E_{\mathbb{k}^{(1)}})] = 2 \cdot 1 = 2.$$

(2) *The number of classes of  $\mathbb{K}_3$  which capitulate in  $\mathbb{k}^{(1)}$  is*

$$[\mathbb{k}^{(1)} : \mathbb{K}_3][E_{\mathbb{K}_3} : N_{\mathbb{k}^{(1)}/\mathbb{K}_3}(E_{\mathbb{k}^{(1)}})] = 2 \cdot 1 = 2.$$

**4.3. Main theorem.** We can now state the main result of this section.

**Theorem 4.11.** *Let  $q_1 \equiv q_2 \equiv -1 \pmod{4}$  be two distinct prime integers such that*

$$\left(\frac{2}{q_j}\right) = -\left(\frac{2}{q_k}\right) = \left(\frac{q_j}{q_k}\right) = -\left(\frac{q_k}{q_j}\right) = 1,$$

$1 \leq j \neq k \leq 2$ . Put  $\mathbb{k} = \mathbb{Q}(\sqrt{2q_1q_2}, i)$ . Note that  $\mathbb{K}_j = \mathbb{k}(\sqrt{q_j}) = \mathbb{Q}(\sqrt{q_j}, \sqrt{2q_k}, i)$ ,  $\mathbb{K}_k = \mathbb{k}(\sqrt{q_k}) = \mathbb{Q}(\sqrt{q_k}, \sqrt{2q_j}, i)$  and  $\mathbb{K}_3 = \mathbb{k}(\sqrt{2}) = \mathbb{Q}(\sqrt{2}, \sqrt{q_1q_2}, i)$  are three unramified quadratic extensions of  $\mathbb{k}$ . Let  $m \geq 2$  such that  $2^m = h_2(-2q_j)$ . Then the 2-class field tower of  $\mathbb{k}$  stops at  $\mathbb{k}^{(2)}$  with  $\mathbb{k}^{(1)} \neq \mathbb{k}^{(2)}$  and

$$G_{\mathbb{K}_j} \simeq G_{\mathbb{K}_3} \simeq Q_{m+1}, \quad G_{\mathbb{k}} \simeq Q_{m+2} \quad \text{and} \quad G_{\mathbb{K}_k} \simeq \mathbb{Z}/2^{m+1}\mathbb{Z}.$$

**Proof.** Recall that  $G_{\mathbb{k}} = \text{Gal}(\mathbb{k}^{(2)}/\mathbb{k})$ , where  $\mathbb{k}^{(2)}$  is the second Hilbert 2-class field of  $\mathbb{k}$ . Without loss of generality, we may suppose that the primes  $q_1$  and  $q_2$  satisfy

$$q_1 \equiv q_2 \equiv -1 \pmod{4} \quad \text{and} \quad \left(\frac{2}{q_1}\right) = -\left(\frac{2}{q_2}\right) = \left(\frac{q_1}{q_2}\right) = -\left(\frac{q_2}{q_1}\right) = 1.$$

As the 2-class group  $\mathbb{C}\mathbb{L}_2(\mathbb{k})$  of  $\mathbb{k}$  is of type  $(2, 2)$ , then  $G_{\mathbb{k}}/G'_{\mathbb{k}} \simeq (2, 2)$ . On the other hand, by Theorem 4.5, there are exactly two classes of  $\mathbb{C}\mathbb{L}_2(\mathbb{k})$  which capitulate in each extension  $\mathbb{K}_j$ ,  $1 \leq j \leq 3$ , so Theorem 2.1 implies that  $G_{\mathbb{k}}$  is quaternion or semidihedral and the class field tower of  $\mathbb{k}$  stops at  $\mathbb{k}^{(2)}$  with  $\mathbb{k}^{(1)} \neq \mathbb{k}^{(2)}$ ; and thus, again by Theorem 2.1, one of the three quadratic extensions of  $\mathbb{k}$  has cyclic 2-class group and the two others have 2-class groups of type  $(2, 2)$  which is already proved in Lemma 4.6. The 2-class groups of  $\mathbb{K}_1$  and  $\mathbb{K}_3$  are of type  $(2, 2)$ . They are both sub-extensions of  $\mathbb{k}^{(1)}$  which has a cyclic 2-class group (since  $G'_{\mathbb{k}} \simeq \text{Gal}(\mathbb{k}^{(2)}/\mathbb{k}^{(1)}) \simeq \mathbb{C}\mathbb{L}_2(\mathbb{k}^{(1)})$  is a cyclic group), and there are exactly two classes in  $\mathbb{K}_1$  and  $\mathbb{K}_3$  capitulating in  $\mathbb{k}^{(1)}$  (Corollary 4.10), so neither  $G_{\mathbb{K}_1}$  nor  $G_{\mathbb{K}_3}$  is dihedral. Hence the result comes by Table 2.  $\square$

**Remark 4.12.** At the first step in the above proof, we showed that  $G_{\mathbb{k}}$  is quaternion or semidihedral. Note that it is impossible to decide whether  $G_{\mathbb{k}}$  is quaternion or semidihedral by using the usual method given by Kisilevsky (by determining whether  $\mathbb{K}_i/\mathbb{k}$  is of type A or B,  $i = 1, 3$ ). In fact, it is hard to determine the generators of the 2-class groups. For this reason the authors of [3] couldn't decide whether  $G_k$  is quaternion or semidihedral with  $k = \mathbb{Q}(\sqrt{-2}, \sqrt{pq})$  for two primes  $p \equiv 5 \pmod{8}$  and  $q \equiv 7 \pmod{8}$  (see [3], Corollary 17). Using the same techniques described in the general context in Section 3, we gave the answer in [7], Remark 5.7.

**Acknowledgement.** We are very grateful to the unknown referee for his/her several helpful suggestions, that enabled us to improve our paper, and for calling our attention to some missing details.

### References

- [1] *A. Azizi*: Sur le 2-groupe de classes d'idéaux de  $\mathbb{Q}(\sqrt{d}, i)$ . *Rend. Circ. Mat. Palermo, II. Ser.* 48 (1999), 71–92. (In French.) zbl MR doi
- [2] *A. Azizi*: Unités de certains corps de nombres imaginaires et abéliens sur  $\mathbb{Q}$ . *Ann. Sci. Math. Qué.* 23 (1999), 15–21. (In French.) zbl MR
- [3] *A. Azizi, I. Benhamza*: Sur la capitulation des 2-classes d'idéaux de  $\mathbb{Q}(\sqrt{d}, \sqrt{-2})$ . *Ann. Sci. Math. Qué.* 29 (2005), 1–20. (In French.) zbl MR
- [4] *A. Azizi, A. Zekhnini*: The structure of the second 2-class group of some special Dirichlet fields. *Asian-Eur. J. Math.* 13 (2020), Article ID 2050053, 6 pages. zbl MR doi
- [5] *A. Azizi, A. Zekhnini, M. Taous*: The generators of the 2-class group of some field  $\mathbb{Q}(\sqrt{pq_1q_2}, i)$ : Correction to Theorem 3 of [5]. *Int. J. Pure Appl. Math.* 103 (2015), 99–107. doi

- [6] *M. M. Chems-Eddin, A. Azizi, A. Zekhnini*: Unit groups and Iwasawa lambda invariants of some multiquadratic number fields. *Bol. Soc. Mat. Mex., III. Ser.* 27 (2021), Article ID 24, 16 pages. [zbl](#) [MR](#) [doi](#)
- [7] *M. M. Chems-Eddin, A. Zekhnini, A. Azizi*: Units and 2-class field towers of some multiquadratic number fields. *Turk. J. Math.* 44 (2020), 1466–1483. [zbl](#) [MR](#) [doi](#)
- [8] *P. E. Conner, J. Hurrelbrink*: *Class Number Parity*. Series in Pure Mathematics 8. World Scientific, Singapore, 1988. [zbl](#) [MR](#) [doi](#)
- [9] *D. Gorenstein*: *Finite Groups*. Harper’s Series in Modern Mathematics. Harper and Row, New York, 1968. [zbl](#) [MR](#)
- [10] *F.-P. Heider, B. Schmithals*: Zur Kapitulation der Idealklassen in unverzweigten primzyklischen Erweiterungen. *J. Reine Angew. Math.* 366 (1982), 1–25. (In German.) [zbl](#) [MR](#) [doi](#)
- [11] *D. Hilbert*: Über den Dirichlet’schen biquadratischen Zahlkörper. *Math. Ann.* 45 (1894), 309–340. (In German.) [zbl](#) [MR](#) [doi](#)
- [12] *P. Kaplan*: Sur le 2-groupe des classes d’idéaux des corps quadratiques. *J. Reine Angew. Math.* 283/284 (1976), 313–363. (In French.) [zbl](#) [MR](#) [doi](#)
- [13] *H. Kisilevsky*: Number fields with class number congruent to 4 mod 8 and Hilbert’s Theorem 94. *J. Number Theory* 8 (1976), 271–279. [zbl](#) [MR](#) [doi](#)
- [14] *H. Wada*: On the class number and the unit group of certain algebraic number fields. *J. Fac. Sci., Univ. Tokyo, Sect. I* 13 (1966), 201–209. [zbl](#) [MR](#)

*Authors’ addresses:* *Abdelmalek Azizi*, Department of Mathematics, Faculty of Sciences, Mohammed First University, Oujda, Morocco, e-mail: [abdelmalekazizi@yahoo.fr](mailto:abdelmalekazizi@yahoo.fr); *Mohamed Mahmoud Chems-Eddin*, Department of Mathematics, Faculty of Sciences Dhar El Mahraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco, e-mail: [2m.chemseddin@gmail.com](mailto:2m.chemseddin@gmail.com); *Abdelkader Zekhnini*, Department of Mathematics, Faculty of Sciences, Mohammed First University, Oujda, Morocco, e-mail: [zekha1@yahoo.fr](mailto:zekha1@yahoo.fr).