

P. Houston; L. O'Raiheartaigh

On the charge distribution of static axial and mirror symmetric monopole systems

In: Zdeněk Frolík (ed.): Abstracta. 8th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1980. pp. 96--103.

Persistent URL: <http://dml.cz/dmlcz/701186>

## Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1980

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On the Charge Distribution of Static Axial and  
Mirror Symmetric Monopole Systems

P. Houston and L. O'Raiheartaigh

It has been shown that the imposition of axial and mirror symmetry on the static magnetic SU(2) Yang-Mills-Higgs system reduces the usual first order ("self-dual") Bogomolny equations to five equations for five unknown (gauge-invariant) functions. In this note we show that, for a differentiable Higgs field, the five field equations and the topology rule out all charge-distributions along the axis of symmetry except a single-point distribution, and the boundary conditions at the single point distribution are discussed.

Ever since the discovery [1] that the static Yang-Mills-Higgs equations of unified gauge-theory admit finite-energy magnetic monopole solutions, there has been interest in the question of solutions which go beyond the original ones, which are all spherically symmetric. The next simplest configuration seems to be that in which the fields are axially symmetric and symmetric under reflexions in all planes containing the z-axis, and in this note we wish to consider the field equations for such systems.

To obtain the simplest possible system of this kind we do not consider the full second-order YMH equations but rather the first-order equations

$$(1) \quad \underline{F} = D \underline{\Phi} ,$$

which are obtained by minimizing the Bogomolny-Prasad-Sommerfield [2] Hamiltonian

$$(2) \quad H = \frac{1}{2} \int d_3x (\vec{F}^2 + (D_\mu \Phi)^2),$$

in which the gauge-group is SU(2), the Higgs field  $\Phi$  is assumed to belong to the adjoint representation, and the scalar potential is set equal to zero and replaced by the boundary condition  $(\Phi, \Phi) \rightarrow c^2$  as  $r \rightarrow \infty$ .

It has been shown [3,4] that for axial and mirror symmetry the field equations (1) reduce to the five equations

$$(3) \quad ku_\alpha = b_{,\alpha} + \rho \epsilon_{\alpha\beta} h_{,\beta}$$

$$(4) \quad -k_{,\alpha} = bu_\alpha + \rho h \epsilon_{\alpha\beta} u_\beta$$

$$(5) \quad hk = \rho \epsilon_{\alpha\beta} u_{\alpha\beta}$$

for the five (gauge-invariant) unknown functions  $h, k, b, u_\alpha$  defined as

$$h^2 = (\Phi, \Phi), \quad h^2 k^2 = (D_\gamma \Phi, D_\gamma \Phi),$$

$$(6) \quad h^2 k^2 (b^2 + k^2) = (D_\gamma^2 \Phi, D_\gamma^2 \Phi), \quad h_{,\alpha} h_{,\beta} + h^2 u_\alpha u_\beta = (D_\alpha \Phi, D_\beta \Phi).$$

It was also shown that the first four equations are equivalent to the existence of a master-potential  $W$  such that

$$(7) \quad \frac{\partial W}{\partial z} = hb, \quad h^2 = c^2 + \Delta W,$$

$$2\rho \frac{\partial W}{\partial \rho} = \rho^2 (h^2 - c^2) + n^2 - k^2 - b^2.$$

In particular

$$(8) \quad (k^2 + b^2)_{,\alpha} = \rho^2 (h^2)_{,\alpha} - 2\rho \epsilon_{\alpha\beta} (bh)_{,\beta}$$

Furthermore, the ordinary Maxwell magnetic field  $\vec{B}$ , defined as usual [5], to be

$$(9) \quad B_i = (F_i, \theta) - \frac{1}{2} \epsilon_{ijk} (\theta, D_j \theta \wedge D_k \theta), \quad \text{where } \theta = \Phi/h,$$

turns out to be just ..

$$(10) \quad B_\alpha = \frac{1}{\rho} \epsilon_{\alpha\beta} b_{,\beta} \quad , \quad (B_\gamma = 0) \quad .$$

Thus the magnetic charge contained in any volume of revolution  $V$  around the  $z$ -axis and intersecting the axis at two points  $z_1$  and  $z_2$  is just

$$(11) \quad \Delta n = \frac{1}{4\pi} \int d_3x \nabla_\alpha B_\alpha = \frac{1}{2} (b(z_1) - b(z_2)) \quad .$$

Of course, since (8) is a curl the charge can be located only at points where is not well defined i.e. where the Higgs field  $\Phi$  is zero. The boundary conditions for the functions  $h, b, k, u_\alpha$  as  $r \rightarrow 0$  are

$$(12) \quad k, u_\alpha \rightarrow 0 \quad , \quad h \rightarrow c - \frac{n}{r} \quad , \quad b \rightarrow -n \cos \theta \quad ,$$

where  $n$  is the total topological charge (in agreement with (9)). The boundary conditions on the  $z$ -axis are determined by the finiteness of the energy [2], which in terms of the variables  $h, k, b, u_\alpha$  takes the form

$$(13) \quad H = \frac{1}{2} \int d_3x \left[ (\nabla_\alpha h)^2 + h^2 u_\alpha^2 + (\epsilon_{\alpha\beta} u_{\alpha,\beta})^2 + \frac{1}{\rho^2} \left\{ h^2 k^2 + (b_{,\alpha} - k u_\alpha)^2 + (k_{,\alpha} + b u_\alpha)^2 \right\} \right] \quad .$$

From this equation we see that (subject to uniform behaviour with respect to  $z$ ) we have

$$(14) \quad \rho^{3/2} \nabla_\alpha h \rightarrow 0 \quad , \quad u_\alpha \rightarrow b^{-1} k_{,\alpha} \rightarrow -k^{-1} b_{,\alpha} \quad , \quad hk \rightarrow 0 \quad ,$$

as  $\rho \rightarrow 0$  .

From the central equation in (14) and from (10) we have in particular — —

$$(15) \quad b^2 + k^2 = n^2 \quad \text{for} \quad \rho = 0 \quad .$$

This equation also follows from (3,4,5) and the first eqn. in (14).

In a previous note [6] it was shown that topological considerations alone implied that for an axially symmetric system

with charges located on the z-axis, the Higgs field  $h$  had to vanish everywhere along the axis joining these points, and it is interesting to note that this result can be verified directly from (14) and (15) in a gauge-invariant manner. For if  $h \neq 0$  on the axis, then from (14) we have  $k=0$  and hence from (15)  $b = \pm n$ . It then follows from (11) that between any two points where  $h \neq 0$  the charge must be either 0, or  $n$ . Since  $n$  is the total charge this establishes the result.

The fact that the Higgs field must vanish between charges suggests that actually there can be no such distribution, except at a single point, and we now wish to show that, provided that the Higgs field  $h$  has a continuous derivative such is indeed the case. Although this condition is quite mild, it is a little more than is required by finite energy.

The basis for our result will be the two equations

$$(16) \quad \Delta(hb) = (h^2)_{,z} ,$$

$$(17) \quad \Delta(hk) = \frac{2(ff_{,\alpha} - g\epsilon_{\alpha\beta} f_{,\beta})}{h^2 D^2} (hk)_{,\alpha} + \\ + \left[ \frac{D^2}{\rho^2} + \frac{2\epsilon_{\alpha\beta} g_{,\alpha} f_{,\beta}}{h^2 D^2} \right] (hk) ,$$

$$(18) \quad f \equiv \rho h^2 , \quad g \equiv hb , \quad D^2 = b^2 + k^2 + \rho^2 h^2 ,$$

which follow directly from (3) - (7). If we now assume that at any zero,  $\underline{z}_0$ , of the Higgs field on the z-axis,  $h$  is  $C^1$  then (17) has the small distance approximation

$$(19) \quad \Delta(hk) = \frac{n^2}{\rho^2} (hk) .$$

Since finite energy requires that  $hk \rightarrow 0$ , as  $\underline{x} \rightarrow \underline{z}_0$ , the solution of (19) is an associated Legendre function of the first kind. Thus wherever  $P_m^n(\cos \theta) \neq 0$  we have

$$(20) \quad hk \rightarrow Kr^m P_m^n(\cos \theta), \quad m \geq n,$$

where  $r, \theta$  are polar coordinates about  $\underline{z}_0$ . Since  $k$  is bounded, equation (20) already shows that  $h$  does not vanish infinitely fast at  $\underline{z}_0$ , but between the powers  $r$  and  $r^m$ .

Let us now consider those points  $\underline{z}_0$ , where  $h \rightarrow 0$ , but  $b \neq 0$  for small  $r$  and either  $\theta=0$  or  $\pi$  (or both). It follows from (14) and (15) that such points exist. Also from (15) we see that, as  $r \rightarrow 0$ ,

$$(21) \quad (hb)^2 + (hk)^2 \rightarrow n^2 k^2,$$

and that, since  $b \not\rightarrow 0$ ,  $hk$  can not dominate  $hb$ , for all  $\theta$ . From (21) we can write (16) in the form

$$(22) \quad \Delta(hb) = \frac{1}{n^2} [(h^2 b^2)_{,z} + (h^2 k^2)_{,z}] + \dots$$

Since  $hk$  does not dominate  $hb$  and  $hb \rightarrow 0$  as  $r \rightarrow 0$ , it follows from (22) that the leading behaviour of  $hk$  is a solution of the homogeneous equation  $\Delta(hb) = 0$ . In other words

$$(23) \quad hb \rightarrow K_s r^s P_s(\cos \theta) \quad \text{for some } s \leq m,$$

where  $K_s$  is a non-zero constant. It then follows from (21) that  $h^2$  is of the form

$$(24) \quad h^2 \rightarrow \frac{1}{n^2} [(K_s r^s P_s(\cos \theta))^2 + (Kr^m P_m^n(\cos \theta))^2], \quad s \leq m,$$

as  $r \rightarrow 0$ . Equation (24) shows that  $h^2$  is a regular function at  $\underline{z}_0$  and that

$$(25) \quad h^2(r, 0) = h^2(r, \pi), \quad r \neq 0.$$

Thus the field equations require that  $h^2$  be symmetric about any point on the  $z$ -axis where it vanishes. It is indeed possible for a single point charge distribution to satisfy this. However, now consider a charge which is not distributed at a single point on the  $z$ -axis. According to the previous discussion of the topology,  $h^2$ , in this case, must vanish along a

finite section of the z-axis. On the z-axis in a neighbourhood of an end point  $z_0$  of such a section,  $h^2=0$  inside  $z_0$ , and  $h^2 \neq 0$ ,  $b^2 = n^2 \neq 0$  outside. Thus  $h^2$  can not be symmetric about  $z_0$ . Therefore equation (25) rules out all but the single point charge distribution.

Let us now assume that the charge distribution is concentrated at a single point (superimposed monopoles), which we take to be the origin. The equations derived above for  $z_0$  are still valid, so that near the origin we have the leading behaviour

$$\begin{aligned}
 (26) \quad & hk \rightarrow Kr^m P_m^n(\cos \theta), \quad m \geq n, \\
 & hb \rightarrow K_s r^s P_s(\cos \theta), \quad m \geq s \geq 1, \\
 & h \rightarrow \frac{1}{n} r^s |K_s P_s(\cos \theta)|, \quad \text{for } s < m, \\
 & h \rightarrow \frac{1}{n} r^m \left\{ (K_m P_m(\cos \theta))^2 + (K P_m^n(\cos \theta))^2 \right\}^{1/2} \\
 & \qquad \qquad \qquad \text{for } s = m,
 \end{aligned}$$

where the condition  $s \leq m$  comes from the fact that if  $s > m$ , we have  $b \rightarrow 0$  and hence no topological charge. The most natural case (and possibly only stable case) is  $m=n$ , and then we have

$$\begin{aligned}
 (27) \quad & hk \rightarrow K \rho^n, \\
 & hb \rightarrow K_s r^s P_s(\cos \theta), \\
 & h \rightarrow \frac{1}{n} r^s |K_s P_s(\cos \theta)|, \quad \text{for } s < m, \\
 & h \rightarrow \frac{1}{n} r^n \left\{ (K_n P_n(\cos \theta))^2 + K^2 \rho^{2n} \right\}^{1/2}, \\
 & \qquad \qquad \qquad \text{for } s = m.
 \end{aligned}$$

It has been pointed out by P. Rossi that when  $s=1$  the behaviour at the origin can be determined more precisely, namely as

$$(28) \quad h^2 = \left( \sum_{l=1}^n K_l \frac{r^1}{n} P_l(\cos \theta) \right)^2,$$

$$b^2 = (n + \sum_{l=1}^n \frac{\sin^2 \theta r^{l+1}}{n(l+1)} K_1 P_1'(\cos \theta))^2 .$$

This result can be understood in our context, and be slightly generalized, by noting that if  $s < m$  then  $k \sim r^{m-s}$  and hence from the field equations (3) - (5)

$$(29) \quad b_{,\alpha} + \rho \in_{\alpha\beta} h_{,\beta} = O(r^{2m-2s-1}) ,$$

which implies that, to the order shown, there exists a harmonic function

$$(30) \quad U = \sum_{l=1}^{n+1} \gamma_l r^l P_l(\cos \theta) ,$$

such that

$$(31) \quad h^2 = \left(\frac{\partial U}{\partial z}\right)^2 , \quad \text{and} \quad b^2 = \rho^2 \left(\frac{\partial U}{\partial \rho}\right)^2 .$$

Alternatively, we may note that to the order in which  $k$  (and therefore  $ru_\alpha$ ) may be neglected the field equations (7) reduce to the two equations  $\Delta h = 0$  and (16) and that these two equations are identically satisfied by (31) if  $U$  is harmonic.

A further symmetry which is carried by the field equations (3) - (5) is invariance under reflexion in the  $(xy)$ -plane, where  $h, k$  are even and  $b$  is odd with respect to this reflexion (in agreement with the boundary conditions as  $r \rightarrow \infty$ ). If we impose this condition on the fields then all the integers,  $m, s, n$  above must be odd. In particular, only odd topological charges  $n$  are permitted. Since this symmetry agrees naturally with the equation and boundary conditions, it suggests that even topological charges might not be allowed [7] (under the assumptions of axial symmetry).

Finally we should mention that the results mentioned here are only very slightly modified when the assumption of mirror



symmetry is removed. Indeed, apart from the complication due to the introduction of four new fields, the results go through for axial symmetry alone. This generalization and a more detailed treatment of the above results, will be given in a forthcoming paper.

#### References

- [1] G. 't Hooft, Nucl. Phys. B79, 276 (1976)  
A. Polyakov, JETP letters, 20, 194 (1974)
- [2] E. Bogomolny, Sov. J. Nucl. Phys. 24, 861 (1976)
- [3] P. Houston, L. O'Raifeartaigh, Proc. Seminar on Group Th. and Quant. Fields, Zvenigorod, Moscow, Nov. 1979. DIAS-STP-79-35  
N. Manton, Nucl. Phys. B135, 319 (1978)
- [4] S. Adler, Private Communication
- [5] Arafune, P. Freund, C. Goebel, J. Math. Phys. 16, 433 (1975)
- [6] P. Houston, L. O'Raifeartaigh, DIAS-STP-80-14 (1980)  
See also P.S. Jang, S.Y. Park, K.C. Wali, Phys. Rev. D17, 1641 (1978)
- [7] See also J. Madore, Gen. Rel. and Grav. 8, 655 (1977)